

**THE UNIVERSITY OF WESTERN ONTARIO
BIOLOGICAL AGENTS REGISTRY FORM**
Approved Biohazards Subcommittee: April 9, 2010
Biosafety Website: www.uwo.ca/humanresources/biosafety/

This form must be completed by each Principal Investigator who is in charge of a laboratory/facility where the work described in the laboratory or animal work proposed. The form is required for all work involving animals carrying zoonotic agents infectious to humans as defined by the Public Health Agency of Canada (PHAC) or Canadian Food Inspection Agency (CFIA).

Statements in yellow
are new/modified.
(for sept meeting)

This form must be updated at least every 3 years or when there are changes to the biohazards being used.

Containment Levels will be established in accordance with Laboratory Biosafety Guidelines, 3rd edition, 2004, Public Health Agency of Canada (PHAC) or Containment Standards for Veterinary Facilities, 1st edition 1996, Canadian Food Inspection Agency (CFIA).

Completed forms are to be returned to Occupational Health and Safety, (OHS), (Support Services Building, Room 4190) for distribution to the Biohazard Subcommittee. For questions regarding this form, please contact the Biosafety Officer at extension 81135 or biosafety@uwo.ca. If there are changes to the information on this form (excluding grant title and funding agencies), contact Occupational Health and Safety for a modification form. See website: www.uwo.ca/humanresources/biosafety

PRINCIPAL INVESTIGATOR	Peeyush K Lala
DEPARTMENT	Anatomy and Cell biology
ADDRESS	Medical sciences Bldg , Rm 433
PHONE NUMBER	519-661-3015 or Ext 83015
EMERGENCY PHONE NUMBER(S)	519-472-4501 (Dr Lala), 519-850-2071 (Xiping Xin)
EMAIL	pklala@uwo.ca

Location of experimental work to be carried out: Building(s) MSB Room(s) 433

*For work being performed at Institutions affiliated with the University of Western Ontario, the Safety Officer for the Institution where experiments will take place must sign the form prior to its being sent to the University of Western Ontario Biosafety Officer (See Section 12.0, Approvals).

FUNDING AGENCY/AGENCIES(1) CIHR, (2) Canadian Breast Cancer Foundation (CBCF)_(3) Upjohn Canada

GRANT TITLE(S): (1) Biology of the feto-maternal interface; (2) Regulation of lymphangiogenesis and lymphatic metastasis in a mouse breast cancer . model. (3) Natural host defence to neoplasia and its manipulations for immunotherapy

PLEASE ATTACH A BRIEF DESCRIPTION OF YOUR WORK THAT EXPLAINS THE BIOHAZARDS USED AND HOW THEY WILL BE STORED, USED AND DISPOSED OF. PROJECTS SUBMITTED WITHOUT A SUMMARY WILL NOT BE REVIEWED.

(1) The human placenta is a highly invasive "Tumor-like" structure in which a migratory and invasive cell population known as the "Extra-villous trophoblast (EVT)" cells invades the pregnant uterus and its arteries to establish an efficient exchange of oxygen and nutrients between the maternal and fetal circulations. However, unlike tumors, placental invasion of the uterus is exquisitely controlled to maintain a healthy utero-placental homeostasis. Inadequate invasiveness of EVT cells results in poor perfusion of the placenta with maternal blood, which in turn can cause intrauterine growth restriction (IUGR) of the fetus and a serious maternal disease called pre-eclampsia. Our long standing research efforts have been to identify molecular mechanisms regulating EVT cell invasiveness and their

derangements in preeclampsia and placenta cancer. For these studies we use occasional samples of human placenta, but more routinely human placenta-derived EVT cell lines produced by our lab many years ago (which we have been distributing world-wide to other researchers as well), and commercially available placental cancer (choriocarcinoma) cell lines. For example, we have discovered that two molecules , Transforming Growth factor (TGF) - β and a TGF- β binding proteoglycan decorin – both produced by the decidual tissue of the pregnant uterus negatively regulates EVT cell proliferation, migration and invasiveness in a mutually independent manner. We have found that our precancerous EVT cell lines as well as commercially available cancerous EVT (choriocarcinoma) cell lines (see list later) are resistant to the negative regulation by both molecules. We have discovered the genetic alterations responsible for TGF- β resistance in choriocarcinoma cells. Our current efforts using normal, precancerous and cancerous EVT cells (see list below) are to define the molecular mechanisms in decorin action and their alterations in trophoblast hyperinvasive disorders (precancer and cancer) as well as trophoblast hypoinvasive disorders such as pre-eclampsia.

(2) Breast cancer afflicts one in nine to ten women in Canada, and still accounts for the second highest cancer-related deaths in females, because of the difficulty in curing metastasis (spread to distant sites). This spread can occur to other organs via the new blood vessels recruited by the cancerous tissue to feed itself (a process called angiogenesis) or to lymph nodes via new lymphatic vessels draining the tumor site (a process called lymphangiogenesis). Spread to lymph nodes is often the earliest mode of spread of breast cancer. Our goals are to identify molecular mechanisms in blood-borne and lymphatic metastasis so that we can exploit this information to identify new therapeutic targets. To achieve our goals we use commercially available human breast cancer cell lines and also mouse breast cancer cell lines produced in our own laboratory from spontaneous mouse breast cancers. We conduct experiments in tissue culture and also in mice. We use immuno-deficient mice to transplant human breast cancer cell lines and syngeneic mice to transplant mouse breast cancer cell lines. We have discovered that expression of an enzyme Cyclo-oxygenase (COX)-2 promotes breast cancer progression and metastasis in mice and the human by activating prostaglandin-receptors on cancer cells. This is due to multiple mechanisms: inactivation of cancer fighting immune cells of the host, stimulation of cancer cell migration and cancer-associated angiogenesis and lymphangiogenesis. Using human breast cancer tissues and numerous human breast cancer cell lines varying in COX-2 expression (see list below) we discovered that COX-2 is responsible for upregulation of vascular Endothelial growth factors stimulating angiogenesis and lymphangiogenesis. Therapeutic value of drugs blocking COX-2 enzyme such as celecoxib and drugs blocking certain prostaglandin receptors are currently being successfully tested in our mouse breast cancer models bearing transplants of mouse or human breast cancer cell lines expressing COX-2. These cell lines are listed later. For example, murine C3L5 cells (transplanted in C3H/HeJ mice) and human MDA-MB-231 cells (transplanted in nude mice) express high levels of COX-2 and are highly metastatic in vivo.

(3) We had shown that prostaglandin E2 produced by cancer cells or host macrophages in the cancer bearing host inactivates cancer-fighting immune lymphocytes of the host. This could be reversed by therapeutic use of prostaglandin inhibitors blocking cyclo-oxygenase enzymes (such as indomethacin, ibuprofen or Celicoxib) to activate the lymphocytes. We have used these drugs in combination with an immune cell activating cytokine interleukin (IL)-2 to cure different types of advanced mouse cancers and human cancers grown in immuno-deficient mice. Subsequently we applied this combination therapy protocol to advanced human kidney cancer and melanoma patients in a phase to clinical trial at the London regional Cancer centre (1988-92) with highly promising results. This combination therapy is now practiced as one of the standard protocols in these types of cancer patients at the NCI , USA. Currently we are testing similar immunotherapy protocols in mice bearing mouse or human breast cancer transplants. These transplants include syngeneic mouse breast cancer cell lines such as C3L5 developed in our lab, as well as xenotransplants of COX-2 expressing (PGE-2 producing) human breast cancer lines such as MDA-MB-231 cells (listed later).

Names of all personnel working under Principal Investigators supervision in this location: (All of them have taken biosafety training)

Xiping Xin, post-doctoral fellow

Rabindranath Bhattacharjee. Post-doctoral fellow

Gannareddy Girish, post-doctoral,fellow
 Neena Lala, Graduate student
 Jose Tores-Garcia, undergraduate research assistant

Mousumi Bhattacharjee, post-doctoral fellow
 Elena Tutunea-Fatan, graduate student
 Lida Raden, undergraduate research assistant

1.0 Microorganisms

1.1 Does your work involve the use of biological agents? YES NO
 (non-pathogenic and pathogenic biological agents including but not limited to bacteria and other microorganisms, viruses, prions, parasites or pathogens of plant or animal origin)? If no, please proceed to Section 2.0

Do you use microorganisms that require a permit from the CFIA? YES NO

If YES, please give the name of the species. _____

What is the origin of the microorganism(s)? _____

Please describe the risk (if any) of escape and how this will be mitigated:

Please attach the CFIA permit.

Please describe any CFIA permit conditions:

1.2 Please complete the table below:

Name of Biological agent(s)*	Is it known to be a human pathogen? YES/NO	Is it known to be an animal pathogen? YES/NO	Is it known to be a zoonotic agent? YES/NO	Maximum quantity to be cultured at one time? (in Litres)	Source/ Supplier	PHAC or CFIA Containment Level
Human placental tissue samples	<input type="radio"/> Yes <input checked="" type="radio"/> X No	<input type="radio"/> Yes <input checked="" type="radio"/> X No	<input type="radio"/> Yes <input checked="" type="radio"/> X No	Not cultured	Local hospitals (as approved, see files)	<input type="radio"/> 1 <input checked="" type="radio"/> 2 <input type="radio"/> X <input type="radio"/> 2+ <input type="radio"/> 3
Human placental trophoblast cell lines HTR-SVneo RSVT-2C,	<input type="radio"/> Yes <input checked="" type="radio"/> XNo	<input type="radio"/> Yes <input checked="" type="radio"/> x No	<input type="radio"/> Yes <input checked="" type="radio"/> X No	In mLs	Produced 18 years ago in my lab (see files)	<input type="radio"/> 1 <input checked="" type="radio"/> 2 <input type="radio"/> X <input type="radio"/> 2+ <input type="radio"/> 3
Choriocarcin oma cell lines Jeg3, JaR	<input type="radio"/> Yes <input checked="" type="radio"/> X No	<input type="radio"/> Yes <input checked="" type="radio"/> X No	<input type="radio"/> Yes <input checked="" type="radio"/> XNo	In mLs	ATCC (see files)	<input type="radio"/> 1 <input checked="" type="radio"/> 2 <input type="radio"/> X <input type="radio"/> 2+ <input type="radio"/> 3
MDA-MB-231, MCF-7, Hs57T, SKBR-3, T47D, MDA-MB-468 human breast cancer cell lines	<input type="radio"/> Yes <input checked="" type="radio"/> XNo	<input type="radio"/> Yes <input checked="" type="radio"/> X No	<input type="radio"/> Yes <input checked="" type="radio"/> XNo	In mLs	ATCC (see files)	<input type="radio"/> 1 <input checked="" type="radio"/> 2 <input type="radio"/> X <input type="radio"/> 2+ <input type="radio"/> 3

C3-L5 and C10 mouse breast cancer cell lines					Produced in my lab 20 years ago (see files)	Level 1
--	--	--	--	--	---	---------

*Please attach a Material Safety Data Sheet or equivalent from the supplier. These agents have already been approved earlier (see earlier files)

2.0 Cell Culture

2.1 Does your work involve the use of cell cultures?

YES X

NO

If no, please proceed to Section 3.0

2.2 Please indicate the type of primary cells (i.e. derived from fresh tissue) that will be grown in culture:

Cell Type	Is this cell type used in your work?	Source of Primary Cell Culture Tissue	AUS Protocol Number
Human	<input type="radio"/> Yes <input checked="" type="radio"/> NoX		Not applicable
Rodent	<input type="radio"/> Yes <input checked="" type="radio"/> NoX		
Non-human primate	<input type="radio"/> Yes <input checked="" type="radio"/> NoX		
Other (specify)	<input type="radio"/> Yes <input checked="" type="radio"/> NoX		

2.3 Please indicate the type of established cells that will be grown in culture in:

Cell Type	Is this cell type used in your work?	Specific cell line(s)*	Supplier / Source
Human	<input checked="" type="radio"/> YesX <input type="radio"/> No	HTR-8/SVneo, RSVT-2C human placental trophoblast cell lines JEG3 and JaR choriocarcinoma cell line MDA-MB-231, MCF-7, Hs57T, SKBR-3, T47D, MDA-MB-468 breast cancer cell lines MDA-MB-468 and its lymphatic metastatic variant MDA-MB-468-LN Breast cancer cell lines, (currently being used for studies of molecular mechanisms in metastasis)	Derived from human placenta in our lab some 18 years ago (see below) ATCC ATCC Dr Anne Chambers, LRCP
Rodent	<input checked="" type="radio"/> YesX <input type="radio"/> No	C3-L5 and C-10 mouse breast cancer cell lines	Derived from a spontaneous breast

* DESCRIPTION MUST BE ATTACHED TO THIS FORM OR PROJECT WILL NOT BE REVIEWED*

			cancer in a C3H/HEJ mouse in our lab some 20 years ago (see below)
Non-human primate	<input type="radio"/> Yes	<input type="radio"/> NoX	
Other (specify)	<input type="radio"/> Yes	<input type="radio"/> NoX	

*Please attach a Material Safety Data Sheet or equivalent from the supplier. (For more information, see www.atcc.org)---- (see earlier files) .

N.B. Cell lines derived in P.K.Lala's lab:

(1) HTR-8/SVneo Human Trophoblast Cell Line:

This cell line was derived at Dr P.K.Lala's laboratory by SV40 Tag immortalization of a short-lived primary human extravillous trophoblast (EVT) cell line HTR-8 isolated from a first trimester human chorionic villus explant in culture. This cell line has all the phenotypic and functional properties of the primary cell line except for being immortal. It is neither precancerous nor cancerous.

Reference: Graham, C.H., Hawley, T.S., Hawley, R.G., MacDougall, J.R., Kerbel, R.S., Khoo, N.K.S. and Lala, P.K. Establishment and characterization of first trimester human trophoblast cells with extended life span. *Exp. Cell Res.* 206: 204-211, 1993.

This cell line is recognized as the gold standard of normal EVT cells used in our past and current experiments outlined in section 1 above, currently also used by many other labs world-wide

(2) RSVT-2 C Human Trophoblast Cell line. RSVT-2 cell line was derived in a similar manner as above, but was selected by long life rather than selection for neomycin. It was found to be long-lived (living for 45 passages when it went through a crisis. A surviving clone was then propagated as RSVT-2C cell line

Both cell lines were found to be precancerous but not cancerous. We have been using this cell line as a model for precancerous trophoblast in experiments outlined in section 1 above.

References : (1)Khoo, N.K.S., Bechberger, J.F., Shepherd, T., Bond, S.L., McCrae, K., Hamilton, G.S. and Lala, P.K. SV40 Tag transformation of the normal invasive trophoblast results in a premalignant phenotype: Mechanisms responsible for hyperinvasiveness and resistance to antiinvasive action of TGFβ. *Intl. J. Cancer.* 77, 429-439, 1998

(2) Khoo, N.K.S., Zhang, Y., Bechberger, J.F., Bond, S.L., Hum, K. and Lala, P.K. SV40 Tag transformation of the normal invasive trophoblast results in a premalignant phenotype: II. Changes in gap junctional intercellular communication. *Intl. J. Cancer.* 77, 440-448, 1998.

(3) C3-L5 and C10 mouse breast cancer cell lines:

C3-L5 cell line was derived by Dr P.K.Lala's laboratory from a spontaneous mammary tumor appearing in a retired female breeder mouse of C3/HeJ strain and showing lung metastasis in the original host as follows. A clone C3 produced from the primary tumor (1983) exhibited high ability for spontaneous lung metastasis from the subcutaneous (SC)transplants in the groin. With repeated passage over several years, the metastatic ability of C3 cells declined gradually, after which C3 cells were reselected in vivo by five cycles of SC to lung micro-metastasis (1987-88), and the resulting line C3L5 has since maintained its highly metastatic ability from SC transplant site to the lungs.

References: (1) Saarloos, M.N., Khoo, N.K.S., and Lala, P.K. Effects of cancer immunotherapy with indomethacin and IL-2 on murine hemopoietic stem cells. *Cancer Research*, 52, 6452-6462, 1992

(2) Lala, P.K. and Parhar, R.S. Eradication of spontaneous and experimental adenocarcinoma metastasis with chronic indomethacin and intermittent IL-2 therapy. *Int. J. Cancer*, 54, 667-684, 1993

C10 cell line was derived as a non-metastatic clone from the same spontaneous mammary tumor. Over the years with repeated passage, it has become weakly metastatic.

Reference: Jadeski, L.C., Hum, K.D., Chakraborty, C. and Lala, P.K. Nitric oxide promotes murine mammary tumor growth and metastasis by stimulating tumor cell migration, invasiveness and angiogenesis. *Int. J. Cancer.* 86, 30-39, 2000.

We have been using C3L5 cells as an in vitro model of metastatic breast cancer, and transplants of these cells in syngeneic C3H/HeJ mice are used in experiments as a preclinical model for COX-2 expressing breast cancer to understand molecular mechanisms in vascular and lymphatic metastasis and their abrogation with various therapies, as outlined in section 1 We have also been using C3L5 and C10 cells for differential gene arrays and have identified a good number of metastasis – promoting and suppressor genes, which are being tested in functional assays.

We are using MDA-MB-468 and MDA-MB-468 LN (supplied by Dr Ann Chambers at the LRCP) for in vitro experiments to understand molecular mechanisms in lymphatic metastasis.

We have given out HTR-8/SVneo human trophoblast and C3-L5 murine breast cancer cell lines to various research labs all over the world on the basis of MTAs approved by the biosafety office,UWO.

2.3 For above named cell types(s) indicate PHAC or CFIA containment level required 1 2 X 2+ 3

3.0 Use of Human Source Materials

3.1 Does your work involve the use of human source materials? YES x NO
 If no, please proceed to Section 4.0

3.2 Indicate in the table below the Human Source Material to be used.

Human Source Material	Source/Supplier /Company Name	Is Human Source Material Infected With An Infectious Agent? YES/NO	Name of Infectious Agent (If applicable)	PHAC or CFIA Containment Level (Select one)
Human Blood (whole) or other Body Fluid	None	<input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> Unknown		<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 2+ <input type="radio"/> 3
Human Blood (fraction) or other Body Fluid	None	<input type="radio"/> Yes <input type="radio"/> No <input type="radio"/> Unknown		<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 2+ <input type="radio"/> 3
Human Organs or Tissues (unpreserved) Human placenta We use the unfixed tissue for extracting RNA to examine mRNA expression of certain invasion-regulating genes (such as decorin) by qPCR, and also fixed tissue for immunocytochemistry of protein expression, or in situ hybridization for mRNA localization of the same molecules (see section I for an outline of the studies)	From UWO hospitals (as approved, see files). Before release, they are Confirmed as non-infectious by the Obstretician, based on the screening of the mothers.for HIV or HBV	<input type="radio"/> Yes <input type="radio"/> No X <input type="radio"/> Unknown		<input type="radio"/> 1 <input type="radio"/> 2 X <input type="radio"/> 2+ <input type="radio"/> 3
Human Organs or Tissues (preserved)	None	Not Applicable		Not Applicable

4.0 Genetically Modified Organisms and Cell lines

4.1 Will genetic modifications be made to the microorganisms, biological agents, or cells described in Sections 1.0 and 2.0? YES NO If no, please proceed to Section 5.0

* DESCRIPTION MUST BE ATTACHED TO THIS FORM OR PROJECT WILL NOT BE REVIEWED*

4.2 Will genetic modification(s) involving plasmids be done? YES X, complete table below NO

Bacteria Used for Cloning *	Plasmid(s) *	Source of Plasmid	Gene Transfected	Describe the change that results
<i>E coli</i>	Human COX-2 cDNA cloned into the eukaryotic expression vector pIRES2-EGFP	Kindly provided by Dr Michael Archer University of Toronto	We introduced Cyclo-oxygenase-(COX)2 gene into human MCF-7 and SKBR-3 breast cancer cell lines by electroporation	These cell lines (named as MCF-7-COX-2 and SKBR-3-COX-2 respectively) show more aggressive behavior such as migratory and invasive capacity and tumosphere formation in tissue culture.

See e-mail

* Please attach a Material Data Sheet or equivalent if available. (No MDS available)

4.3 Will genetic modification(s) involving viral vectors be made? YES, complete table below NOX

Virus Used for Vector Construction	Vector(s) *	Source of Vector	Gene(s) Transduced	Describe the change that results

* Please attach a Material Safety Data Sheet or equivalent.

4.4 Will genetic sequences from the following be involved?

- ◆ HIV YES, please specify _____ NO X
- ◆ HTLV 1 or 2 or genes from any Level 1 or Level 2 pathogens YES, specify _____ NO X
- ◆ SV 40 Large T antigen YES X --had been used to generate our HTR-8/SV neo human placental trophoblast cell line 18 years ago (see files, and documentation above) NO
- ◆ E1A oncogene YES NO X
- ◆ Known oncogenes YES, please specify _____ NO X
- ◆ Other human or animal pathogen and or their toxins YES, please specify _____ NO X

4.5 Will virus be replication defective? YES N/A

4.6 Will virus be infectious to humans or animals? YES N/A

4.7 Will this be expected to increase the containment level required? YES N/A

5.0 Human Gene Therapy Trials

5.1 Will human clinical trials be conducted involving a biological agent? YES NOX
(including but not limited to microorganisms, viruses, prions, parasites or pathogens of plant or animal origin)
If no, please proceed to Section 6.0

5.2 If YES, please specify which biological agent will be used: _____
Please attach a full description of the biological agent.

5.2 Will the biological agent be able to replicate in the host? YES NO

5.3 How will the biological agent be administered? _____

5.4 Please give the Health Care Facility where the clinical trial will be conducted: _____

5.5 Has human ethics approval been obtained? YES, number: _____ NO PENDING

6.0 Animal Experiments

6.1 Will live animals be used? YESX NO If no, please proceed to section 7.0

6.2 Name of animal species to be used _____ Mice _____

6.3 AUS protocol # _2007-057, 2007-036 _____

6.4 Will any of the agents listed in section 4.0 be used in live animals YES, specify: _____ NOX

6.5 Will the agent(s) be shed by the animal: YES NOX, please justify:
Tumors will be transplanted in mice with breast cancer cell lines. They will not shed cells. _____

7.0 Use of Animal species with Zoonotic Hazards

7.1 Will any animals with zoonotic hazards or their organs, tissues, lavages or other body fluids including blood be used (see list below)? YES NoX If no, please proceed to section 8.0

7.2 Please specify the animal(s) used:

- ◆ Pound source dogs YES NO
- ◆ Pound source cats YES NO
- ◆ Cattle, sheep or goats YES NO
- ◆ Non-human primates YES, please specify species _____ NO
- ◆ Wild caught animals YES, please specify species & colony # _____ NO
- ◆ Birds YES NO
- ◆ Others (wild or domestic) YES, please specify _____ NO

8.0 Biological Toxins

8.1 Will toxins of biological origin be used? YES NO X If no, please proceed to Section 9.0

8.2 If YES, please name the toxin(s) _____
Please attach information, such as a Material Safety Data Sheet, for the toxin(s) used.

8.3 What is the LD₅₀ (specify species) of the toxin _____

8.4 How much of the toxin is handled at one time*? _____

8.5 How much of the toxin is stored*? _____

8.6 Will any biological toxins be used in live animals? YES, Please provide details: _____ NO

*For information on biosecurity requirements, please see:

http://www.uwo.ca/humanresources/docandform/docs/healthandsafety/biosafety/Biosecurity_Requirements.pdf

9.0 Insects

9.1 Do you use insects? YES NO X If no, please proceed to Section 10.0

9.2 If YES, please give the name of the species. _____

9.3 What is the origin of the insect? _____

9.4 What is the life stage of the insect? _____

9.5 What is your intention? Initiate and maintain colony, give location: _____
 "One-time" use, give location: _____

9.6 Please describe the risk (if any) of escape and how this will be mitigated:

9.7 Do you use insects that require a permit from the CFIA permit? YES NO
If YES, Please attach the CFIA permit & describe any CFIA permit conditions:

10.0 Plants

10.1 Do you use plants? YES NO X If no, please proceed to Section 11.0

10.2 If YES, please give the name of the species. _____

10.3 What is the origin of the plant? _____

10.4 What is the form of the plant (seed, seedling, plant, tree...)? _____

10.5 What is your intention? Grow and maintain a crop "One-time" use

10.6 Do you do any modifications to the plant? YES NO
If yes, please describe: _____

10.7 Please describe the risk (if any) of loss of the material from the lab and how this will be mitigated:

10.8 Is the CFIA permit attached? YES NO
If YES, Please attach the CFIA permit & describe any CFIA permit conditions:

11.0 Import Requirements

11.1 Will any of the above agents be imported? YES, please give country of origin _____ NO X
If no, please proceed to Section 12.0

11.2 Has an Import Permit been obtained from HC for human pathogens? YES NO

11.3 Has an import permit been obtained from CFIA for animal or plant pathogens? YES NO

11.4 Has the import permit been sent to OHS? YES, please provide permit # _____ NO

12.0 Training Requirements for Personnel Named on Form

All personnel named on the above form who will be using any of the above named agents are required to attend the following training courses given by OHS:

- ◆ Biosafety
- ◆ Laboratory and Environmental/Waste Management Safety
- ◆ WHMIS (Western or equivalent)
- ◆ Employee Health and Safety Orientation

As the Principal Investigator, I have ensured that all of the personnel named on the form who will be using any of the biohazardous agents in Sections 1.0 to 9.0 have been trained.

SIGNATURE _____ P.K.Lala _____

13.0 Containment Levels

13.1 For the work described in sections 1.0 to 9.0, please indicate the highest HC or CFIA Containment Level required. 1 2 2+ 3

13.2 Has the facility been certified by OHS for this level of containment?
 YES, permit # if on-campus **BIO-UWO-0018, MSB 433A inspected Sept 3, 2009**
 NO, please certify
 NOT REQUIRED for Level 1 containment

14.0 Procedures to be Followed

14.1 As the Principal Investigator, I will ensure that this project will follow the Western Biosafety Guidelines and Procedures Manual for Containment Level 1 & 2 Laboratories (and the Level 3 Facilities Manual for Level 3 projects). I will ensure that UWO faculty, staff and students working in my laboratory have an up-to-date Hazard Communication Form, found at <http://www.wph.uwo.ca/>

SIGNATURE _____ P.K.Lala _____ Date: _____ 12 July
2010 _____

14.2 Please describe additional risk reduction measures will be taken beyond containment level 1, 2, 2+ or 3 measures, that are unique to this agent.--- N/A

14.3 Please outline what will be done if there is an exposure to the biohazards listed, such as a needlestick injury: Immediate local cleansing and sterilization with soap and running water, squeezing the area to drain out blood, followed by the use of University health services.

15.0 Approvals

1) UWO Biohazard Subcommittee: SIGNATURE: _____
Date: _____

2) Safety Officer for the University of Western Ontario
SIGNATURE: _____
Date: _____

3) Safety Officer for Institution where experiments will take place (if not UWO):

SIGNATURE: _____

Date: _____

Approval Number: _____ Expiry Date (3 years from Approval): _____


Special Conditions of Approval:

I did the tranfection of MCF 7 cells. I used DH5alpha E.Coli cell for transformation of plasmid (to increase volume of plasmid) and used that plasmid (with the microporator machine) to transfect MCF 7 cell.

Hope this will help you.

Sincerely
Mousumi

Mousumi Majumder PhD
Post Doctoral Fellow
The University of Western Ontario
Department of Anatomy & Cell Biology
Medical Sciences Building, M430
London, Ontario
N6A 5C1 Canada
Tel.No: 519-661-2111 extn.80517 office
519-204-3741 resi.
Email: mmajumde@uwo.ca
majumder.mousumi@gmail.com



Section 4.2

Cell line info

Cell Biology

ATCC® Number: **HTB-26™** [Order this Item](#) Price: **\$256.00**

Designations: MDA-MB-231
Depositors: R Cailleau
Biosafety Level: 1
Shipped: frozen
Medium & Serum: [See Propagation](#)
Growth Properties: adherent
Organism: *Homo sapiens* (human)

epithelial

Morphology:



Organ: mammary gland; breast

Disease: adenocarcinoma

Source:

Derived from metastatic site: pleural effusion

Cell Type: epithelial

In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Permits/Forms:

Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

Receptors: epidermal growth factor (EGF), expressed
transforming growth factor alpha (TGF alpha), expressed

Tumorigenic: Yes

Amelogenin: X

CSF1PO: 12,13

D13S317: 13

D16S539: 12

DNA Profile (STR): D5S818: 12

D7S820: 8,9

TH01: 7,9.3

TPOX: 8,9

vWA: 15,18

Cytogenetic Analysis:

The cell line is aneuploid female (modal number = 64, range = 52 to 68), with chromosome counts in the near-triploid range. Normal chromosomes N8 and N15 were absent. Eleven stable rearranged marker chromosomes are noted as well as unassignable chromosomes in addition to the majority of autosomes that are trisomic. Many of the marker chromosomes are identical to those shown in the karyotype reported by K.L. Satya-Prakash, et al.

Related Links ▶

[NCBI Entrez Search](#)

[Cell Micrograph](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[Related Cell](#)

[Culture Products](#)

Login

Required ▶

[Product Information Sheet](#)

[BioProducts](#)

[Cell, microbial and molecular genomics products for the life](#)

- [sciences BioServices](#)

[Bio-materials management: basic](#)

[repository to complex partnership-](#)

- [level services BioStandards](#)

[Biological Reference Material and Consensus Standards for the life science](#)

- [community](#)

Cell Biology

ATCC® Number: **HTB-22™** Price: **\$272.00**

Designations: MCF7
 Depositors: CM McGrath
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)

epithelial

Morphology:



Organ: mammary gland; breast

Disease: adenocarcinoma

Source:

Derived from metastatic site: pleural effusion

Cell Type: epithelial

Cellular Products:

insulin-like growth factor binding proteins (IGFBP) BP-2;
BP-4; BP-5

In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this

Permits/Forms:

ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Applications:

transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

Receptors:

estrogen receptor, expressed

Antigen Expression: Blood Type O; Rh+

Amelogenin: X

CSF1PO: 10

D13S317: 11

D16S539: 11,12

DNA Profile (STR): D5S818: 11,12

D7S820: 8,9

TH01: 6

TPOX: 9,12

vWA: 14,15

modal number = 82; range = 66 to 87.

The stemline chromosome numbers ranged from hypertriploidy to hypotetraploidy, with the 2S component occurring at 1%.

There were 29 to 34 marker chromosomes per S metaphase; 24 to 28 markers occurred in at least 30% of cells, and generally one large submetacentric (M1) and 3 large subtelocentric (M2, M3, and M4) markers were recognizable in over 80% of metaphases. No DM were detected. Chromosome 20 was nullisomic and X was disomic.

Cytogenetic Analysis:

Related Links ▶

[NCBI Entrez Search](#)

[Cell Micrograph](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[Related Cell](#)

[Culture Products](#)

Login

Required ▶

[Product](#)

[Information Sheet](#)

[BioProducts](#)

[Cell, microbial and molecular genomics products for the life](#)

- [sciences](#)

[BioServices](#)

[Bio-materials management; basic repository to complex partnership-](#)

- [level services](#)

[BioStandards](#)

[Biological Reference Material and Consensus Standards for the life science](#)

- [community](#)

Cell Biology

ATCC® Number: **HTB-133™** [Order this Item](#) Price: **\$256.00**

Designations: T-47D
 Depositors: I Keydar
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)

epithelial

Morphology:



Organ: mammary gland; breast

Tissue: duct

Source:

Disease: ductal carcinoma

Derived from metastatic site: pleural effusion

In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Permits/Forms:

Applications:

transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

calcitonin, expressed

androgen receptor, expressed

estrogen receptor, expressed

progesterone receptor, expressed

Receptors:

glucocorticoid receptor, positive, expressed

prolactin, expressed

calcitonin; androgen receptor, positive; progesterone receptor, positive; glucocorticoid; prolactin; estrogen receptor, positive

Amelogenin: X

CSF1PO: 11,13

D13S317: 12

D16S539: 10

DNA Profile (STR): D5S818: 12

D7S820: 11

THO1: 6

TPOX: 11

vWA: 14

Cytogenetic
 Analysis:

Related Links ▶

[NCBI Entrez Search](#)

[Cell Micrograph](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[Related Cell](#)

[Culture Products](#)

Login

Required ▶

[Product Information Sheet](#)

[BioProducts](#)

[Cell, microbial and molecular genomics products for the life](#)

- [sciences](#)

[BioServices](#)

[Bio-materials management: basic repository to complex partnership-](#)

- [level services](#)

[BioStandards](#)

[Biological Reference Material and Consensus Standards for the life](#)

[science](#)

- [community](#)

Cell Biology

ATCC® Number: **HTB-132™** Order this Item Price: **\$272.00**

Designations: MDA-MB-468
 Depositors: R Cailleau
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)
 Morphology: epithelial

Source: **Organ:** mammary gland; breast
Disease: adenocarcinoma
 In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Permits/Forms:

Isolation: **Isolation date:** 1977
 Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))
 Receptors: epidermal growth factor (EGF)
 transforming growth factor alpha (TGF alpha)
 Tumorigenic: Yes
 Antigen Expression: Blood Type AB; HLA Aw23, Aw30, B27, Bw35, Cw2, Cw4 (patient)
 Amelogenin: X
 CSF1PO: 12
 D13S317: 12
 D16S539: 9
 DNA Profile (STR): D5S818: 12
 D7S820: 8
 THO1: 7
 TPOX: 8,9
 vWA: 18

modal number = 64; range = 60 to 67.
 The cell line is aneuploid human, presumably female (X, abnormal X) with most chromosome counts in the hypotriploid range.; Normal chromosomes X, N2, N3, N7, N8, N10, and N22 are clearly under-represented due to their involvement in the formation of the many marker (19) chromosomes present in this cell line.; A normal chromosome N1 (or two) is identified in each karyotype, but, in addition, regions of chromosome N1 are also present in five different marker chromosomes.; Variation is evident in the normal and marker chromosome copy number from karyotype to karyotype.

Cytogenetic Analysis:

Related Links ▶

[NCBI Entrez Search](#)
[Make a Deposit](#)
[Frequently Asked Questions](#)
[Material Transfer Agreement](#)
[Technical Support](#)
[Related Cell Culture Products](#)

Login Required ▶

[Product Information Sheet](#)


[BioProducts](#)

[Cell, microbial and molecular genomics products for the life sciences](#)
[BioServices](#)

- [Bio-materials management; basic repository to complex partnership-level services](#)
- **[BioStandards](#)**

- [Biological Reference Material and Consensus Standards for the life science community](#)

Plasmid Map(s)



45