

**THE UNIVERSITY OF WESTERN ONTARIO
BIOLOGICAL AGENTS REGISTRY FORM**
Approved Biohazards Subcommittee: October 14, 2010
Biosafety Website: www.uwo.ca/humanresources/biosafety/

This form must be completed by each Principal Investigator holding a grant administered by the University of Western Ontario (UWO) or in charge of a laboratory/facility where the use of Level 1, 2 or 3 biological agents is described in the laboratory or animal work proposed. The form must also be completed if any work is proposed involving animals carrying zoonotic agents infectious to humans or involving plants, fungi, or insects that require Public Health Agency of Canada (PHAC) or Canadian Food Inspection Agency (CFIA) permits.

This form must be updated at least every 3 years or when there are changes to the biological agents being used.

Containment Levels will be established in accordance with Laboratory Biosafety Guidelines, 3rd edition, 2004, Public Health Agency of Canada (PHAC) or Containment Standards for Veterinary Facilities, 1st edition 1996, Canadian Food Inspection Agency (CFIA).

Completed forms are to be returned to Occupational Health and Safety, (OHS), (Support Services Building, Room 4190) for distribution to the Biohazards Subcommittee. For questions regarding this form, please contact the Biosafety Officer at extension 81135 or biosafety@uwo.ca. If there are changes to the information on this form (excluding grant title and funding agencies), contact Occupational Health and Safety for a modification form. See website: www.uwo.ca/humanresources/biosafety

PRINCIPAL INVESTIGATOR	<u>David Litchfield</u>
DEPARTMENT	<u>Biochemistry</u>
ADDRESS	<u>The University of Western Ontario</u> <u>London ON</u> <u>N6A 5C1</u>
PHONE NUMBER	<u>519-661-2111 ext 86849 (lab) ext 84186 (office)</u>
EMERGENCY PHONE NUMBER(S)	<u>519-657-8751</u>
EMAIL	<u>litchfi@uwo.ca</u>

Location of experimental work to be carried out: Building(s) _____ MSB _____ Room(s) 355/359/365A/380

*For work being performed at Institutions affiliated with the University of Western Ontario, the Safety Officer for the Institution where experiments will take place must sign the form prior to its being sent to the University of Western Ontario Biosafety Officer (See Section 15.0, Approvals).

FUNDING AGENCY/AGENCIES: Canadian Cancer Society Research Institute, Canadian Institutes of Health Research, Cancer Research Society, Inc

GRANT TITLE(S): Regulation and role of protein kinase CK2 in cell cycle progression (Canadian Cancer Society Research Institute), Control of cell proliferation & survival: Convergence of CK2 & caspase signaling pathways (Canadian Institutes of Health Research), Rational Design of Novel Inhibitors of Cancer Cell Proliferation (Cancer Research Society, Inc)

List all personnel working under Principal Investigators supervision in this location:

<u>Name</u>	<u>UWO E-mail Address</u>	<u>Date of Biosafety Training</u>
<u>Laszlo Gyenis</u>	<u>lgyenis@uwo.ca</u>	<u>2007</u>
<u>Michelle Gabriel</u>	<u>mgabrie5@uwo.ca</u>	<u>2009</u>
<u>Dana Onica</u>	<u>donica2@uwo.ca</u>	<u>2010</u>
<u>Brenden Innes</u>	<u>binnes22@uwo.ca</u>	<u>2008</u>
<u>Stephanie Zukowski</u>	<u>szukowsk@uwo.ca</u>	<u>2010</u>
<u>Jacob Turowec</u>	<u>jtrowec@uwo.ca</u>	<u>2006</u>
<u>Kathryn Garside</u>	<u>kgarside@uwo.ca</u>	<u>2009</u>

Please explain the biological agents and/or biohazardous substances used and how they will be stored, used and disposed of. Projects without this description will not be reviewed.

Use: All biological agents and/or biohazardous substances are used according to the UWO Biosafety Guidelines and Procedures Manual following good microbiological laboratory practices. All trained laboratory personnel are working with biological agents and/or biohazardous substances only inside the containment Level 2 laboratories. The cell lines are handled in the biological safety cabinets that are re-certified annually. **Storage:** All biological agents and/or biohazardous substances are stored only inside the containment laboratory that has lockable doors. The containment laboratory is always kept locked when it is unoccupied. **Disposal:** All biological agents and/or biohazardous substances are autoclaved, incinerated or disinfected to inactivate the biological agents prior to disposal. Autoclaving of biological agents are done on-site in functioning autoclaves.

Please include a one page research summary or teaching protocol.

Regulation and role of protein kinase CK2 in cell cycle progression (Canadian Cancer Society Research Institute). The growth and division of cells is controlled by a network of enzymes known as protein kinases that frequently contribute to tumorigenesis when they become active at the wrong time in cancer cells. Our research is focused on one group of these protein kinases known as CK2 because this group of enzymes is present at abnormally high levels in many forms of human cancer and because CK2 has been shown to promote breast cancer and leukemia in mice. We are working to understand how CK2 contributes to the inappropriate division of cancer cells with the hope that this information may provide new ideas for improving cancer treatment. To complement this work, we have been involved in the evaluation of novel CK2 inhibitors to determine whether these compounds can inhibit CK2 functions in cells and to examine whether they have any other undesirable effects on the cells. The latter studies are essential if CK2-targeted therapies are to move forward for clinical testing. One aspect of this project is focused on performing a detailed characterization of new compounds that have been designed, on the basis of the three-dimensional structure of CK2, to inhibit CK2 in cells. We will employ new methods, involving proteomics and genetic approaches, to characterize the effects of these compounds on cancer cells to ensure that these compounds do indeed target CK2 in cells and to determine whether these compounds have other cellular targets or side effects. A second aspect of this project will be focused on determining how CK2 is involved in the control of cell division since this function of CK2 is likely to be related to the ability of CK2 to promote tumorigenesis. This aspect of the work will be fostered by an examination of unique experimental systems (human cancer cells that have been engineered to express mutant forms of CK2) that we have developed.

Control of cell proliferation & survival: Convergence of CK2 & caspase signaling pathways (Canadian Institutes of Health Research). The overall objective of our ongoing studies is directed towards elucidation of the molecular mechanisms that control cell proliferation and survival. Biochemical and genetic studies have demonstrated the existence of networks of protein kinases that are involved in the regulation of cellular events that determine whether cells live or die. Cells also contain networks of proteases, including aspartic-acid specific proteases known as caspases that play a major role in the regulation of cell survival through their involvement in the initiation and execution phases of apoptosis. Perturbations in regulatory protein kinase and caspase networks induce alterations in cell survival and frequently accompany transformation and tumorigenesis. Furthermore, the convergence of protein kinase and caspase signaling pathways has become increasingly evident, as phosphorylation of caspase substrates within caspase recognition motifs has been shown to prevent caspase-catalyzed cleavage of a number of proteins. To systematically investigate the global role of phosphorylation in the regulation of caspase signaling, we employed computational approaches to identify protein sequences from the human proteome that contained overlapping protein kinase and caspase recognition motifs. Notably, the constitutively active and oncogenic protein kinase CK2 emerged as the most prominent kinase displaying an overlapping consensus for phosphorylation of caspase recognition motifs. HYPOTHESIS: On the basis of these observations and a striking overlap between the substrate specificity of CK2 and caspases, the working hypothesis for our studies is that CK2 protects cells from apoptosis through its ability to modulate the susceptibility of regulatory cellular proteins to caspase cleavage. To test this hypothesis, we have employed systematic computational and functional proteomic approaches that revealed many novel candidate targets with overlapping CK2 and caspase sites including pro-caspase 3 and pro-caspase 8. As a logical extension of our work, the two major aims of this project will be: i) to validate candidate proteins as direct CK2 and caspase targets, and ii) to elucidate mechanisms controlling CK2 phosphorylation of apoptotic proteins. The first aim will be addressed by studies to evaluate candidates as targets for CK2 and caspase in vitro and in cells, and an examination of the functional effects of CK2 phosphorylation on apoptosis. The second aim will involve studies to determine whether CK2 activity changes in response to apoptotic stimuli and to characterize complexes between CK2 and its substrates during apoptosis. Given the emergence of CK2 as a promising candidate for molecular-targeted therapy, we envisage that our studies could yield insights that will contribute to improvements in cancer treatment in leukemia or tumors with elevated levels of CK2.

Rational Design of Novel Inhibitors of Cancer Cell Proliferation (Cancer Research Society, Inc). The long-term goal of our studies is to develop innovative knowledge-based strategies for the control of cell proliferation and those diseases such as cancer that are characterized by inappropriate cell proliferation. These studies focus on Pin1, a peptidyl-prolyl isomerase that has emerged as a potential prognostic indicator and promising therapeutic target for cancer. Based on its essential role in cell division, elevated expression in several cancers and regulatory interactions with oncogene and tumor suppressor gene product, the underlying hypothesis for our work is that Pin1 will be an effective target for the development of novel molecular-targeted cancer therapies. Towards this objective, we undertook studies employing a broad base of experimental strategies ranging from yeast genetics to structural biology that generated the most extensive mutagenic analysis of Pin1 and new insights regarding its catalytic mechanism. We also performed an unbiased screen to identify novel inhibitors of Pin1 that yielded cyclic peptides able to inhibit Pin1 at sub-micromolar concentrations. Using NMR spectroscopy, we have determined the high-resolution structure of the inhibitory peptide (CRYPEVEIC designated CCP-1) and characterized its mode of interaction with Pin1. To capitalize on our advances, the objectives of this proposal are: i) to develop specific cell-permeable inhibitors of Pin1, and ii) to perform inhibitor validation studies in cancer cells (ie. verify inhibition of Pin1 and evaluate off-target effects).

1.0 Microorganisms

1.1 Does your work involve the use of biological agents? YES NO
 (non-pathogenic and pathogenic biological agents including but not limited to bacteria and other microorganisms, viruses, prions, parasites or pathogens of plant or animal origin)? If no, please proceed to Section 2.0

Do you use microorganisms that require a permit from the CFIA? YES NO

If YES, please give the name of the species. _____

What is the origin of the microorganism(s)? _____

Please describe the risk (if any) of escape and how this will be mitigated:

Please attach the CFIA permit.

Please describe any CFIA permit conditions:

1.2 Please complete the table below:

Name of Biological Agent(s)* (Be specific)	Is it known to be a human pathogen? YES/NO	Is it known to be an animal pathogen? YES/NO	Is it known to be a zoonotic agent? YES/NO	Maximum quantity to be cultured at one time? (in Litres)	Source/ Supplier	PHAC or CFIA Containment Level
E. coli (DH5a)	<input type="radio"/> Yes <input checked="" type="radio"/> No	<input type="radio"/> Yes <input checked="" type="radio"/> No	<input type="radio"/> Yes <input checked="" type="radio"/> No	10		<input checked="" type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 2+ <input type="radio"/> 3
E. coli (XL 1 Blue)	<input type="radio"/> Yes <input checked="" type="radio"/> No	<input type="radio"/> Yes <input checked="" type="radio"/> No	<input type="radio"/> Yes <input checked="" type="radio"/> No	2		<input checked="" type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 2+ <input type="radio"/> 3
E. coli (BL21)	<input type="radio"/> Yes <input checked="" type="radio"/> No	<input type="radio"/> Yes <input checked="" type="radio"/> No	<input type="radio"/> Yes <input checked="" type="radio"/> No	10		<input checked="" type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 2+ <input type="radio"/> 3
S. cerevisia	<input type="radio"/> Yes <input checked="" type="radio"/> No	<input type="radio"/> Yes <input checked="" type="radio"/> No	<input type="radio"/> Yes <input checked="" type="radio"/> No	2		<input checked="" type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 2+ <input type="radio"/> 3

*Please attach a Material Safety Data Sheet or equivalent from the supplier.

2.0 Cell Culture

2.1 Does your work involve the use of cell cultures? YES NO

If no, please proceed to Section 3.0

2.2 Please indicate the type of primary cells (i.e. derived from fresh tissue) that will be grown in culture:

Cell Type	Is this cell type used in your work?	Source of Primary Cell Culture Tissue	AUS Protocol Number
Human	<input type="radio"/> Yes <input checked="" type="radio"/> No		Not applicable
Rodent	<input type="radio"/> Yes <input checked="" type="radio"/> No		
Non-human primate	<input type="radio"/> Yes <input checked="" type="radio"/> No		
Other (specify)	<input type="radio"/> Yes <input checked="" type="radio"/> No		

2.3 Please indicate the type of established cells that will be grown in culture in:

Cell Type	Is this cell type used in your work?	Specific cell line(s)*	Containment Level of each cell line	Supplier / Source of cell line(s)
Human	X Yes O No	A431 A549 HEK293T HeLa HeLa (Tet-off) HeLa S3 Hela-Fas Jurkat LNCaP MCF7 PANC-1 Saos-2 U2OS UTA6 (U2OS-derived)	1 1 2 2 2 2 1 1 1 1 1 1 1	ATCC, Clontech, collaborative labs
Rodent	X Yes O No	3T3-L1 (LTA5) [#] BV2 [#] C2C12 [#] C57B1/6 [#] E7 hybridoma [#] F9 Cells [#] G182 [#] Morphs [#] NIH 3T3 [#] Swiss 3T3 [#] Note: [#] for storage purposes only	1 1 1 1 1	ATCC, Clontech, collaborative labs
Non-human primate	X Yes O No	Cos7	2	ATCC
Other (specify)	X Yes O No	BK3A (chicken) [#] Note: [#] for storage purposes only		collaborative labs

*Please attach a Material Safety Data Sheet or equivalent from the supplier. (For more information, see www.atcc.org)

2.4 For above named cell types(s) indicate PHAC or CFIA containment level required 1 2 2+ 3

3.0 Use of Human Source Materials

3.1 Does your work involve the use of human source materials? YES NO
If no, please proceed to Section 4.0

3.2 Indicate in the table below the Human Source Material to be used.

Human Source Material	Source/Supplier /Company Name	Is Human Source Material Infected With An Infectious Agent? YES/UNKNOWN	Name of Infectious Agent (If applicable)	PHAC or CFIA Containment Level (Select one)
Human Blood (whole) or		<input type="radio"/> Yes		<input type="radio"/> 1 <input type="radio"/> 2

other Body Fluid		<input type="radio"/> Unknown		<input type="radio"/> 2+ <input type="radio"/> 3
Human Blood (fraction) or other Body Fluid		<input type="radio"/> Yes <input type="radio"/> Unknown		<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 2+ <input type="radio"/> 3
Human Organs or Tissues (unpreserved)		<input type="radio"/> Yes <input type="radio"/> Unknown		<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 2+ <input type="radio"/> 3
Human Organs or Tissues (preserved)		Not Applicable		Not Applicable

4.0 Genetically Modified Organisms and Cell lines

4.1 Will genetic modifications be made to the microorganisms, biological agents, or cells described in Sections 1.0 and 2.0? YES NO If no, please proceed to Section 5.0

4.2 Will genetic modification(s) involving plasmids be done? YES, complete table below NO

Bacteria Used for Cloning *	Plasmid(s) **	Source of Plasmid	Gene Transfected	Describe the change that results from transformation or tranfection
E. coli (DH5a, XL 1 Blue, BL21)	pBI, pRc/CMV, pTRE, pEGFP, pcDNA3.1, NME3	Invitrogen, Clontech, Addgene	Different genes depending on the experiment	Overexpression of gene product

* Please attach a Material Data Sheet or equivalent if available.

** Please attach a plasmid map.

4.3 Will genetic modification(s) of bacteria and/or cells involving viral vectors be made?

YES, complete table below NO

Virus Used for Vector Construction	Vector(s) *	Source of Vector	Gene(s) Transduced	Describe the change that results from transduction

* Please attach a Material Safety Data Sheet or equivalent.

4.4 Will genetic sequences from the following be involved?

- ◆ HIV YES, please specify _____ NO
- ◆ HTLV 1 or 2 or genes from any Level 1 or Level 2 pathogens YES, specify _____ NO
- ◆ SV 40 Large T antigen YES, please specify __Cos7 NO
- ◆ E1A oncogene YES NO
- ◆ Known oncogenes YES, please specify __CK2 _____ NO
- ◆ Other human or animal pathogen and or their toxins YES, please specify _____ NO

4.5 Will virus be replication defective? YES NO N/A

4.6 Will virus be infectious to humans or animals? YES NO N/A

4.7 Will this be expected to increase the containment level required? YES NO

5.0 Human Gene Therapy Trials

5.1 Will human clinical trials be conducted involving a biological agent? YES NO
 (including but not limited to microorganisms, viruses, prions, parasites or pathogens of plant or animal origin)
 If no, please proceed to Section 6.0

5.2 If YES, please specify which biological agent will be used: _____
 Please attach a full description of the biological agent.

5.2 Will the biological agent be able to replicate in the host? YES NO

5.3 How will the biological agent be administered? _____

5.4 Please give the Health Care Facility where the clinical trial will be conducted: _____

5.5 Has human ethics approval been obtained? YES, number: _____ NO PENDING

6.0 Animal Experiments

6.1 Will live animals be used? YES NO If no, please proceed to section 7.0

6.2 Name of animal species to be used _____

6.3 AUS protocol # _____

6.4 Will any of the agents listed in section 4.0 be used in live animals YES, specify: _____ NO

6.5 Will the agent(s) be shed by the animal: YES NO, please justify:

7.0 Use of Animal species with Zoonotic Hazards

7.1 Will any animals with zoonotic hazards or their organs, tissues, lavages or other body fluids including blood be used (see list below)? YES No If no, please proceed to section 8.0

7.2 Will live animals be used? YES No

7.3 If yes, please specify the animal(s) used:

- ◆ Pound source dogs YES NO
- ◆ Pound source cats YES NO
- ◆ Cattle, sheep or goats YES, please specify species _____ NO
- ◆ Non-human primates YES, please specify species _____ NO
- ◆ Wild caught animals YES, please specify species & colony # _____ NO
- ◆ Birds YES, please specify species _____ NO
- ◆ Others (wild or domestic) YES, please specify _____ NO

7.4 If no live animals are used, please specify the source of the specimens:

8.0 Biological Toxins

8.1 Will toxins of biological origin be used? YES NO If no, please proceed to Section 9.0

8.2 If YES, please name the toxin(s)_Okadaic Acid (phosphatase inhibitor) _____
Please attach information, such as a Material Safety Data Sheet, for the toxin(s) used.

8.3 What is the LD₅₀ (specify species) of the toxin _____ Oral - mouse - 0.15 mg/kg _____

8.4 How much of the toxin is handled at one time*? _____ 5ug _____

8.5 How much of the toxin is stored*? _____ 100ug _____

8.6 Will any biological toxins be used in live animals? YES, Please provide details: _____ NO

*For information on biosecurity requirements, please see:

http://www.uwo.ca/humanresources/docandform/docs/healthandsafety/biosafety/Biosecurity_Requirements.pdf

9.0 Insects

9.1 Do you use insects? YES NO If no, please proceed to Section 10.0

9.2 If YES, please give the name of the species. _____

9.3 What is the origin of the insect? _____

9.4 What is the life stage of the insect? _____

9.5 What is your intention? Initiate and maintain colony, give location: _____
 "One-time" use, give location: _____

9.6 Please describe the risk (if any) of escape and how this will be mitigated:

9.7 Do you use insects that require a permit from the CFIA permit? YES NO

If YES, Please attach the CFIA permit & describe any CFIA permit conditions:

10.0 Plants

10.1 Do you use plants? YES NO If no, please proceed to Section 11.0

10.2 If YES, please give the name of the species. _____

10.3 What is the origin of the plant? _____

10.4 What is the form of the plant (seed, seedling, plant, tree...)? _____

10.5 What is your intention? Grow and maintain a crop "One-time" use

10.6 Do you do any modifications to the plant? YES NO

If yes, please describe: _____

10.7 Please describe the risk (if any) of loss of the material from the lab and how this will be mitigated:

10.8 Is the CFIA permit attached? YES NO

If YES, Please attach the CFIA permit & describe any CFIA permit conditions:

11.0 Import Requirements

11.1 Will any of the above agents be imported? YES, please give country of origin _____ NO

If no, please proceed to Section 12.0

11.2 Has an Import Permit been obtained from HC for human pathogens? YES NO

11.3 Has an import permit been obtained from CFIA for animal or plant pathogens? YES NO


11.4 Has the import permit been sent to OHS? YES, please provide permit # _____ NO

12.0 Training Requirements for Personnel Named on Form

All personnel named on the above form who will be using any of the above named agents are required to attend the following training courses given by OHS:

- ◆ Biosafety
- ◆ Laboratory and Environmental/Waste Management Safety
- ◆ WHMIS (Western or equivalent)
- ◆ Employee Health and Safety Orientation

As the Principal Investigator, I have ensured that all of the personnel named on the form who will be using any of the biological agents in Sections 1.0 to 9.0 have been trained.



SIGNATURE

13.0 Containment Levels

13.1 For the work described in sections 1.0 to 9.0, please indicate the highest HC or CFIA Containment Level required. 1 2 2+ 3

13.2 Has the facility been certified by OHS for this level of containment?
 YES, date of most recent biosafety inspection: July 22, 2010
 NO, please certify
 NOT REQUIRED for Level 1 containment

13.3 Please indicate permit number (not applicable for first time applicants): BIO-UWO-0069

14.0 Procedures to be Followed

14.1 Please describe additional risk reduction measures will be taken beyond containment level 1, 2, 2+ or 3 measures, that are unique to this agent.

The general precautions that are outlined in UWO biosafety manual section 3.0 SAFETY PRACTICES AND PROCEDURES will be followed

14.2 Please outline what will be done if there is an exposure to the biological agents listed, such as a needlestick injury or an accidental splash:

The emergency response procedures will be followed that are outlined in the UWO biosafety manual section 3.5 MEDICAL PROCEDURES AND INCIDENT REPORTING

14.3 As the Principal Investigator, I will ensure that this project will follow the Western Biosafety Guidelines and Procedures Manual for Containment Level 1 & 2 Laboratories (and the Level 3 Facilities Manual for Level 3 projects). I will ensure that UWO faculty, staff and students working in my laboratory have an up-to-date Hazard Communication Form, found at <http://www.wph.uwo.ca/>



SIGNATURE:

Date: March 8, 2011

15.0 Approvals

1) UWO Biohazards Subcommittee: SIGNATURE: _____
Date: _____

2) Safety Officer for the University of Western Ontario
SIGNATURE: _____
Date: _____

3) Safety Officer for Institution where experiments will take place (if not UWO):
SIGNATURE: _____
Date: _____

Approval Number: _____ Expiry Date (3 years from Approval): _____

Special Conditions of Approval:

Control of Receptor-induced Signaling Complex Formation by the Kinetics of Ligand/Receptor Interaction*

Received for publication, July 23, 2002, and in revised form, September 3, 2002
Published, JBC Papers in Press, September 4, 2002, DOI 10.1074/jbc.M207399200

Anja Krippner-Heidenreich, Fabian Tübing, Susanne Bryde, Sylvia Willi, Gudrun Zimmermann,
and Peter Scheurich‡

From the Institute of Cell Biology and Immunology, University of Stuttgart, Allmandring 31, 70569 Stuttgart, Germany

Tumor necrosis factor (TNF) exists both as a membrane-integrated type II precursor protein and a soluble cytokine that have different bioactivities on TNFR2 (CD120b) but not on TNFR1 (CD120a). To identify the molecular basis of this disparity, we have investigated receptor chimeras comprising the cytoplasmic part of Fas (CD95) and the extracellular domains of the two TNF receptors. The membrane form of TNF, but not its soluble form, was capable of inducing apoptosis as well as activation of c-Jun N-terminal kinase and NF- κ B via the TNFR2-derived chimera. In contrast, the TNFR1-Fas chimera displayed strong responsiveness to both TNF forms. This pattern of responsiveness is identical to that of wild type TNF receptors, demonstrating that the underlying mechanisms are independent of the particular type of the intracellular signaling machinery and rather are controlled upstream of the intracellular domain. We further demonstrate that the signaling strength induced by a given ligand/receptor interaction is regulated at the level of adaptor protein recruitment, as shown for FADD, caspase-8, and TRAF2. Since both incidents, strong signaling and robust adaptor protein recruitment, are paralleled by a high stability of individual ligand-receptor complexes, we propose that half-lives of individual ligand-receptor complexes control signaling at the level of adaptor protein recruitment.

The majority of cell surface receptors initiate intracellular signals after ligand-mediated homo- or heteromultimerization. Members of the tumor necrosis factor (TNF)¹ ligand family typically form stable homotrimers capable of multimerizing their respective receptors (1). Essential for intracellular signal induction is the subsequent recruitment of adaptor proteins (e.g. members of the TNF receptor-associated factor (TRAF) family or death domain-containing proteins like TNF receptor-associated death domain protein (TRADD) or Fas-associated

death domain protein (FADD)) (1). These two protein groups define the two major signaling pathways leading to either gene induction (TRAFs) or the activation of the apoptotic program via autoproteolytic cleavage of initiator caspases by induced proximity (1).

TNF binds to two membrane receptors, TNFR1 (CD120a; p55/60) and TNFR2 (CD120b; p75/80). Whereas TNFR1 seems to be constitutively expressed in most tissues, TNFR2 expression is more restricted and can be found especially in immune cells but also in endothelial and neuronal tissue (2). TNFR2 is a typical member of the non-death domain-containing subgroup of the TNF receptor family. It directly binds TRAF2, leading to the activation of NF- κ B and the c-Jun N-terminal kinase (JNK). TNFR1 carries a death domain in its cytoplasmic part and therefore represents a direct activator of apoptotic caspases after recruitment of TRADD and FADD. In parallel to its cytotoxic activity (and this seems to be unique within the TNF receptor family), TNFR1 is a strong activator of gene induction. Receptor-bound TRADD serves as an assembly platform also for recruitment of TRAF2 and receptor-interacting protein (3), which act together in the activation of the inhibitor of κ B kinases, leading to the activation of NF- κ B (4).

Many aspects of the initial events during signal initiation of TNF, however, are poorly understood. After ligand binding, TNF receptor complexes are internalized or may be, alternatively, proteolytically cleaved (5–7). Internalization has been shown to be important for intracellular signal initiation by TNFR1 (8) but not for others like Fas (CD95, APO-1) (8, 9). Further, in some cellular systems the activation of the initiator caspase-8 was found to be necessary for Fas cluster formation (9).

Remarkably, most ligands of the TNF family are expressed as type II transmembrane proteins from which soluble ligands are formed by proteolytic cleavage. In a recent publication, we have compared the signaling capacity of the membrane-bound (memTNF) versus the soluble form of TNF (sTNF) and have demonstrated that TNFR2, but not TNFR1, differentially responds to these two ligand forms (10). It was shown that memTNF, when acting on TNFR2, displays a superior capacity to initiate various cellular responses in a positive cooperative manner with TNFR1. Full TNFR2 activation can even cause a shift in the phenotype of the respective cellular response to sTNF (10). In addition, we have developed a TNFR2-specific monoclonal antibody, termed 80M2, that mimics the bioactivity of memTNF when combined with sTNF. Kinetic studies with iodinated TNF revealed that 80M2 stabilizes ligand binding in terms of a prolonged receptor complex half-life (10). These data raised the hypothesis that the kinetics of receptor ligand complex formation and disintegration might at least in part determine intracellular signaling strength. Accordingly, transient binding of sTNF to TNFR2 would only allow formation of short

* This work was supported by Deutsche Forschungsgemeinschaft Grant GR 1307/3–3 and Sonderforschungsbereich 495, project AA. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

‡ To whom correspondence should be addressed. Tel.: 49-711-685-6987; Fax: 49-711-685-7484.

¹ The abbreviations used are: TNF, tumor necrosis factor; TNFR, TNF receptor; EGFP, enhanced green fluorescent protein; FADD, Fas-associated protein with death domain; JNK, c-Jun amino-terminal kinase; mAb, monoclonal antibody; MF, mouse fibroblast; RISC, receptor-induced signaling complex; TRAF, TNF receptor-associated factor; z-VAD-fmk, benzyloxycarbonyl-Val-Ala-Asp fluoromethyl ketone; TRADD, TNF receptor-associated death domain protein; memTNF, membrane-bound TNF; sTNF, soluble TNF; CHO, Chinese hamster ovary; PBS, phosphate-buffered saline; MOPS, 4-morpholinepropane-sulfonic acid.

living complexes that may inefficiently induce intracellular signals (10). On the other hand, the bioactivity of 80M2 is dependent on its dimeric IgG1 structure, since Fab fragments derived thereof are inactive (data not shown). These data raised the possibility that secondary clustering of TNF·TNFR2 complexes by 80M2 might reflect the underlying mechanism by which memTNF gains superior signaling capability on TNFR2.

Differences in the bioactivity of soluble and membrane-bound forms of other members of the TNF ligand family have also been described (e.g. for Fas ligand (FasL) (11), CD40L (12), and TRAIL (13)). Soluble FasL can even exert antiapoptotic activity by serving as an antagonist for the membrane-bound form of FasL (11).

In this study, we have analyzed receptor chimeras derived from the extracellular domains of the two TNF receptors and the intracellular domain of Fas. TNFR2-Fas chimera, comprising the extracellular domain of TNFR2 and the cytoplasmic part of Fas, strongly induced apoptosis only after treatment with memTNF-like stimuli but not with sTNF. In contrast, both ligand forms induced a strong apoptotic signal in TNFR1-Fas chimeras. These data show that the individual responsiveness of TNFR1 and TNFR2 for soluble or membrane-bound TNF can be transferred to a distinct intracellular signaling system, indicating that responsiveness is dominantly controlled by the extracellular domains.

EXPERIMENTAL PROCEDURES

Cell Culture and Reagents—HeLa cells stably transfected with human TNFR2 (HeLa80) or TNFR2 plus Fas (HeLa80Fas) (14, 15) and Chinese hamster ovary (CHO) cells expressing TNF Δ 1–12 (CHO_{TNF Δ 1–12}) (10) have been described elsewhere. Simian virus 40 large T-immortalized murine fibroblasts have been generated from TNFR1 and TNFR2 double knockout mice and were generously supplied by Daniela Männel (University of Regensburg, Regensburg, Germany). HeLa cells, CHO cells, and immortalized mouse fibroblasts were grown in RPMI 1640 medium supplemented with 5% (v/v) heat-inactivated fetal calf serum and 2 mM L-glutamine. 1 μ g/ml puromycin A was added once a week to TNFR1-Fas and TNFR2-Fas expressing mouse fibroblasts routinely. Kym-1 cells were grown in Clicks-RPMI medium containing 10% fetal calf serum and 2 mM L-glutamine (10). Recombinant human TNF (2 \times 10⁷ units/mg) was provided by Knoll AG (Ludwigshafen, Germany). Cys-TNF and mutants derived thereof (Cys-TNF143N/145R and Cys-TNF32W/36T) have been generated in *Escherichia coli* and purified to homogeneity using a nickel-chelate column.² The monoclonal mouse antibody 80M2 (16), H398 (17), and Htr9 (18) have been described. The TNFR2-specific monoclonal antibody MR2-1 was kindly provided by W. Buurman (University of Limburg, Maastricht, The Netherlands). Additional antibodies specific for TNFR2 were purchased from R&D (goat anti-huTNFR2), TRAF2-specific antibodies (mouse anti-huTRAF2, clone C90-481) were from Pharmingen, and antibodies specific for JNK (rabbit anti-huJNK; C-17) were from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA). z-VAD-fmk was purchased from Bachem. The expression plasmids pFADD-EGFP and pCaspase-8-EGFP were kind gifts from Michael Lenardo (National Institutes of Health, Bethesda, MD), and pTRAF2-EGFP was from Harald Wajant (University of Stuttgart, Stuttgart, Germany), and they have been described elsewhere (19, 20).

Measurement of the Metabolic Activity by Microphysiometry—Description and operation of the Cytosensor microphysiometer (Molecular Devices Corp., Sunnyvale, CA) have appeared elsewhere (21, 22). HeLa80Fas (3 \times 10⁵ cells) were seeded into chambers (Molecular Devices). The cell capsule was placed into a flow- and temperature (37 °C)-regulated sensing chamber of the microphysiometer, and pH changes were monitored. Cells were perfused with low phosphate-buffered RPMI medium (Irvine Scientific, Irvine, CA) in a cyclic manner. The pump cycle was 120 s, comprising a flow-on period (100 μ l/min for 80 s) followed by a flow-off period (40 s). The extracellular acidification rate was mathematically normalized to 100% at the data point just prior to stimulation with TNF143N/145R (150 ng/ml) (basal acidification rate). The addition of mAb 80M2 (2 μ g/ml) was per-

formed 30 min prior to treatment with TNF mutants and extended to the whole TNF stimulation time of 30 min.

Generation of Stably Transfected Cell Lines—Expression constructs encoding the fusion proteins TNFR1-Fas and TNFR2-Fas were generated by PCR cloning. A *Kpn*I site was introduced by silent mutagenesis into the coding region of pBS-TNFR1 and pBS-TNFR2 at bp 704 and 899, respectively, 3' of the potential transmembrane region. In addition, the cytoplasmic region of Fas was amplified by PCR introducing 5' and 3' appropriate restriction sites for TNFR ligation (pBS TNFR1-Fas and pBS TNFR2-Fas). TNFR1-Fas and TNFR2-Fas were subcloned into the expression vector pEF PGKpuro (23), using *Bam*HI and *Eco*RV, generating pEFpuroTNFR1-Fas and pEFpuroTNFR2-Fas. All constructs generated by PCR were verified by sequencing. Immortalized mouse fibroblasts (4 \times 10⁵ cells) from TNFR1 and TNFR2 double knockout cells were transfected with pEFpuroTNFR1-Fas or pEFpuroTNFR2-Fas using LipofectAMINE Plus (Invitrogen) according to the manufacturer's recommendations. The day after, cells were selected for stably expressing cells by 1–5 μ g/ml puromycin A, and 2 weeks later they were sorted for TNFR1-Fas- and TNFR2-Fas-positive cells by flow cytometry using a FACStar⁺ (Becton Dickinson, San Jose, CA). Briefly, 5 \times 10⁵ cells were harvested and resuspended in PBA (0.025% bovine serum albumin, 0.02% Na₂S₂O₃ in PBS) containing mouse 5 μ g/ml anti-huTNFR1 (Htr 9) or mouse anti-TNFR2 antibodies (MR2-1). After incubation for 1 h at 4 °C, cells were washed once with PBA, resuspended in PBA containing secondary fluorescein isothiocyanate-labeled goat anti-mouse IgG plus IgM antibodies (Dianova, Germany), and incubated at least for 30 min at 4 °C. Cells were washed again and subjected to fluorescence-activated cell sorting. 10,000–30,000 positive cells were collected and grown in cell culture medium containing 1 μ g/ml puromycin A. TNFR1-Fas and TNFR2-Fas expressing mouse fibroblasts (MF-R1-Fas and MF-R2-Fas cells, respectively) were stable for at least 3 weeks.

Cell Death Assays—Mouse fibroblasts (1.5 \times 10⁴ cells/well) were grown in 96-well plates overnight. Cells were then treated as indicated and cultivated overnight. The next day, cells were washed three times with PBS followed by crystal violet staining (20% methanol, 0.5% crystal violet) for 15 min. The wells were washed with H₂O and air-dried. The dye was resolved with methanol for 15 min, and optical density at 550 nm was determined with an enzyme-linked immunosorbent assay plate reader (SPECTRAMax 340PC; Molecular Devices).

For coculture experiments, CHO cells (1.5 \times 10⁵) were seeded into a 12-well plate. The next day, MF-R2-Fas cells (1.5 \times 10⁵) were given on top and cultivated at 37 °C. Cells were visualized by light microscopy, and pictures were taken 1 h after seeding of the mouse fibroblasts.

Coimmunoprecipitation and Western Blotting—HeLa80Fas cells (5 \times 10⁶ cells) were pretreated where indicated with antagonistic TNFR1-Fab fragment (H398-Fab) for 30 min followed by stimulation with sTNF (100 ng/ml) or Cys-TNF (116 ng/ml). After incubation at 37 °C for the indicated times, cells were washed with ice-cold PBS and scraped off the plate in PBS. Cells were pelleted and lysed in 300 μ l of lysis buffer (20 mM Tris, pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 30 mM NaF, 0.5 mM phenylmethylsulfonyl fluoride, protease inhibitor mix (Roche Molecular Biosciences). Proteins were extracted by vortexing three times for 40 s, and TNFR2 complexes were immunoprecipitated for 3 h at 4 °C while tumbling. Immune complexes were washed in lysis buffer and assessed by Western blotting using TRAF2-specific antibodies. Proteins were visualized by chemiluminescence (Super Signal; Pierce).

Electrophoretic Mobility Shift Assays of NF- κ B Activation—5 \times 10⁵ cells/well HeLa80Fas cells or mouse fibroblasts were seeded in six-well plates and grown overnight. Where indicated, cells were pretreated with 2 μ g/ml mAb 80M2 for 30 min. The cells were then stimulated for various times with the indicated reagents. Nuclear extracts were prepared as described (24), and samples were adjusted for identical protein levels. As probe, [³²P]ATP-end-labeled NF- κ B-specific oligonucleotides (5'-AGTTGAGGGGACTTCCAGGC-3') were used.

Immunocomplex JNK Assay—JNK assays were performed basically as described (25). Briefly, following stimulation, cells (5 \times 10⁶ cells) were lysed in kinase lysis buffer (20 mM Tris, pH 7.4, 5 mM MgCl₂, 1% Triton X-100, 150 mM NaCl, 0.1 mM phenylmethylsulfonyl fluoride, 1 mM NaVO₄, and 1 mM NaF) by sonification. JNK was immunoprecipitated with 0.6 μ g of JNK-specific antiserum (Santa Cruz Biotechnology) and subjected to kinase assays using GST-Jun-(5–89) (0.5 μ g/assay) and 5 μ Ci of [³²P]ATP as substrate. The reactions were carried out in assay buffer (20 mM MOPS, pH 7.2, 10 mM EGTA, 2 mM MgCl₂, 0.1% Triton X-100, and 1 mM dithiothreitol) at 37 °C for 20 min.

Transient Transfections and Confocal Microscopy—Cells were harvested and transiently transfected with 10 μ g of expression plasmids of

² A. B. Hammer, J. Gerspach, P. Scheurich, and K. Pfizenmaier, manuscript in preparation.




FIG. 1. Stable expression of chimeric receptors in mouse fibroblasts. *A*, schematic representation of TNFR1-Fas and TNFR2-Fas chimeric proteins. The cytoplasmic domain of Fas, amino acids 191–335, was fused to the C terminus of the potential transmembrane region of TNFR1 (amino acids 236) or TNFR2 (amino acids 301). Amino acids in the fused region are indicated. *B*, immortalized mouse fibroblasts were stably transfected with TNFR1-Fas or TNFR2-Fas expression plasmids. Expression of the chimeras was analyzed by flow cytometry using TNFR1-specific (Htr9) and TNFR2-specific (MR2-1) antibodies. The percentage of TNFR-Fas-positive cells is indicated.

pFADD-EGFP and pTRAF2-EGFP, respectively, or with 4 μ g of pCaspase-8(C369S)-EGFP plus 4 μ g of murine pFADD by electroporation. Cells (800 μ l; 1.25×10^6 cells/ml) were electroporated at 1800 microfarad and 250 V in a 0.4-cm cuvette (Peqbio Easyject Plus; Peqlab). After electroporation, cells were immediately transferred into cell culture medium, and 3×10^5 cells/dish (35 mm; Maltek) were grown for 18 h before analysis. Electroporation with pCaspase-8(C369S)-EGFP required the addition of 20 μ M z-VAD-fmk. For live imaging TNF, mAb 80M2 and Cys-TNF143N/145R were coupled with AlexaFluor-546 (Molecular Probes, Inc., Eugene, OR) according to the manufacturer's instructions. Cells were washed in PBS and, where indicated, preincubated with AlexaFluor-546-coupled 80M2 (80M2_{red}) (6 μ g/ml) for 4 min at room temperature or 1.7 μ g/ml AlexaFluor-546-coupled sTNF (sTNF_{red}) on ice followed by two washings with cell culture medium without phenol red. Cells were placed into a chamber held constantly at 37 °C and 5% CO₂ and treated where indicated with 300 ng/ml TNFR2-specific TNF mutant (Cys-TNF143N/145R), and pictures were taken at the indicated time points. Alternatively, 10^5 cells were seeded onto coverslips in a 24-well plate and treated as above. Instead of performing live imaging of the cells, cells were washed with ice-cold PBS, fixed with 3.5% paraformaldehyde in PBS for 15 min at 37 °C at various time points, mounted with Fluoromount-G (Biozol, Germany) onto glass slides, and examined using a Leica DM IRBE confocal immunofluorescence microscope. Pictures were taken at a resolution of 1024 \times 1024 pixels with a magnification of \times 630 for live imaging and \times 1000 for fixed cells.

Binding Kinetics—Receptor-ligand studies were performed as described (26). Briefly, TNF was labeled with ¹²⁵I by the chloramine-T method. Murine fibroblasts were incubated with 0.2 nM ¹²⁵I-TNF in the presence or absence of mAb 80M2 (2 μ g/ml) for 1 h on ice. Cells were incubated for several time periods at 37 °C, and cell-bound ¹²⁵I-TNF was determined after centrifugation of the cells through a phthalate oil mixture.

RESULTS

Construction of Fibroblasts Expressing Chimeric TNF Receptor/Fas Proteins—To get a better understanding for the molecular basis of the differential signaling capacity of sTNF and memTNF, we constructed chimeras of Fas and TNFR1 and TNFR2, respectively. These fusion proteins consist of the extracellular and transmembrane region of the respective human TNFRs fused to the cytoplasmic domain of human Fas (Fig. 1A). Hybrid constructs TNFR1-Fas and TNFR2-Fas were stably expressed in large T antigen-immortalized fibroblasts de-

rived from TNFR1/TNFR2 double knockout animals, obtaining a cellular system devoid of any TNF background responsiveness. Fluorescence analyses of the TNFR1-Fas- and TNFR2-Fas-expressing mouse fibroblast cells, MF-R1-Fas and MF-R2-Fas, respectively, are shown in Fig. 1B. Equilibrium binding studies using iodinated TNF revealed ligand binding sites of 15,000 for MF-R1-Fas cells and 45,000 for MF-R2-Fas cells per cell (data not shown).

Induction of Apoptosis in MF-R1-Fas and MF-R2-Fas Cells—As expected, we could not detect any TNF responsiveness in the parental mouse fibroblasts devoid of both mouse TNF receptors (data not shown). MF-R1-Fas cells, however, developed a strong cytotoxic response after treatment with serial dilutions of sTNF (Fig. 2A). Development of cell death was nearly maximum at TNF concentrations as low as 1 ng/ml, and the majority of the cells showed typical blebbing already after 3 h of sTNF treatment (data not shown). Further, cell death could be blocked by the inhibitor z-VAD-fmk, demonstrating the involvement of caspases (data not shown). Wild type TNFR1 is equally well activated by sTNF and memTNF (10). We confirmed this for TNFR1-Fas using the TNF mutein Cys-TNF32W/86T, derived from a TNFR1-specific mutant of sTNF (27), which allows additional receptor cross-linking due to the formation of cysteine-linked multimers (Fig. 2A). The respective TNFR2-specific mutein of TNF, Cys-TNF143N/145R, did not induce apoptosis in MF-R1-Fas cells (Fig. 2A).

In a parallel set of experiments, we investigated the chimeric receptor TNFR2-Fas. MF-R2-Fas cells were treated with sTNF up to concentrations of 300 ng/ml, but no significant cytotoxic response could be observed after overnight culture (Fig. 2B). In the presence of the antibody 80M2, however, a strong cytotoxic response was observed with a half-maximum effect at a TNF concentration of about 100 pg/ml, whereas 80M2 on its own was not toxic (Fig. 2B). Cell death developed rapidly; most cells showed massive signs of disintegration after 1 h of memTNF-like stimulation (data not shown). Again, induction of cytotoxicity could be efficiently blocked with the pan caspase inhibitor z-VAD-fmk (data not shown). The TNFR2-selective derivative Cys-TNF143N/145R also induced a significant apoptotic re-




FIG. 2. Response pattern of TNFR1-Fas and TNFR2-Fas expressing cells to sTNF and memTNF. *A* and *B*, MF-R1-Fas (*A*) and MF-R2-Fas (*B*) cells were treated with serial dilutions of sTNF or TNFR1-specific (Cys-TNF32W/86T) and TNFR2-specific (Cys-TNF143N/145R) TNF mutants, respectively. For costimulation with the monoclonal antibody 80M2, cells had been preincubated with 2 μ g/ml 80M2 for 30 min at 37 $^{\circ}$ C before sTNF treatment. Cell viability was determined by crystal violet staining the next day. All experimental groups shown were performed in parallel, one representative experiment out of three is shown. *C*, wild type Chinese hamster ovary (CHO_w) cells (*left*) or CHO cells stably expressing a noncleavable form of memTNF (CHO_{TNFM1-12}) (*right*) were grown overnight. MF-R2-Fas cells were seeded on top, and induction of apoptosis was followed by light microscopy. Pictures were taken after 1 h of coculture. The percentage of apoptotic mouse fibroblasts was calculated after counting about 200 cells. Bars, 75 μ m. *D* and *E*, MF-R1-Fas (*D*) and MF-R2-Fas (*E*) cells were treated with TNF (100 ng/ml), Cys-TNF (112 ng/ml), without or after pretreatment with mAb 80M2 (2 μ g/ml) for 30 min as indicated. Cell lysates were prepared directly before (–) or after the indicated time points of TNF stimulation, and JNK activity was measured by immunocomplex kinase assay with GST-c-Jun(5–89) as a substrate (*upper panels*), or the translocation of NF- κ B was investigated from isolated nuclei by gel shift analysis (*lower panels*). The positive control (*pos. c*) corresponds to HeLa cells stimulated with TNF (100 ng/ml) for 30 min. The specificity was controlled by the addition of a 100-fold excess of unlabeled oligonucleotides (100 \times cold).

sponse, although with reduced efficacy when compared with sTNF plus 80M2 (Fig. 2*B*). In contrast, the TNFR1-specific mutein Cys-TNF32W/86T was ineffective (Fig. 2*B*). To confirm that the transmembrane form of TNF is also able to induce apoptosis in MF-R2-Fas cells, these were cocultured for 1 h with CHO cells expressing a mutant form of memTNF that cannot be processed by the tumor necrosis factor α -converting enzyme (10). These cells (Fig. 2*C*, *right panel*), but not control CHO cells (Fig. 2*C*, *left panel*), induced a strong cytotoxic response in TNFR2-Fas-expressing mouse fibroblasts. When apoptotic cells were counted after 3 h of coculture, <5% of apoptotic MF-R2-Fas cells were determined in cocultures with control CHO cells, whereas between 68 and 79% ($n = 3$) of MF-R2-Fas cells showed an apoptotic phenotype in cocultures

with memTNF-expressing CHO cells (data not shown). These values are in good agreement with the percentage of the MF-R2-Fas cells expressing high numbers of chimeric receptors as estimated from flow cytometry data (Fig. 1*B*). As expected, memTNF-expressing CHO cells were also toxic for MF-R1-Fas cells (data not shown). In summary, these results show that the divergent responsiveness of TNFR1 and TNFR2 to sTNF is independent of the cytoplasmic domain of the receptors, since it is transferable to the intracellular part of Fas.

Gene Induction Pathways Initiated by TNFR-Fas Chimeras—Although Fas represents the prototype of a death receptor, it is also known to activate gene expression (e.g. via the activation of the transcription factor NF- κ B and activation of mitogen-activated protein kinases) (25, 28). To investigate




FIG. 3. Differential RISC formation of TNFR-Fas chimeras. MF-R1-Fas (A and B) or MF-R2-Fas (C–H) cells were transiently transfected with constructs expressing human FADD-EGFP (A–F) or human caspase-8(C360S)-EGFP plus murine FADD in the presence of 20 μ M z-VAD-fmk (G and H). The day after, cells were pretreated with Alexa 546-labeled 80M2 (80M2_(red); 6 μ g/ml) (E–H) or sTNF (TNF_(red); 1.7 μ g/ml) (A–D) for 4 min on ice followed by washing with PBS. Subsequently, unlabeled sTNF (40 ng/ml) was added to the 80M2_(red)-treated cells, and cells were examined by live imaging (A–F) or fixed at the indicated time points (G and H) and examined using a confocal laser-scanning microscope.

whether the differential responsiveness of TNFR2-Fas to sTNF and memTNF also holds true for noncytotoxic Fas responses, we analyzed the activation of NF- κ B and JNK in both MF-R1-Fas and MF-R2-Fas cells after treatment with sTNF and memTNF-like stimuli. The activations of NF- κ B and JNK were investigated by electromobility shift assays and immunocomplex kinase assays, respectively, revealing an identical response pattern for both cellular responses (Fig. 2, D and E) as compared with the induction of apoptosis (Fig. 2, A and B). These results strongly suggest that the molecular basis of the difference in signal initiation by memTNF and sTNF is at or upstream of the formation of the receptor-induced signaling complex (RISC). In accordance with literature data, JNK activation induced by TNFR-Fas chimera could be blocked by z-VAD-fmk (25, 29, 30), implicating the dependence on the activation of caspases, whereas nuclear translocation of NF- κ B was rather augmented by this caspase inhibitor (data not shown).

Recruitment of FADD to TNF Receptor/Fas Chimeras—We then asked whether the observed differences in the signaling strength of the various ligand/receptor chimera combinations are mirrored at the level of RISC formation. One of the first intracellular reactions after ligand-induced oligomerization of Fas is the recruitment of FADD, leading to caspase-8 activation. We therefore monitored transiently expressed human FADD-EGFP in mouse fibroblast cell lines by confocal microscopy after stimulation with AlexaFluor-546-labeled TNF receptor agonists. Fig. 3 shows the overlays of the green and the red fluorescence observed. As expected, MF-R1-Fas cells transiently transfected with FADD-EGFP showed a green cytosolic fluorescence (Fig. 3A). These cells had been preincubated with Alexa-546-labeled sTNF, detectable as a weak red cell surface staining. No prominent colocalization with FADD-EGFP at the cell surface can be observed. After incubation of the cells for 15 min at 37 $^{\circ}$ C, however, the majority of the TNFR1-Fas-bound sTNF had formed clusters mostly colocalized with FADD-EGFP (Fig. 3B). These data demonstrate that sTNF is able to

recruit significant amounts of FADD-EGFP to TNFR1-Fas molecules, leading to the formation of large RISC aggregates.

In contrast, the respective sTNF treatment of MF-R2-Fas cells did not reveal a significant colocalization of sTNF and FADD-EGFP, and no signs of cell surface located cluster formation could be observed (Fig. 3, C and D). Stimulation for only 2–8 min with a memTNF-like agent, however, consisting of sTNF and the red fluorescent antibody 80M2, induced rapid formation of cell surface-associated clusters of colocalized TNFR2-Fas and FADD-EGFP (data not shown; see Fig. 3, E and F). Similar results were obtained using MF-R2-Fas cells transiently transfected with pCaspase-8(C360S)-EGFP, expressing a catalytic inactive caspase-8, and murine FADD expression constructs in the presence of z-VAD-fmk. Again, sTNF was able to induce recruitment of caspase-8 to TNFR1-Fas but not to TNFR2-Fas chimeras (data not shown). In contrast, memTNF-like stimuli were efficient in both cellular systems (data not shown; Fig. 3, G and H). Together, these data demonstrate that the different signaling capacities of the two TNF forms are directly reflected at the level of RISC formation.

Enhanced Recruitment of TRAF2 to Wild-type TNFR2 by memTNF—We next investigated whether a lack of significant adaptor protein recruitment to TNFR2-Fas by sTNF can also be observed in wild type TNFR2-positive cells. In HeLa cells stably overexpressing TNFR2 (HeLa80), transiently expressed TRAF2-EGFP was also primarily located in the cytosol (Fig. 4, upper panel). The addition of the TNFR2-selective sTNF mutant sTNF143N/145R, marked with a red fluorescing dye, revealed a staining of the plasma membrane but did not result in any visible recruitment of TRAF2-EGFP to the cell membrane (Fig. 4A, upper panel). Cellular stimulation with sTNF143N/145R in the presence of the antibody 80M2, however, resulted in the strong formation of membrane-associated TRAF2-EGFP aggregates showing colocalization with TNFR2 (Fig. 4A, lower panel). Significant TRAF2-EGFP recruitment could also be induced using the secondary cross-linked TNFR2-selective TNF mutein Cys-TNF143N/154R (data not shown). Similar




FIG. 4. Recruitment of TRAF2 by stimulation of TNFR2 with a memTNF-like agent but not with sTNF. *A*, HeLa cells, stably expressing TNFR2 (HeLa80), were transiently transfected with constructs expressing huTRAF2-EGFP. The day after, cells were pretreated with Alexa 546-labeled mAb 80M2 (6 μ g/ml) or TNFR2-specific TNF (sTNF143N/145R; 1.7 μ g/ml) for 4 min on ice followed by washing with PBS. Subsequently unlabeled membrane-like TNF (Cys-TNF143N/145R) was added to the mAb 80M2-treated cells. Live imaging was performed with a confocal microscope, and pictures were taken at the indicated times. *B*, HeLa cells stably expressing TNFR2 plus Fas (HeLa80Fas) were treated with sTNF (100 ng/ml) or Cys-TNF143N/145R (116 ng/ml) for the indicated times, followed by cell lysis. Where indicated, cells had been pretreated with antagonistic TNFR1-Fab fragments (α TNFR1-Fab; 14 μ g/ml) to prevent binding of sTNF to TNFR1 and subsequent TRAF2 recruitment. Lysates were subjected to coimmunoprecipitation with TNFR2-specific antibodies, and Western blot analysis was performed using TRAF2-specific antibodies. As controls, TNFR2-specific antibodies (control IgG) and lysates from cells overexpressing huTRAF2 (TRAF2) were used.

data were obtained using mouse fibroblasts transfected with wild type TNFR2 and TRAF2-EGFP (data not shown).

To study TRAF2 recruitment to TNFR2 after memTNF-like stimulation also at physiological TRAF2 levels, we assessed coimmunoprecipitation studies with endogenously expressed TRAF2. TNFR2 was immunoprecipitated from HeLa cells, stably expressing TNFR2, and the precipitates were investigated for TRAF2 by Western blotting. 15 and 30 min after stimulation of the cells with sTNF, only slightly enhanced TRAF2 amounts were detectable in the immunoprecipitates as compared with unstimulated cells (Fig. 4*B*). TRAF2 coimmunoprecipitation could not be blocked with a TNFR1-specific antagonistic Fab fragment (α TNFR1-Fab), capable of inhibiting TNF binding to TNFR1 (Fig. 4*B*). In parallel experiments, we used TNFR2-specific Cys-TNF143N/145R for a memTNF-like stimulation of TNFR2. Immunoprecipitates from these cells contained significantly larger amounts of TRAF2 as compared with that obtained from sTNF-treated cells (Fig. 4*B*), confirming the data obtained by confocal microscopy (Fig. 4*A*). Together, these results show that also in wild type TNFR2, the enhanced signaling capacity of memTNF-like stimuli, like Cys-TNF143N/145R, is linked to an enhanced recruitment of adaptor proteins and not dependent on overexpression of these intracellular signaling molecules.

We verified that enhanced adaptor recruitment to wild type





FIG. 5. Enhanced NF- κ B activation and metabolic activity induced by memTNF-like stimuli. *A*, KYM-1 cells (*top*) and HeLa cells expressing TNFR2 (HeLa80, *bottom*) were treated with various sTNF-like stimuli (sTNF, 30 ng/ml; TNFR1-specific sTNF32W/86T, 30 ng/ml; TNFR2-specific sTNF143N/145R, 300 ng/ml) or memTNF-like stimuli (Cys-TNF, 30 ng/ml; sTNF/80M2, 30 ng/ml/2 μ g/ml; TNFR1-specific Cys-TNF32W/86T, 30 ng/ml; TNFR2-specific Cys-TNF143N/145R, 300 ng/ml) for 30 min. An agonistic TNFR2-specific serum (M80, 1:200) and a rabbit control serum (1:200) were also included. Nuclear extracts were prepared, and 10 μ g of protein was subjected to gel shift analysis using 32 P-labeled NF- κ B-specific oligonucleotides. *B*, the metabolic activity of HeLa80Fas cells was analyzed using a microphysiometer. HeLa80Fas cells (3×10^5) had been pretreated with mAb 80M2 for 30 min at 37 $^{\circ}$ C where indicated. Cells were then stimulated with the TNFR2-specific TNF mutein sTNF143N/145R in the presence or absence of mAb 80M2 for 30 min (*arrow*). The change in the metabolic activity was followed over time and is expressed as percentage acidification.

TNFR2, as depicted in Fig. 4, *A* and *B*, is in fact paralleled by an enhanced cellular response. To this, HeLa80 cells and KYM-1 cells, known to activate NF- κ B after appropriate stimulation via TNFR2 (16, 20), were examined for the activation of NF- κ B by electrophoretic mobility shift assay. Nuclear translocation and DNA binding of NF- κ B was determined after 30 min of receptor stimulation (*i.e.* within the same time range also investigated in the confocal microscopy experiments). These experiments revealed a similar stimulatory capacity of TNFR1 and TNFR2 for both cell lines (Fig. 5*A*). The TNFR2-selective sTNF143N/145R on its own showed only a marginal, if any, capacity to activate NF- κ B, whereas the TNFR2-selective Cys-TNF143N/145R possessed an intermediate stimulatory capacity. This confirms that in HeLa and KYM-1 cells a memTNF-like activity is mandatory for full activation of TNFR2, whereas sTNF is sufficient to fully activate TNFR1. Finally, in TNFR2-expressing HeLa cells the overall metabolic response to a selective stimulation of TNFR2 by sTNF in the absence and presence of 80M2 was determined using a microphysiometer. The results confirm a strong cellular (*i.e.* metabolic) response initiated by TNFR2 only when stimulated with sTNF143N/145R in the presence of 80M2 (Fig. 5*B*).

Ligand Association and Dissociation Studies—In a previous

Fig. 6. Time course of TNF dissociation at 37 °C. MF-R1-Fas (A) and MF-R2-Fas (B) cells were incubated for 1 h with 10 ng/ml 125 I-TNF at 4 °C in the presence or absence of mAb 80M2 (2 μ g/ml). Dissociation of the radiolabeled ligand was measured at 37 °C in the presence of 2 μ g/ml unlabeled sTNF. Non-specific binding determined in the presence of a 200-fold excess of unlabeled sTNF was less than 5% of total binding and has been subtracted. Half-lives of the TNF-receptor complexes were calculated from exponential decay curves.



publication, we showed that the enhanced signaling capacity of sTNF in the presence of 80M2, representing a memTNF-like stimulus, correlates with the stabilization of ligand-receptor complexes due to a strongly reduced dissociation rate (10). Using MF-R1-Fas and MF-R2-Fas cells, respectively, association and dissociation studies with iodinated sTNF at 37 °C were performed. The association kinetics of iodinated sTNF to both receptors were rapid and similar, with half-maximum binding after about 2 min for TNFR1-Fas and about 1 min for TNFR2-Fas at a ligand concentration of 0.3 nM (data not shown). The presence of the antibody 80M2 did not significantly change ligand association kinetics of TNFR2 ($k_{on}(sTNF) = 3.0 \cdot 10^9 \text{ M}^{-1} \text{ min}^{-1}$ and $k_{on}(sTNF + 80M2) = 2.5 \cdot 10^9 \text{ M}^{-1} \text{ min}^{-1}$). To study ligand dissociation, untreated MF-R1-Fas and MF-R2-Fas cells, as well as 80M2-pretreated MF-R2-Fas cells, were incubated with 0.2 nM iodinated sTNF for 1 h on ice to allow receptor complex formation. The temperature was then shifted to 37 °C, and in the presence of a 200-fold excess of unlabeled sTNF, the release of the radiolabeled ligand was followed. Fig. 6A shows that sTNF dissociates only slowly from MF-R1-Fas cells (half-life = 32 min). In contrast, sTNF-TNFR2-Fas complexes have a low half-life of about 1.2 min, whereas 80M2 pretreatment stabilizes ligand-TNFR2-Fas complexes more than 10-fold (half-life = 14.5 min). These association and dissociation data are in good agreement with the results obtained with wild type TNF receptors (10). Accordingly, also the dissociation constants (K_d values) at 37 °C, as calculated for the chimeric receptors, are very similar to those values determined for the wild type TNF receptors (29 μ M for TNFR1-Fas, 19 μ M for TNFR1; 286 μ M for TNFR2-Fas, 420 μ M for TNFR2) (26). In summary, we show that the stability of ligand-receptor complexes correlates with the formation of the RISC and signal capacities of TNFR1-Fas and TNFR2-Fas chimeras.

DISCUSSION

In contrast to TNFR1, which is equally well activated by both sTNF and memTNF, TNFR2 can only be efficiently activated by memTNF. To address the molecular mechanisms underlying these different bioactivities, we have constructed receptor chimeras containing the extracellular domains of the TNF receptors fused to the intracellular part of Fas. In the present study, we demonstrate that these chimeric receptors exhibit identical activation requirements regarding the two TNF forms as the wild type TNF receptors. Moreover, our data suggest that the half-life of TNF-TNFR complexes becomes translated into the efficiency of intracellular adaptor recruitment, thus controlling the intensity of the transmitted signal.

Experimental systems quantitatively assessing the signal capacity of cell surface expressed memTNF are difficult to handle. We therefore used available tools that mimic memTNF signaling. We have recently described the TNFR2-specific monoclonal antibody 80M2 that, in combination with sTNF, induces intracellular signals comparable with the transmem-

brane form of TNF (10). Furthermore, we have utilized mutants of sTNF that form intermolecular disulfide bonds via an N-terminal cysteine residue (Cys-TNF and the receptor-specific derivatives Cys-TNF32W/86T and Cys-TNF143N/145R), resulting in the formation of secondary cross-linked TNF trimers. These TNF derivatives also show an enhanced signaling capacity via TNFR2 as compared with sTNF (Fig. 2B).²

To obtain a suitable molecular system for investigation of sTNF and memTNF action, we transferred the extracellular domains of the two TNF receptors to the cytoplasmic part of Fas, another TNF receptor family member. A simple exchange of the intracellular TNF receptor domains turned out to be inappropriate due to the constitutive cytotoxic activity of the intracellular region of TNFR1 (data not shown). To omit problems with endogenous TNF responsiveness, receptor chimeras were expressed in a fibroblast-derived cell line from TNFR1/TNFR2 double knockout mice. As expected, both chimeric receptors, TNFR1-Fas and TNFR2-Fas, could be expressed in quite high numbers in these cells, with about 15,000 and 45,000 TNF binding sites for MF-R1-Fas and -R2-Fas cells, respectively (data not shown). In the absence of any TNF receptor-specific stimulus, both cell lines proliferate well without indications of spontaneous apoptosis (data not shown).

In various studies on TNFR1- and Fas-expressing cells, conflicting results have been obtained regarding the involvement of membrane rafts or the requirement of internalization for signal initiation. Recently, the possible arrangement of Fas in lipid rafts was shown (31, 32), which may be true for type II but not for type I cells (9, 33). Furthermore, Fas signaling has been reported to be independent of receptor complex internalization (8, 9), whereas TNFR1 was not (8). In the present study, both TNFR-Fas chimeras have been expressed in reasonable receptor numbers in mouse fibroblasts and are therefore supposed to induce a very strong initial signal upon appropriate stimulation. This is in accordance with the rapid TNF-induced morphological changes, typical for apoptosis, within 1 (TNFR2-Fas) to 3 h (TNFR1-Fas). In addition, pretreatment of MF-R2-Fas cells with methyl- β -cyclodextrine, which disrupts lipid rafts by cholesterol depletion (34), or monodansylcadaverine, which blocks receptor internalization (8), did not affect cell death kinetics of MF-R2-Fas cells treated with sTNF plus 80M2 in a 3-h assay (data not shown). In agreement, using radioiodinated sTNF, we did not find significant internalization of ligand-receptor complexes within 30 min of incubation at 37 °C (<5% of TNFR1-Fas when stimulated with sTNF; <10% of TNFR2-Fas when stimulated with sTNF with or without 80M2; data not shown). Together, all of these data suggest that our cellular model displays a very rapid apoptotic response after appropriate TNF treatment that is largely independent of secondary processes following RISC formation and/or cofactors.

The Studies Performed with the TNFR-Fas Chimeras Allow Two Direct Conclusions—First, the differential response pattern of the two TNF receptors to sTNF and memTNF could be

fully transferred to the Fas signaling system (*i.e.* in the case of TNFR2 from a gene inductive pathway, acting via TRAF2 binding, to the apoptotic pathway of Fas, acting via FADD mediated caspase-8 activation). Identical patterns were found for three different cellular responses (*i.e.* the induction of apoptosis, activation of NF- κ B, and activation of the mitogen-activated protein kinase JNK) (Fig. 2). These results clearly show that the responsiveness of the TNF receptors to the soluble *versus* the membrane bound form of TNF is independent of the particular intracellular signaling machinery. A direct consequence is that the decisive process, able to distinguish between sTNF and memTNF in the case of TNFR2 and TNFR2-Fas, is located upstream of the recruitment of cytoplasmic factors. This strongly argues for the existence of a general principle able to control the signaling strength of a given receptor within the TNF receptor family, which is not determined solely by the affinity of ligand/receptor interaction, since sTNF acting at saturating concentrations on TNFR2-Fas elicits only weak, if any, responses (Fig. 2, B and E).

Second, exogenously initiated cross-linking of ligand-receptor complexes is not necessary for induction of full signaling and receptor cluster formation. Cross-linking reagents like antibodies are commonly used for the efficient signal induction of Fas, TNFR2, or CD40 (11, 12, 16). Due to their multivalent nature, the treatment of cells with antibodies is paralleled by the formation of large receptor clusters (32, 35). However, the functional role of these clusters for signal initiation and strength has not been fully elucidated. In our studies, the efficient recruitment of FADD and TRAF2, respectively, is also paralleled by the formation of large receptor clusters, visible by colocalization of receptors with EGFP-tagged adaptor proteins (Figs. 3 (B, F, and H) and 4A) and in coimmunoprecipitation studies (Fig. 4B). This cluster formation always correlated with the particular signaling strength. In the case of TNFR2 and the TNFR2-derived Fas chimera, exogenously initiated cross-linking of ligand-receptor complexes was necessary for induction of full signaling and receptor cluster formation (Figs. 3 (F and H) and 4A). However, sTNF on its own is sufficient to initiate a strong intracellular signal via TNFR1-Fas, whereas in parallel triggering the formation of large receptor clusters (Figs. 2A and 3B). Since there are no observable differences in the efficiency of sTNF and memTNF upon TNFR1 stimulation (10),³ the results argue against a mandatory role of an external, additional cross-linking agent for full activation of a given receptor. Recent observations by the group of Peter, demonstrating that formation of large Fas clusters is not directly dependent on the cross-linking properties of the stimulating agent (9), are consistent with these observations.

As discussed above, cluster formation occurs in parallel with enhanced signaling and is independent of the cytoplasmic part of the TNF receptors. It is therefore likely that the extracellular parts of ligand-bound receptors mediate cluster formation via additional interactions possibly involving supplementary molecules. Since the TNFR-Fas chimeras used in our studies contain the transmembrane parts of the respective TNF receptors, the possibility cannot be excluded that these domains are involved in receptor aggregate formation. When chimeric fusion proteins comprising the extracellular domain of the erythropoietin receptor and the intracellular part from TNFR2 were investigated, an exchange of the respective transmembrane domains had moderate effects on the signaling capacity (36). However, the present structural data of ligand trimerized receptors indicate that the transmembrane domains are unlikely

to directly interact with each other between individual complexes (37, 38). More likely, a direct interaction of the ligated extracellular receptor domains might occur. Based on the dimeric crystal structure of the extracellular part of TNFR1, such a cluster formation has been already proposed by Naismith *et al.* (39). A possible candidate for receptor/receptor interaction is the recently defined preligand assembly domain, identified in both TNF receptors and in Fas, which is most likely present also in other members of the TNF receptor family (40).

What mechanisms beside additional cross-linking of receptor complexes could determine the enhanced signaling capability of memTNF? We propose that the stability of individual receptor-ligand complexes could be the important factor. Interactions of sTNF with TNFR1 and TNFR2 differ strongly in this regard. Whereas sTNF forms very stable complexes with TNFR1 (half-life = 33 min) (10) and TNFR1-Fas (half-life = 32 min) (Fig. 6A), ligation of TNFR2 (half-life = 1.1 min) (10) and TNFR2-Fas (half-life = 1.2 min) (Fig. 6B) occurs only very transiently. These differences are most likely inherent properties of the TNF/TNF receptor interactions rather than caused by different stoichiometries of complexes or subsequent steps like receptor cluster formation. This is evident from studies with TNF receptor-derived IgG fusion proteins that revealed very similar results (41). The exchange rates of radiolabeled sTNF complexed with TNFR-IgG fusion proteins showed a half-life of about 7 min for TNFR2-derived complexes, whereas TNFR1 complexes were extremely stable (half-life = 8 h) (41). Accordingly, the stability of individual ligand receptor complexes, most likely comprising three receptor molecules, would control subsequent steps leading to the formation of large, stable clusters, capable of effectively recruiting adaptors and initiating a strong intracellular signal. Certain conditions must be met, however. First, the initial sTNF binding must be rapid as compared with the subsequent steps. This holds true, since sTNF binding occurs very fast (26) and is mainly controlled by diffusion.³ Second, receptor cluster formation should be dependent on preformed ligand-receptor complexes and should not occur spontaneously with unligated receptors. Most likely, this also holds true, since we observe no receptor patches directly after ligand binding on ice in our microscopy studies (Figs. 3 and 4). Third, formation of large receptor clusters must stabilize ligand binding in TNFR2 and TNFR2-Fas molecules. This also seems very reasonable, since integration of single receptor-ligand complexes into a lattice-like structure should inhibit dissociation of individual ligand-receptor complexes. We therefore suggest a causal relationship between the stability of individual receptor complexes and the efficiency of adaptor protein recruitment, the latter being directly translated into signaling strength. This principle might hold true for many, if not all, members of the TNF receptor family but is not necessarily restricted to these molecules.

Acknowledgments—We thank I.-W. von Broen (Knoll AG, Ludwigshafen, Germany) for recombinant TNF, D. Männel for murine fibroblasts, M. Lenardo for pCaspase-8-EGFP and pFADD-EGFP, H. Wajant for pTRAF2-EGFP, and A. Strasser (The Walter and Eliza Hall Institute, Melbourne, Australia) for pEF PGKpuro. We also thank K. Pfizenmaier and H. Wajant for helpful discussion.

REFERENCES

- Locksley, R. M., Killeen, N., and Lenardo, M. J. (2001) *Cell* **104**, 487–501
- Grell, M., and Scheurich, P. (1997) *Growth Factors and Cytokines in Health and Disease* (LeRoith, D., and Bondy, G. P., eds) pp. 669–726, JAI Press Inc., Greenwich, CT
- Hsu, H., Shu, H. B., Pan, M. C., and Goeddel, D. V. (1996) *Cell* **84**, 299–308
- Wajant, H., and Scheurich, P. (2001) *Int. J. Biochem. Cell Biol.* **33**, 19–32
- Pennica, D., Lam, V. T., Mize, N. K., Weber, R. F., Lewis, M., Fendly, B. M., Lipari, M. T., and Goeddel, D. V. (1992) *J. Biol. Chem.* **267**, 21172–21178
- Higuchi, M., and Aggarwal, B. B. (1994) *J. Immunol.* **152**, 3550–3558
- Dri, P., Gasparini, C., Menegazzi, R., Cramer, R., Alberi, L., Presani, G., Garbisa, S., and Patriarca, P. (2000) *J. Immunol.* **165**, 2165–2172
- Schutze, S., Machleidt, T., Adam, D., Schwandner, R., Wiegmann, K., Kruse,

³ A. Krippner-Heidenreich, F. Tübing, S. Bryde, S. Willi, G. Zimmermann, and P. Scheurich, unpublished data.

- M. L., Heinrich, M., Wickel, M., and Kronke, M. (1999) *J. Biol. Chem.* **274**, 10203-10212
9. Algeciras-Schimmich, A., Shen, L., Barnhart, B. C., Murnann, A. E., Burkhardt, J. K., and Peter, M. E. (2002) *Mol. Cell. Biol.* **22**, 207-220
 10. Grell, M., Douni, E., Wajant, H., Lohden, M., Clauss, M., Maxeiner, B., Georgopoulos, S., Lesslauer, W., Kollias, G., and Pfizenmaier, K., and Scheurich, P. (1995) *Cell* **83**, 793-802
 11. Schneider, P., Holler, N., Bodmer, J. L., Hahne, M., Frei, K., Fontana, A., and Tschopp, J. (1998) *J. Exp. Med.* **187**, 1205-1213
 12. Pound, J. D., Challa, A., Holder, M. J., Armitage, R. J., Dower, S. K., Fanslow, W. C., Kikutani, H., Paulie, S., Gregory, C. D., and Gordon, J. (1999) *Int. Immunol.* **11**, 11-20
 13. Wajant, H., Moosmayer, D., Wuest, T., Bartke, T., Gerlach, E., Schonherr, U., Peters, N., Scheurich, P., and Pfizenmaier, K. (2001) *Oncogene* **20**, 4101-4106
 14. Weiss, T., Grell, M., Hessabi, B., Bourteele, S., Muller, G., Scheurich, P., and Wajant, H. (1997) *J. Immunol.* **158**, 2398-2404
 15. Weiss, T., Grell, M., Sieminski, K., Muhlenbeck, F., Durkop, H., Pfizenmaier, K., Scheurich, P., and Wajant, H. (1998) *J. Immunol.* **161**, 3136-3142
 16. Grell, M., Scheurich, P., Meager, A., and Pfizenmaier, K. (1993) *Lymphokine Cytokine Res.* **12**, 143-148
 17. Thoma, B., Grell, M., Pfizenmaier, K., and Scheurich, P. (1990) *J. Exp. Med.* **172**, 1019-1023
 18. Brockhaus, M., Schoenfeld, H. J., Schlaeger, E. J., Hunziker, W., Lesslauer, W., and Loetscher, H. (1990) *Proc. Natl. Acad. Sci. U. S. A.* **87**, 3127-3131
 19. Siegel, R. M., Martin, D. A., Zheng, L., Ng, S. Y., Bertin, J., Cohen, J., and Lenardo, M. J. (1998) *J. Cell Biol.* **141**, 1243-1253
 20. Fotin-Mleczek, M., Henkler, F., Samel, D., Reichwein, M., Hausser, A., Parmryd, I., Scheurich, P., Schmid, J. A., and Wajant, H. (2002) *J. Cell Sci.* **115**, 2757-2770
 21. Parce, J. W., Owicki, J. C., Kereso, K. M., Sigal, G. B., Wada, H. G., Muir, V. C., Bousse, L. J., Ross, K. L., Sikic, B. I., and McConnell, H. M. (1989) *Science* **246**, 243-247
 22. Parce, J. W., Owicki, J. C., and Kereso, K. M. (1990) *Ann. Biol. Clin. (Paris)* **48**, 639-641
 23. Huang, D. C., Cory, S., and Strasser, A. (1997) *Oncogene* **14**, 405-414
 24. Dignam, J. D., Lebovitz, R. M., and Roeder, R. G. (1983) *Nucleic Acids Res.* **11**, 1475-1489
 25. Muhlenbeck, F., Haas, E., Schwenzer, R., Schubert, G., Grell, M., Smith, C., Scheurich, P., and Wajant, H. (1998) *J. Biol. Chem.* **273**, 33091-33098
 26. Grell, M., Wajant, H., Zimmermann, G., and Scheurich, P. (1995) *Proc. Natl. Acad. Sci. U. S. A.* **95**, 570-575
 27. Loetscher, H., Stueber, D., Banner, D., Mackay, F., and Lesslauer, W. (1993) *J. Biol. Chem.* **268**, 26350-26357
 28. Ahn, J. H., Park, S. M., Cho, H. S., Lee, M. S., Yoon, J. B., Vilcek, J., and Lee, T. H. (2001) *J. Biol. Chem.* **276**, 47100-47106
 29. Lenczowski, J. M., Dominguez, L., Eder, A. M., King, L. B., Zacharechuk, C. M., and Ashwell, J. D. (1997) *Mol. Cell. Biol.* **17**, 170-181
 30. Deak, J. C., Cross, J. V., Lewis, M., Qian, Y., Parrott, L. A., Distelhorst, C. W., and Templeton, D. J. (1993) *Proc. Natl. Acad. Sci. U. S. A.* **95**, 5595-5600
 31. Grassme, H., Schwarz, H., and Gulbins, E. (2001) *Biochem. Biophys. Res. Commun.* **284**, 1016-1030
 32. Grassme, H., Jekle, A., Riehle, A., Schwarz, H., Berger, J., Sandhoff, K., Kolesnick, R., and Gulbins, E. (2001) *J. Biol. Chem.* **276**, 20539-20596
 33. Scaffidi, C., Fulda, S., Srinivasan, A., Friesen, C., Li, F., Tomaselli, K. J., Debatin, K. M., Kramer, P. H., and Peter, M. E. (1998) *EMBO J.* **17**, 1675-1687
 34. Simons, K., and Toomre, D. (2000) *Nat. Rev. Mol. Cell. Biol.* **1**, 31-39
 35. Grassme, H., Jendrosseck, V., Bock, J., Riehle, A., and Gulbins, E. (2002) *J. Immunol.* **168**, 298-307
 36. Declercq, W., Vandenaebelle, P., and Fiers, W. (1995) *Cytokine* **7**, 701-709
 37. Park, Y. C., Ye, H., Hsia, C., Segal, D., Rich, R. L., Liou, H. C., Myszka, D. G., and Wu, H. (2000) *Cell* **101**, 777-787
 38. Ye, H., Park, Y. C., Kreishman, M., Kieff, E., and Wu, H. (1999) *Mol. Cell* **4**, 321-330
 39. Naismith, J. H., Devine, T. Q., Brandhuber, B. J., and Sprang, S. R. (1995) *J. Biol. Chem.* **270**, 13303-13307
 40. Chan, P. K., Chun, H. J., Zheng, L., Siegel, R. M., Bui, K. L., and Lenardo, M. J. (2000) *Science* **288**, 2351-2354
 41. Evans, T. J., Moyes, D., Carpenter, A., Martin, R., Loetscher, H., Lesslauer, W., and Cohen, J. (1994) *J. Exp. Med.* **180**, 2173-2179


Material Safety Data Sheet



Stratagene BL21 Competent Cells, Catalog # 200133

1. Product and company identification

Product name	: Stratagene BL21 Competent Cells, Catalog # 200133
Part No.	: BL21 competent cells 200133-41 pUC18 Control Plasmid 200231-42 DNA 1.42 M 2-Mercaptoethanol 210200-43
Manufacturer / Supplier	: Agilent Technologies, Inc. 1834 State Highway 71 West Cedar Creek, TX 78612
Emergency telephone number	: 1-800-894-1304
Use of the substance/preparation	: Chemical Kit
Validation date	: 11/21/2008



2. Hazards identification

Physical state	: BL21 competent cells Liquid. pUC18 Control Plasmid Liquid. DNA 1.42 M 2-Mercaptoethanol Liquid.
Odor	: BL21 competent cells Not available. pUC18 Control Plasmid Not available. DNA 1.42 M 2-Mercaptoethanol Not available.
OSHA/HCS status	: BL21 competent cells This material is considered hazardous by the OSHA Hazard Communication Standard (29 CFR 1910.1200). pUC18 Control Plasmid While this material is not considered hazardous by the OSHA Hazard Communication Standard (29 CFR 1910.1200), this MSDS contains valuable information critical to the safe handling and proper use of the product. This MSDS should be retained and available for employees and other users of this product. DNA 1.42 M 2-Mercaptoethanol This material is considered hazardous by the OSHA Hazard Communication Standard (29 CFR 1910.1200).
Emergency overview-Signal Word	: WARNING !
Emergency overview-Label Statement	: BL21 competent cells HARMFUL IF SWALLOWED. CONTAINS MATERIAL THAT MAY CAUSE TARGET ORGAN DAMAGE, BASED ON ANIMAL DATA. pUC18 Control Plasmid NOT EXPECTED TO PRODUCE SIGNIFICANT ADVERSE HEALTH EFFECTS WHEN THE RECOMMENDED INSTRUCTIONS FOR USE ARE FOLLOWED. DNA 1.42 M 2-Mercaptoethanol HARMFUL IF SWALLOWED. CAUSES EYE AND SKIN IRRITATION. BL21 competent cells Toxic if swallowed. Avoid exposure - obtain special instructions before use. Do not breathe vapor or mist. Do not ingest. Avoid contact with eyes, skin and clothing. Contains material that may cause target organ damage, based on animal data. Wash thoroughly after handling. pUC18 Control Plasmid No known significant effects or critical hazards. Avoid prolonged contact with eyes, skin and clothing. DNA 1.42 M 2-Mercaptoethanol Toxic if swallowed. Irritating to eyes and skin. Do not breathe vapor or mist. Do not ingest. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling.

2. Hazards identification

Routes of entry	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Eye contact. Inhalation. Ingestion. Eye contact. Ingestion. Eye contact. Ingestion.
Potential acute health effects		
Eyes	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	No known significant effects or critical hazards. No known significant effects or critical hazards. Irritating to eyes.
Skin	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	No known significant effects or critical hazards. No known significant effects or critical hazards. Irritating to skin.
Inhalation	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	No known significant effects or critical hazards. No known significant effects or critical hazards. No known significant effects or critical hazards.
Ingestion	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Toxic if swallowed. No known significant effects or critical hazards. Toxic if swallowed.
Medical conditions aggravated by over-exposure	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Repeated or prolonged exposure to the substance can produce target organs damage. Not applicable. Repeated skin exposure can produce local skin destruction or dermatitis. Repeated or prolonged contact with spray or mist may produce chronic eye irritation and severe skin irritation.
Over-exposure signs/symptoms	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Not applicable. Not applicable. Not applicable.

See toxicological information (section 11)

3. Composition/information on ingredients

<u>Name</u>	<u>CAS number</u>	<u>%</u>
BL21 competent cells		
Glycerol	56-81-5	5 - 10
Manganese dichloride	7773-01-5	5 - 10
Sucrose	57-50-1	5 - 10
Dimethyl sulfoxide	67-68-5	5 - 10
Potassium chloride	7447-40-7	1 - 5
1.42 M 2-Mercaptoethanol		
2-Mercaptoethanol	60-24-2	10

There are no ingredients or additional ingredients present which, within the current knowledge of the supplier and in the concentrations applicable, are classified as hazardous to health or the environment and hence require reporting in this section.

4 . First aid measures

Eye contact	: BL21 competent cells	In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if adverse health effects persist or are severe.
	pUC18 Control Plasmid DNA	In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if adverse health effects persist or are severe.
	1.42 M 2-Mercaptoethanol	In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if adverse health effects persist or are severe.
Skin contact	: BL21 competent cells	In case of contact, immediately flush skin with plenty of water. Remove contaminated clothing and shoes. Wash clothing before reuse. Clean shoes thoroughly before reuse. Get medical attention if adverse health effects persist or are severe.
	pUC18 Control Plasmid DNA	In case of contact, immediately flush skin with plenty of water. Remove contaminated clothing and shoes. Wash clothing before reuse. Clean shoes thoroughly before reuse. Get medical attention if adverse health effects persist or are severe.
	1.42 M 2-Mercaptoethanol	In case of contact, immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse. Clean shoes thoroughly before reuse. Get medical attention if adverse health effects persist or are severe.
Inhalation	: BL21 competent cells	If inhaled, remove to fresh air. If breathing is difficult, give oxygen. If not breathing, give artificial respiration. Get medical attention if adverse health effects persist or are severe.
	pUC18 Control Plasmid DNA	If inhaled, remove to fresh air. If breathing is difficult, give oxygen. If not breathing, give artificial respiration. Get medical attention if adverse health effects persist or are severe.
	1.42 M 2-Mercaptoethanol	If inhaled, remove to fresh air. If breathing is difficult, give oxygen. If not breathing, give artificial respiration. Get medical attention if adverse health effects persist or are severe.
Ingestion	: BL21 competent cells	Do not induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person. Get medical attention if adverse health effects persist or are severe.
	pUC18 Control Plasmid DNA	Do not induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person. Get medical attention if adverse health effects persist or are severe.
	1.42 M 2-Mercaptoethanol	Do not induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person. Get medical attention if adverse health effects persist or are severe.
Protection of first-aiders	: BL21 competent cells	Not applicable.
	pUC18 Control Plasmid DNA	Not applicable.
	1.42 M 2-Mercaptoethanol	Not applicable.
Notes to physician	: No specific treatment. Treat symptomatically. Contact poison treatment specialist immediately if large quantities have been ingested or inhaled.	

5 . Fire-fighting measures

Flammability of the product	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Non-flammable. Non-flammable. Non-flammable.
Products of combustion	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Decomposition products may include the following materials: carbon oxides halogenated compounds metal oxide/oxides No specific data. Decomposition products may include the following materials: carbon oxides sulfur oxides
<u>Extinguishing media</u>		
Suitable	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Use an extinguishing agent suitable for the surrounding fire. Use an extinguishing agent suitable for the surrounding fire. Use an extinguishing agent suitable for the surrounding fire.
Not suitable	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Not applicable. Not applicable. Not applicable.
Special protective equipment for fire-fighters	: Fire-fighters should wear	appropriate protective equipment and self-contained breathing apparatus (SCBA) with a full face-piece operated in positive pressure mode.
Special remarks on fire hazards	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Not available. Not available. Not available.
Special remarks on explosion hazards	: Not available.	

6 . Accidental release measures

Personal precautions	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	No action shall be taken involving any personal risk or without suitable training. Evacuate surrounding areas. Keep unnecessary and unprotected personnel from entering. Do not touch or walk through spilled material. Avoid breathing vapor or mist. Provide adequate ventilation. Wear appropriate respirator when ventilation is inadequate. Put on appropriate personal protective equipment (see section 8). No action shall be taken involving any personal risk or without suitable training. Evacuate surrounding areas. Keep unnecessary and unprotected personnel from entering. Do not touch or walk through spilled material. Avoid breathing vapor or mist. Provide adequate ventilation. Wear appropriate respirator when ventilation is inadequate. Put on appropriate personal protective equipment (see section 8). No action shall be taken involving any personal risk or without suitable training. Evacuate surrounding areas. Keep unnecessary and unprotected personnel from entering. Do not touch or walk through spilled material. Avoid breathing vapor or mist. Provide adequate ventilation. Wear appropriate respirator when ventilation is inadequate. Put on appropriate personal protective equipment (see section 8).
-----------------------------	--	--

6 . Accidental release measures

Environmental precautions	: BL21 competent cells	Avoid dispersal of spilled material and runoff and contact with soil, waterways, drains and sewers. Inform the relevant authorities if the product has caused environmental pollution (sewers, waterways, soil or air).
	pUC18 Control Plasmid DNA	Avoid dispersal of spilled material and runoff and contact with soil, waterways, drains and sewers. Inform the relevant authorities if the product has caused environmental pollution (sewers, waterways, soil or air).
	1.42 M 2-Mercaptoethanol	Avoid dispersal of spilled material and runoff and contact with soil, waterways, drains and sewers. Inform the relevant authorities if the product has caused environmental pollution (sewers, waterways, soil or air).

Methods for cleaning up

Small spill	: BL21 competent cells	Stop leak if without risk. Move containers from spill area. Dilute with water and mop up if water-soluble or absorb with an inert dry material and place in an appropriate waste disposal container. Dispose of via a licensed waste disposal contractor.
	pUC18 Control Plasmid DNA	Stop leak if without risk. Move containers from spill area. Dilute with water and mop up if water-soluble or absorb with an inert dry material and place in an appropriate waste disposal container. Dispose of via a licensed waste disposal contractor.
	1.42 M 2-Mercaptoethanol	Stop leak if without risk. Move containers from spill area. Dilute with water and mop up if water-soluble or absorb with an inert dry material and place in an appropriate waste disposal container. Dispose of via a licensed waste disposal contractor.

7 . Handling and storage

Handling	: BL21 competent cells	Do not ingest. Wash thoroughly after handling.
	pUC18 Control Plasmid DNA	Wash thoroughly after handling.
	1.42 M 2-Mercaptoethanol	Do not ingest. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling.
Storage	:	Store in accordance with local regulations. Store in original container protected from direct sunlight in a dry, cool and well-ventilated area, away from incompatible materials (see section 10) and food and drink. Keep container tightly closed and sealed until ready for use. Containers that have been opened must be carefully resealed and kept upright to prevent leakage. Do not store in unlabeled containers. Use appropriate containment to avoid environmental contamination.

8 . Exposure controls/personal protection

Product name

Exposure limits

United States

BL21 competent cells
Glycerol

ACGIH TLV (United States, 1/2008).
TWA: 10 mg/m³ 8 hour(s). Form: Mist
OSHA PEL (United States, 11/2006).
TWA: 5 mg/m³ 8 hour(s). Form: Respirable fraction
TWA: 15 mg/m³ 8 hour(s). Form: Total dust
OSHA PEL 1989 (United States, 3/1989).
TWA: 5 mg/m³ 8 hour(s). Form: Respirable fraction
TWA: 10 mg/m³ 8 hour(s). Form: Total dust

Manganese dichloride

ACGIH TLV (United States, 1/2008).
TWA: 0.2 mg/m³, (as Mn) 8 hour(s).
OSHA PEL 1989 (United States, 3/1989).
CEIL: 5 mg/m³, (as Mn)
NIOSH REL (United States, 12/2001).

8 . Exposure controls/personal protection

Sucrose	<p>TWA: 1 mg/m³, (as Mn) 10 hour(s). STEL: 3 mg/m³, (as Mn) 15 minute(s). OSHA PEL (United States, 11/2006). CEIL: 5 mg/m³, (as Mn) ACGIH TLV (United States, 1/2008). TWA: 10 mg/m³ 8 hour(s). OSHA PEL 1989 (United States, 3/1989). TWA: 15 mg/m³ 8 hour(s). Form: Total dust TWA: 5 mg/m³ 8 hour(s). Form: Respirable fraction NIOSH REL (United States, 12/2001). TWA: 10 mg/m³ 10 hour(s). Form: Total TWA: 5 mg/m³ 10 hour(s). Form: Respirable fraction OSHA PEL (United States, 11/2006). TWA: 15 mg/m³ 8 hour(s). Form: Total dust TWA: 5 mg/m³ 8 hour(s). Form: Respirable fraction</p>
Dimethyl sulfoxide	<p>AIHA WEEL (United States, 1/2008). TWA: 250 ppm 8 hour(s).</p>
1.42 M 2-Mercaptoethanol 2-Mercaptoethanol	<p>AIHA WEEL (United States, 1/2008). TWA: 0.2 ppm 8 hour(s).</p>

Consult local authorities for acceptable exposure limits.

Engineering measures : If user operations generate dust, fumes, gas, vapor or mist, use process enclosures, local exhaust ventilation or other engineering controls to keep worker exposure to airborne contaminants below any recommended or statutory limits.

Personal protection

- Eyes** : Safety eyewear complying with an approved standard should be used when a risk assessment indicates this is necessary to avoid exposure to liquid splashes, mists, gases or dusts.
- Skin** : Chemical resistant protective gloves and clothing are recommended. The choice of protective gloves or clothing must be based on chemical resistance and other use requirements. Generally, BUNA-N offers acceptable chemical resistance. Individuals who are acutely and specifically sensitive to this chemical may require additional protective clothing.
- Respiratory** : Use a properly fitted, air-purifying or air-fed respirator complying with an approved standard if a risk assessment indicates this is necessary. Respirator selection must be based on known or anticipated exposure levels, the hazards of the product and the safe working limits of the selected respirator.
- Hands** : Chemical-resistant, impervious gloves complying with an approved standard should be worn at all times when handling chemical products if a risk assessment indicates this is necessary.
- Other protection** : Not available.
- Hygiene measures** : Handle as biohazard material (Biosafety level 1). Wash hands, forearms and face thoroughly after handling chemical products, before eating, smoking and using the lavatory and at the end of the working period. Appropriate techniques should be used to remove potentially contaminated clothing. Wash contaminated clothing before reusing. Ensure that eyewash stations and safety showers are close to the workstation location.

9 . Physical and chemical properties

Physical state	<p>: BL21 competent cells Liquid. pUC18 Control Plasmid Liquid. DNA 1.42 M 2-Mercaptoethanol Liquid.</p>
-----------------------	--

9 . Physical and chemical properties

Flash point	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Not applicable. Not applicable. Not applicable.
Color	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Not available. Not available. Not available.
Odor	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Not available. Not available. Not available.
pH	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Neutral. Neutral. Neutral.
Boiling/condensation point	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Lowest known value: 100°C (212°F) (Water). Weighted average: 116.29°C (241.3°F) Lowest known value: 100°C (212°F) (Water). Lowest known value: 100°C (212°F) (Water). Weighted average: 105.7°C (222.3°F)
Melting/freezing point	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	May start to solidify at the following temperature: 19.8°C (67.6°F) This is based on data for the following ingredient: Glycerol. Weighted average: 1.7°C (35.1°F) May start to solidify at the following temperature: 0°C (32°F) This is based on data for the following ingredient: Water. May start to solidify at the following temperature: 0°C (32°F) This is based on data for the following ingredient: Water.
Relative density	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Weighted average: 1.41 (Water = 1) Not available. Only known value: 1.1 (Water = 1) (2-Mercaptoethanol).
Vapor pressure	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Highest known value: 2.3 kPa (17.5 mm Hg) (at 20°C) (Water). Highest known value: 2.3 kPa (17.5 mm Hg) (at 20°C) (Water). Highest known value: 2.3 kPa (17.5 mm Hg) (at 20°C) (Water). Weighted average: 2.08 kPa (15.6 mm Hg) (at 20°C)
Vapor density	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Highest known value: 3.1 (Air = 1) (Glycerol). Weighted average: 0.83 (Air = 1) Highest known value: 0.62 (Air = 1) (Water). Highest known value: 2.7 (Air = 1) (2-Mercaptoethanol). Weighted average: 0.83 (Air = 1)

10 . Stability and reactivity

Stability and reactivity	: The product is stable.	
Incompatibility with various substances	: Reactive or incompatible with the following materials: oxidizing materials and acids.	
Hazardous decomposition products	: BL21 competent cells pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol	Under normal conditions of storage and use, hazardous decomposition products should not be produced. Under normal conditions of storage and use, hazardous decomposition products should not be produced. Under normal conditions of storage and use, hazardous decomposition products should not be produced.

10 . Stability and reactivity

11 . Toxicological information

Acute toxicity

Product/ingredient name	Result	Species	Dose	Exposure
Dimethyl sulfoxide	LD50 Dermal	Rat	40 gm/kg	-
	LD50 Oral	Rat	14500 mg/kg	-
Sucrose	LD50 Oral	Rat	29700 mg/kg	-
Manganese dichloride	LD50 Oral	Rat	250 mg/kg	-
Glycerol	LD50 Dermal	Rabbit	>10 gm/kg	-
	LD50 Oral	Rat	12600 mg/kg	-
Potassium chloride	LD50 Oral	Rat	2600 mg/kg	-

Eyes	: BL21 competent cells	No known significant effects or critical hazards.
	pUC18 Control Plasmid DNA	No known significant effects or critical hazards.
Skin	: BL21 competent cells	No known significant effects or critical hazards.
	pUC18 Control Plasmid DNA	No known significant effects or critical hazards.
Inhalation	: BL21 competent cells	No known significant effects or critical hazards.
	pUC18 Control Plasmid DNA	No known significant effects or critical hazards.
Ingestion	: BL21 competent cells	Toxic if swallowed.
	pUC18 Control Plasmid DNA	No known significant effects or critical hazards.
	: 1.42 M 2-Mercaptoethanol	Irritating to eyes.
	: 1.42 M 2-Mercaptoethanol	Irritating to skin.
	: 1.42 M 2-Mercaptoethanol	No known significant effects or critical hazards.
	: 1.42 M 2-Mercaptoethanol	Toxic if swallowed.

Classification

Product/ingredient name	ACGIH	IARC	EPA	NIOSH	NTP	OSHA
BL21 competent cells						
Sucrose	A4	-	-	-	-	-

Potential chronic health effects

Chronic effects	: Contains material that may cause target organ damage, based on animal data.
Carcinogenicity	: No known significant effects or critical hazards.
Mutagenicity	: No known significant effects or critical hazards.
Teratogenicity	: No known significant effects or critical hazards.
Developmental effects	: No known significant effects or critical hazards.
Fertility effects	: No known significant effects or critical hazards.

Over-exposure signs/symptoms

Inhalation	: No specific data.	
Ingestion	: No specific data.	
Skin	: No specific data.	
Eyes	: No specific data.	
Other adverse effects	: BL21 competent cells	Not available.
	pUC18 Control Plasmid DNA	Not available.
	: 1.42 M 2-Mercaptoethanol	Not available.

12 . Ecological information

Environmental effects : No known significant effects or critical hazards.

Aquatic ecotoxicity

Product/ingredient name	Test	Result	Species	Exposure
Dimethyl sulfoxide	-	Acute LC50 35 to 37 ml/L Fresh water	Fish	96 hours
	-	Acute LC50 34000000 ug/L Fresh water	Fish	96 hours
Manganese dichloride	-	Acute EC50 4700 ug/L Fresh water	Daphnia	48 hours
Glycerol	-	Acute LC50 54 to 57 ml/L Fresh water	Fish	96 hours
Potassium chloride	-	Acute EC50 83000 ug/L Fresh water	Daphnia	48 hours
	-	Acute LC50 337 mg/L Fresh water	Daphnia	48 hours
	-	Acute LC50 435000 ug/L Fresh water	Fish	96 hours

Other adverse effects : No known significant effects or critical hazards.

13 . Disposal considerations

Waste disposal : The generation of waste should be avoided or minimized wherever possible. Dispose of surplus and non-recyclable products via a licensed waste disposal contractor. Disposal of this product, solutions and any by-products should at all times comply with the requirements of environmental protection and waste disposal legislation and any regional local authority requirements. Avoid dispersal of spilled material and runoff and contact with soil, waterways, drains and sewers.

Disposal should be in accordance with applicable regional, national and local laws and regulations. Local regulations may be more stringent than regional or national requirements.

The information presented below only applies to the material as supplied. The identification based on characteristic(s) or listing may not apply if the material has been used or otherwise contaminated. It is the responsibility of the waste generator to determine the toxicity and physical properties of the material generated to determine the proper waste identification and disposal methods in compliance with applicable regulations.

Refer to Section 7: HANDLING AND STORAGE and Section 8: EXPOSURE CONTROLS/PERSONAL PROTECTION for additional handling information and protection of employees.

14 . Transport information

Regulatory information

DOT /IMDG / IATA : Not regulated.

15 . Regulatory information

HCS Classification	: BL21 competent cells	Toxic material
		Target organ effects
	pUC18 Control Plasmid DNA	Not regulated.
	1.42 M 2-Mercaptoethanol	Toxic material
		Irritating material

15 . Regulatory information

<p>U.S. Federal regulations</p>	<p>: BL21 competent cells</p> <p>pUC18 Control Plasmid DNA</p> <p>1.42 M 2-Mercaptoethanol</p> <p>BL21 competent cells</p> <p>pUC18 Control Plasmid DNA</p> <p>1.42 M 2-Mercaptoethanol</p> <p>BL21 competent cells pUC18 Control Plasmid DNA</p> <p>1.42 M 2-Mercaptoethanol</p> <p>BL21 competent cells</p> <p>pUC18 Control Plasmid DNA</p> <p>1.42 M 2-Mercaptoethanol</p> <p>BL21 competent cells</p> <p>pUC18 Control Plasmid DNA</p> <p>1.42 M 2-Mercaptoethanol</p> <p>BL21 competent cells</p> <p>pUC18 Control Plasmid DNA</p> <p>1.42 M 2-Mercaptoethanol</p>	<p>United States inventory (TSCA 8b): All components are listed or exempted.</p> <p>United States inventory (TSCA 8b): All components are listed or exempted.</p> <p>United States inventory (TSCA 8b): All components are listed or exempted.</p> <p>SARA 302/304/311/312 extremely hazardous substances: No products were found.</p> <p>SARA 302/304 emergency planning and notification: No products were found.</p> <p>SARA 302/304/311/312 hazardous chemicals: Potassium chloride; Glycerol; Manganese dichloride; Sucrose</p> <p>SARA 311/312 MSDS distribution - chemical inventory - hazard identification: Potassium chloride: Immediate (acute) health hazard, Delayed (chronic) health hazard; Glycerol: Immediate (acute) health hazard, Delayed (chronic) health hazard; Manganese dichloride: Delayed (chronic) health hazard; Sucrose: Delayed (chronic) health hazard</p> <p>SARA 302/304/311/312 extremely hazardous substances: No products were found.</p> <p>SARA 302/304 emergency planning and notification: No products were found.</p> <p>SARA 302/304/311/312 hazardous chemicals: No products were found.</p> <p>SARA 311/312 MSDS distribution - chemical inventory - hazard identification: No products were found.</p> <p>SARA 302/304/311/312 extremely hazardous substances: No products were found.</p> <p>SARA 302/304 emergency planning and notification: No products were found.</p> <p>SARA 302/304/311/312 hazardous chemicals: 2-Mercaptoethanol</p> <p>SARA 311/312 MSDS distribution - chemical inventory - hazard identification: 2-Mercaptoethanol: Fire hazard, Immediate (acute) health hazard, Delayed (chronic) health hazard</p> <p>Clean Water Act (CWA) 307: No products were found.</p> <p>Clean Water Act (CWA) 307: No products were found.</p> <p>Clean Water Act (CWA) 307: No products were found.</p> <p>Clean Water Act (CWA) 311: No products were found.</p> <p>Clean Water Act (CWA) 311: Edetic acid</p> <p>Clean Water Act (CWA) 311: No products were found.</p> <p>Clean Air Act (CAA) 112 accidental release prevention: No products were found.</p> <p>Clean Air Act (CAA) 112 accidental release prevention: No products were found.</p> <p>Clean Air Act (CAA) 112 accidental release prevention: No products were found.</p> <p>Clean Air Act (CAA) 112 regulated flammable substances : No products were found.</p> <p>Clean Air Act (CAA) 112 regulated flammable substances : No products were found.</p> <p>Clean Air Act (CAA) 112 regulated flammable substances : No products were found.</p>
--	--	--

15 . Regulatory information

BL21 competent cells	Clean Air Act (CAA) 112 regulated toxic substances: No products were found.
pUC18 Control Plasmid DNA	Clean Air Act (CAA) 112 regulated toxic substances: No products were found.
1.42 M 2-Mercaptoethanol	Clean Air Act (CAA) 112 regulated toxic substances: No products were found.

SARA 313

	<u>Product name</u>	<u>CAS number</u>	<u>Concentration</u>
Form R - Reporting requirements	: BL21 competent cells		
	Manganese dichloride	7773-01-5	5 - 10
	Hexaamminecobalt trichloride	10534-89-1	0.1 - 1
Supplier notification	: BL21 competent cells		
	Manganese dichloride	7773-01-5	5 - 10
	Hexaamminecobalt trichloride	10534-89-1	0.1 - 1

SARA 313 notifications must not be detached from the MSDS and any copying and redistribution of the MSDS shall include copying and redistribution of the notice attached to copies of the MSDS subsequently redistributed.

State regulations	: BL21 competent cells	<p>Connecticut Carcinogen Reporting: None of the components are listed.</p> <p>Connecticut Hazardous Material Survey: None of the components are listed.</p> <p>Florida substances: None of the components are listed.</p> <p>Illinois Chemical Safety Act: None of the components are listed.</p> <p>Illinois Toxic Substances Disclosure to Employee Act: None of the components are listed.</p> <p>Louisiana Reporting: None of the components are listed.</p> <p>Louisiana Spill: None of the components are listed.</p> <p>Massachusetts Spill: None of the components are listed.</p> <p>Massachusetts Substances: The following components are listed: Glycerol; Sucrose</p> <p>Michigan Critical Material: None of the components are listed.</p> <p>Minnesota Hazardous Substances: None of the components are listed.</p> <p>New Jersey Hazardous Substances: The following components are listed: Manganese dichloride</p> <p>New Jersey Spill: None of the components are listed.</p> <p>New Jersey Toxic Catastrophe Prevention Act: None of the components are listed.</p> <p>New York Acutely Hazardous Substances: None of the components are listed.</p> <p>New York Toxic Chemical Release Reporting: None of the components are listed.</p> <p>Pennsylvania RTK Hazardous Substances: The following components are listed: Glycerol; Manganese dichloride; Sucrose</p> <p>Rhode Island Hazardous Substances: None of the components are listed.</p>
	pUC18 Control Plasmid DNA	<p>Connecticut Carcinogen Reporting: None of the components are listed.</p> <p>Connecticut Hazardous Material Survey: None of the components are listed.</p> <p>Florida substances: None of the components are listed.</p> <p>Illinois Chemical Safety Act: None of the components are listed.</p> <p>Illinois Toxic Substances Disclosure to Employee Act: None of the components are listed.</p> <p>Louisiana Reporting: None of the components are listed.</p> <p>Louisiana Spill: None of the components are listed.</p> <p>Massachusetts Spill: None of the components are listed.</p>

15 . Regulatory information

Massachusetts Substances: None of the components are listed.

Michigan Critical Material: None of the components are listed.

Minnesota Hazardous Substances: None of the components are listed.

New Jersey Hazardous Substances: None of the components are listed.

New Jersey Spill: None of the components are listed.

New Jersey Toxic Catastrophe Prevention Act: None of the components are listed.

New York Acutely Hazardous Substances: None of the components are listed.

New York Toxic Chemical Release Reporting: None of the components are listed.

Pennsylvania RTK Hazardous Substances: None of the components are listed.

Rhode Island Hazardous Substances: None of the components are listed.

1.42 M 2-Mercaptoethanol

Connecticut Carcinogen Reporting: None of the components are listed.

Connecticut Hazardous Material Survey: None of the components are listed.

Florida substances: None of the components are listed.

Illinois Chemical Safety Act: None of the components are listed.

Illinois Toxic Substances Disclosure to Employee Act: None of the components are listed.

Louisiana Reporting: None of the components are listed.

Louisiana Spill: None of the components are listed.

Massachusetts Spill: None of the components are listed.

Massachusetts Substances: The following components are listed: 2-Mercaptoethanol

Michigan Critical Material: None of the components are listed.

Minnesota Hazardous Substances: None of the components are listed.

New Jersey Hazardous Substances: None of the components are listed.

New Jersey Spill: None of the components are listed.

New Jersey Toxic Catastrophe Prevention Act: None of the components are listed.

New York Acutely Hazardous Substances: None of the components are listed.

New York Toxic Chemical Release Reporting: None of the components are listed.

Pennsylvania RTK Hazardous Substances: The following components are listed: 2-Mercaptoethanol

Rhode Island Hazardous Substances: None of the components are listed.

State regulations -
California Prop. 65

: No products were found.

16 . Other information

Label requirements	: BL21 competent cells	HARMFUL IF SWALLOWED. CONTAINS MATERIAL THAT MAY CAUSE TARGET ORGAN DAMAGE, BASED ON ANIMAL DATA.
	pUC18 Control Plasmid DNA	NOT EXPECTED TO PRODUCE SIGNIFICANT ADVERSE HEALTH EFFECTS WHEN THE RECOMMENDED INSTRUCTIONS FOR USE ARE FOLLOWED.
	1.42 M 2-Mercaptoethanol	HARMFUL IF SWALLOWED. CAUSES EYE AND SKIN IRRITATION.

Date of issue : 11/21/2008

Version : 1

Notice to reader

DISCLAIMER: This Material Safety Data Sheet is offered without charge to the clients of Agilent Technologies. Data is the most current available to Agilent Technologies at the time of preparation and is issued as a matter of information only, no warranty as to its accuracy or completeness is expressed or implied.

 Indicates information that has changed from previously issued version.

1. IDENTIFICATION OF THE SUBSTANCE/PREPARATION AND THE COMPANY/UNDERTAKING

Product code 18258
Product name ME DH5A COMPETENT CELLS

Contact manufacturer
 INVITROGEN CORPORATON
 1600 FARADAY AVENUE
 PO BOX 6482
 CARLSBAD, CA 92008
 760-603-7200

INVITROGEN CORPORATION
 2270 INDUSTRIAL STREET
 BURLINGTON, ONT
 CANADA L7P 1A1
 800-263-6236

GIBCO PRODUCTS
 INVITROGEN CORPORATION
 3175 STALEY ROAD P.O. BOX 68
 GRAND ISLAND, NY 14072
 716-774-6700

2. COMPOSITION/INFORMATION ON INGREDIENTS

Hazardous/Non-hazardous Components

Chemical Name	CAS-No	Weight %
dimethylsulfoxide	67-68-5	3-7

3. HAZARDS IDENTIFICATION

Emergency Overview

Irritating to eyes. Irritating to skin. Components of the product may be absorbed into the body through the skin.

Form
 Liquid

Principle Routes of Exposure/

Potential Health effects

Eyes	Irritating to eyes.
Skin	Irritating to skin. Components of the product may be absorbed into the body through the skin.
Inhalation	May cause irritation of respiratory tract.
Ingestion	May be harmful if swallowed.

Specific effects

Carcinogenic effects	No information available
Mutagenic effects	No information available
Reproductive toxicity	No information available
Sensitization	No information available

Target Organ Effects

Eyes. Skin.

4. FIRST AID MEASURES

Skin contact	Wash off immediately with plenty of water
Eye contact	Rinse thoroughly with plenty of water, also under the eyelids.
Ingestion	Never give anything by mouth to an unconscious person
Inhalation	Move to fresh air
Notes to physician	Treat symptomatically

5. FIRE-FIGHTING MEASURES

Suitable extinguishing media	Dry chemical
Special protective equipment for firefighters	Wear self-contained breathing apparatus and protective suit

6. ACCIDENTAL RELEASE MEASURES

Personal precautions	Use personal protective equipment
Methods for cleaning up	Soak up with inert absorbent material

7. HANDLING AND STORAGE

Handling	Avoid contact with skin and eyes.
Storage	Keep in properly labelled containers

8. EXPOSURE CONTROLS / PERSONAL PROTECTION**Occupational exposure controls****Exposure limits**

Chemical Name	OSHA PEL (TWA)	OSHA PEL (Ceiling)	ACGIH OEL (TWA)	ACGIH OEL (STEL)
dimethylsulfoxide	-	-	-	-

Engineering measures	Ensure adequate ventilation, especially in confined areas
----------------------	---

Personal protective equipment

Respiratory protection	In case of insufficient ventilation wear suitable respiratory equipment
Hand protection	Protective gloves
Eye protection	Safety glasses with side-shields
Skin and body protection	Lightweight protective clothing
Hygiene measures	Handle in accordance with good industrial hygiene and safety practice
Environmental exposure controls	Prevent product from entering drains

9. PHYSICAL AND CHEMICAL PROPERTIES**General Information**

Form	Liquid
------	--------

Important Health Safety and Environmental Information

Boiling point/range	°C 189	°F No data available
Melting point/range	°C 18.4	°F No data available
Flash point	°C 94	°F No data available
Autoignition temperature	°C No data available	°F No data available
Oxidizing properties	No information available	
Water solubility	soluble	

10. STABILITY AND REACTIVITY

Stability	Stable under normal conditions.
Materials to avoid	No information available
Hazardous decomposition products	No information available
Polymerization	Hazardous polymerisation does not occur

11. TOXICOLOGICAL INFORMATION

Acute toxicity

Chemical Name	LD50 (oral, rat/mouse)	LD50 (dermal, rat/rabbit)	LC50 (inhalation, rat/mouse)
dimethylsulfoxide	14500 mg/kg (Rat)	No data available	No data available

Principle Routes of Exposure/

Potential Health effects

Eyes	Irritating to eyes.
Skin	Irritating to skin. Components of the product may be absorbed into the body through the skin.
Inhalation	May cause irritation of respiratory tract.
Ingestion	May be harmful if swallowed.

Specific effects

Carcinogenic effects	No information available
Mutagenic effects	No information available
Reproductive toxicity	No information available
Sensitization	No information available

Target Organ Effects	Eyes. Skin.
----------------------	-------------

12. ECOLOGICAL INFORMATION

Ecotoxicity effects	No information available.
Mobility	No information available.
Biodegradation	Inherently biodegradable.
Bioaccumulation	Does not bioaccumulate.

13. DISPOSAL CONSIDERATIONS

Dispose of in accordance with local regulations

14. TRANSPORT INFORMATION

IATA

Proper shipping name Not classified as dangerous in the meaning of transport regulations
Hazard Class No information available
Subsidiary Class No information available
Packing group No information available
UN-No No information available

Proper shipping name Not classified as dangerous within the meaning of transport regulations

15. REGULATORY INFORMATION

International Inventories

Chemical Name	TSCA	PICCS	ENCS	DSL	NDSL	AICS
dimethylsulfoxide	Listed	Listed	Listed	Listed	-	Listed

U.S. Federal Regulations

SARA 313
Not regulated

Clean Air Act, Section 112 Hazardous Air Pollutants (HAPs) (see 40 CFR 61)

This product contains the following HAPs:

U.S. State Regulations

Chemical Name	Massachusetts - RTK	New Jersey - RTK	Pennsylvania - RTK	Illinois - RTK	Rhode Island - RTK
dimethylsulfoxide	-	-	-	-	-

California Proposition 65

This product contains the following Proposition 65 chemicals:

WHMIS hazard class:

D2B Toxic materials

This product has been classified according to the hazard criteria of the CPR and the MSDS contains all of the information required by the CPR

16. OTHER INFORMATION

This material is sold for research and development purposes only. It is not for any human or animal therapeutic or clinical diagnostic use. It is not intended for food, drug, household, agricultural, or cosmetic use. An individual technically qualified to handle potentially hazardous chemicals must supervise the use of this material.

The above information was acquired by diligent search and/or investigation and the recommendations are based on prudent application of professional judgment. The information shall not be taken as being all inclusive and is to be used only as a guide. All materials and mixtures may be present unknown hazards and should be used with caution. Since Invitrogen Corporation cannot control the actual methods, volumes, or conditions of use, the Company shall not be held liable for any damages or losses resulting from the handling or from contact with the product as described herein. THE INFORMATION IN THIS MSDS DOES NOT CONSTITUTE A WARRANTY, EXPRESS OR IMPLIED, INCLUDING ANY IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE.

End of Safety Data Sheet

Material Safety Data Sheet



Stratagene XL1-Blue Competent Cells, Catalog #200249

1. Product and company identification

Product name : Stratagene XL1-Blue Competent Cells, Catalog #200249
Part No. : pUC18 Control Plasmid 200231-42
 DNA
 1.42 M 2-Mercaptoethanol 210200-43
 XL1-Blue Competent Cells 200236-41
Manufacturer / Supplier : Agilent Technologies, Inc.
 1834 State Highway 71 West
 Cedar Creek, TX 78612

Emergency telephone number : 1-800-894-1304
Use of the substance/preparation : Chemical Kit
Validation date : 01/09/2009

2. Hazards identification

Physical state : pUC18 Control Plasmid DNA Liquid.
 1.42 M 2-Mercaptoethanol Liquid.
 XL1-Blue Competent Cells Liquid.

Odor : pUC18 Control Plasmid DNA Not available.
 1.42 M 2-Mercaptoethanol Not available.
 XL1-Blue Competent Cells Not available.

OSHA/HCS status : pUC18 Control Plasmid DNA While this material is not considered hazardous by the OSHA Hazard Communication Standard (29 CFR 1910.1200), this MSDS contains valuable information critical to the safe handling and proper use of the product. This MSDS should be retained and available for employees and other users of this product.
 1.42 M 2-Mercaptoethanol This material is considered hazardous by the OSHA Hazard Communication Standard (29 CFR 1910.1200).
 XL1-Blue Competent Cells This material is considered hazardous by the OSHA Hazard Communication Standard (29 CFR 1910.1200).

Emergency overview-Signal Word : WARNING !

Emergency overview-Label Statement : pUC18 Control Plasmid DNA NOT EXPECTED TO PRODUCE SIGNIFICANT ADVERSE HEALTH EFFECTS WHEN THE RECOMMENDED INSTRUCTIONS FOR USE ARE FOLLOWED.
 1.42 M 2-Mercaptoethanol HARMFUL IF SWALLOWED. CAUSES EYE AND SKIN IRRITATION. MAY CAUSE ALLERGIC SKIN REACTION.
 XL1-Blue Competent Cells HARMFUL IF SWALLOWED. CONTAINS MATERIAL THAT MAY CAUSE TARGET ORGAN DAMAGE, BASED ON ANIMAL DATA.

2. Hazards identification

	pUC18 Control Plasmid DNA	No known significant effects or critical hazards. Avoid prolonged contact with eyes, skin and clothing.
	1.42 M 2-Mercaptoethanol	Toxic if swallowed. Irritating to eyes and skin. May cause sensitization by skin contact. Do not breathe vapor or mist. Do not ingest. Do not get on skin or clothing. Avoid contact with eyes. Wash thoroughly after handling.
	XL1-Blue Competent Cells	Toxic if swallowed. Avoid exposure - obtain special instructions before use. Do not breathe vapor or mist. Do not ingest. Avoid contact with eyes, skin and clothing. Contains material that may cause target organ damage, based on animal data. Wash thoroughly after handling.
	pUC18 Control Plasmid DNA	Not available.
	1.42 M 2-Mercaptoethanol	Not available.
	XL1-Blue Competent Cells	Contains material which may cause damage to the following organs: blood, kidneys, gastrointestinal tract, upper respiratory tract, skin, central nervous system (CNS), eye, lens or cornea.
Routes of entry	: pUC18 Control Plasmid DNA	Eye contact. Ingestion.
	1.42 M 2-Mercaptoethanol	Dermal contact. Eye contact. Inhalation. Ingestion.
	XL1-Blue Competent Cells	Eye contact. Inhalation. Ingestion.
<u>Potential acute health effects</u>		
Eyes	: pUC18 Control Plasmid DNA	No known significant effects or critical hazards.
	1.42 M 2-Mercaptoethanol	Irritating to eyes.
	XL1-Blue Competent Cells	No known significant effects or critical hazards.
Skin	: pUC18 Control Plasmid DNA	No known significant effects or critical hazards.
	1.42 M 2-Mercaptoethanol	Irritating to skin. May cause sensitization by skin contact.
	XL1-Blue Competent Cells	No known significant effects or critical hazards.
Inhalation	: pUC18 Control Plasmid DNA	No known significant effects or critical hazards.
	1.42 M 2-Mercaptoethanol	No known significant effects or critical hazards.
	XL1-Blue Competent Cells	No known significant effects or critical hazards.
Ingestion	: pUC18 Control Plasmid DNA	No known significant effects or critical hazards.
	1.42 M 2-Mercaptoethanol	Toxic if swallowed.
	XL1-Blue Competent Cells	Toxic if swallowed.
Medical conditions aggravated by over-exposure	: pUC18 Control Plasmid DNA	Not applicable.
	1.42 M 2-Mercaptoethanol	Repeated skin exposure can produce local skin destruction or dermatitis. Repeated or prolonged contact with spray or mist may produce chronic eye irritation and severe skin irritation.
	XL1-Blue Competent Cells	Repeated or prolonged exposure to the substance can produce target organs damage.
Over-exposure signs/symptoms	: pUC18 Control Plasmid DNA	Not applicable.
	1.42 M 2-Mercaptoethanol	Not applicable.
	XL1-Blue Competent Cells	Not applicable.

See toxicological information (section 11)

3. Composition/information on ingredients

<u>Name</u>	<u>CAS number</u>	<u>%</u>
1.42 M 2-Mercaptoethanol		
2-Mercaptoethanol	60-24-2	10
XL1-Blue Competent Cells		
Glycerol	56-81-5	5 - 10
Manganese dichloride	7773-01-5	5 - 10
Sucrose	57-50-1	5 - 10
Dimethyl sulfoxide	67-68-5	5 - 10
Potassium chloride	7447-40-7	1 - 5

There are no ingredients or additional ingredients present which, within the current knowledge of the supplier and in the concentrations applicable, are classified as hazardous to health or the environment and hence require reporting in this section.

4. First aid measures

Eye contact	: pUC18 Control Plasmid DNA	In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if adverse health effects persist or are severe.
	1.42 M 2-Mercaptoethanol	In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if adverse health effects persist or are severe.
	XL1-Blue Competent Cells	In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if adverse health effects persist or are severe.
Skin contact	: pUC18 Control Plasmid DNA	In case of contact, immediately flush skin with plenty of water. Remove contaminated clothing and shoes. Wash clothing before reuse. Clean shoes thoroughly before reuse. Get medical attention if adverse health effects persist or are severe.
	1.42 M 2-Mercaptoethanol	In case of contact, immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse. Clean shoes thoroughly before reuse. Get medical attention if adverse health effects persist or are severe.
	XL1-Blue Competent Cells	In case of contact, immediately flush skin with plenty of water. Remove contaminated clothing and shoes. Wash clothing before reuse. Clean shoes thoroughly before reuse. Get medical attention if adverse health effects persist or are severe.
Inhalation	: pUC18 Control Plasmid DNA	If inhaled, remove to fresh air. If breathing is difficult, give oxygen. If not breathing, give artificial respiration. Get medical attention if adverse health effects persist or are severe.
	1.42 M 2-Mercaptoethanol	If inhaled, remove to fresh air. If breathing is difficult, give oxygen. If not breathing, give artificial respiration. Get medical attention if adverse health effects persist or are severe.
	XL1-Blue Competent Cells	If inhaled, remove to fresh air. If breathing is difficult, give oxygen. If not breathing, give artificial respiration. Get medical attention if adverse health effects persist or are severe.

4. First aid measures

Ingestion	: pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol XL1-Blue Competent Cells	Do not induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person. Get medical attention if adverse health effects persist or are severe. Do not induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person. Get medical attention if adverse health effects persist or are severe. Do not induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person. Get medical attention if adverse health effects persist or are severe.
Protection of first-aiders	: pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol XL1-Blue Competent Cells	Not applicable. Not applicable. Not applicable.
Notes to physician	: No specific treatment. Treat symptomatically. Contact poison treatment specialist immediately if large quantities have been ingested or inhaled.	

5. Fire-fighting measures

Flammability of the product	: pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol XL1-Blue Competent Cells	Non-flammable. Non-flammable. Non-flammable.
Products of combustion	: pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol XL1-Blue Competent Cells	No specific data. Decomposition products may include the following materials: carbon oxides sulfur oxides Decomposition products may include the following materials: carbon oxides sulfur oxides halogenated compounds metal oxide/oxides
Extinguishing media		
Suitable	: pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol XL1-Blue Competent Cells	Use an extinguishing agent suitable for the surrounding fire. Use an extinguishing agent suitable for the surrounding fire. Use an extinguishing agent suitable for the surrounding fire.
Not suitable	: pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol XL1-Blue Competent Cells	Not applicable. Not applicable. Not applicable.
Special protective equipment for fire-fighters	: Fire-fighters should wear appropriate protective equipment and self-contained breathing apparatus (SCBA) with a full face-piece operated in positive pressure mode.	
Special remarks on fire hazards	: pUC18 Control Plasmid DNA 1.42 M 2-Mercaptoethanol XL1-Blue Competent Cells	Not available. Not available. Not available.
Special remarks on explosion hazards	: Not available.	

6 . Accidental release measures

Personal precautions	: pUC18 Control Plasmid DNA	No action shall be taken involving any personal risk or without suitable training. Evacuate surrounding areas. Keep unnecessary and unprotected personnel from entering. Do not touch or walk through spilled material. Avoid breathing vapor or mist. Provide adequate ventilation. Wear appropriate respirator when ventilation is inadequate. Put on appropriate personal protective equipment (see section 8).
	1.42 M 2-Mercaptoethanol	No action shall be taken involving any personal risk or without suitable training. Evacuate surrounding areas. Keep unnecessary and unprotected personnel from entering. Do not touch or walk through spilled material. Avoid breathing vapor or mist. Provide adequate ventilation. Wear appropriate respirator when ventilation is inadequate. Put on appropriate personal protective equipment (see section 8).
	XL1-Blue Competent Cells	No action shall be taken involving any personal risk or without suitable training. Evacuate surrounding areas. Keep unnecessary and unprotected personnel from entering. Do not touch or walk through spilled material. Avoid breathing vapor or mist. Provide adequate ventilation. Wear appropriate respirator when ventilation is inadequate. Put on appropriate personal protective equipment (see section 8).
Environmental precautions	: pUC18 Control Plasmid DNA	Avoid dispersal of spilled material and runoff and contact with soil, waterways, drains and sewers. Inform the relevant authorities if the product has caused environmental pollution (sewers, waterways, soil or air).
	1.42 M 2-Mercaptoethanol	Avoid dispersal of spilled material and runoff and contact with soil, waterways, drains and sewers. Inform the relevant authorities if the product has caused environmental pollution (sewers, waterways, soil or air).
	XL1-Blue Competent Cells	Avoid dispersal of spilled material and runoff and contact with soil, waterways, drains and sewers. Inform the relevant authorities if the product has caused environmental pollution (sewers, waterways, soil or air).
Methods for cleaning up Small spill	: pUC18 Control Plasmid DNA	Stop leak if without risk. Move containers from spill area. Dilute with water and mop up if water-soluble or absorb with an inert dry material and place in an appropriate waste disposal container. Dispose of via a licensed waste disposal contractor.
	1.42 M 2-Mercaptoethanol	Stop leak if without risk. Move containers from spill area. Dilute with water and mop up if water-soluble or absorb with an inert dry material and place in an appropriate waste disposal container. Dispose of via a licensed waste disposal contractor.
	XL1-Blue Competent Cells	Stop leak if without risk. Move containers from spill area. Dilute with water and mop up if water-soluble or absorb with an inert dry material and place in an appropriate waste disposal container. Dispose of via a licensed waste disposal contractor.

7 . Handling and storage

Handling	: pUC18 Control Plasmid DNA	Wash thoroughly after handling.
	1.42 M 2-Mercaptoethanol	Do not ingest. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling.
	XL1-Blue Competent Cells	Do not ingest. Wash thoroughly after handling.

7. Handling and storage

Storage : Store in accordance with local regulations. Store in original container protected from direct sunlight in a dry, cool and well-ventilated area, away from incompatible materials (see section 10) and food and drink. Keep container tightly closed and sealed until ready for use. Containers that have been opened must be carefully resealed and kept upright to prevent leakage. Do not store in unlabeled containers. Use appropriate containment to avoid environmental contamination.

8. Exposure controls/personal protection

Product name

Exposure limits

United States

1.42 M 2-Mercaptoethanol

2-Mercaptoethanol

AIHA WEEL (United States, 1/2008).

TWA: 0.2 ppm 8 hour(s).

XL1-Blue Competent Cells

Glycerol

ACGIH TLV (United States, 1/2008).

TWA: 10 mg/m³ 8 hour(s). Form: Mist

OSHA PEL (United States, 11/2006).

TWA: 5 mg/m³ 8 hour(s). Form: Respirable fraction

TWA: 15 mg/m³ 8 hour(s). Form: Total dust

OSHA PEL 1989 (United States, 3/1989).

TWA: 5 mg/m³ 8 hour(s). Form: Respirable fraction

TWA: 10 mg/m³ 8 hour(s). Form: Total dust

Manganese dichloride

ACGIH TLV (United States, 1/2008).

TWA: 0.2 mg/m³, (as Mn) 8 hour(s).

OSHA PEL 1989 (United States, 3/1989).

CEIL: 5 mg/m³, (as Mn)

NIOSH REL (United States, 12/2001).

TWA: 1 mg/m³, (as Mn) 10 hour(s).

STEL: 3 mg/m³, (as Mn) 15 minute(s).

OSHA PEL (United States, 11/2006).

CEIL: 5 mg/m³, (as Mn)

Sucrose

ACGIH TLV (United States, 1/2008).

TWA: 10 mg/m³ 8 hour(s).

OSHA PEL 1989 (United States, 3/1989).

TWA: 15 mg/m³ 8 hour(s). Form: Total dust

TWA: 5 mg/m³ 8 hour(s). Form: Respirable fraction

NIOSH REL (United States, 12/2001).

TWA: 10 mg/m³ 10 hour(s). Form: Total

TWA: 5 mg/m³ 10 hour(s). Form: Respirable fraction

OSHA PEL (United States, 11/2006).

TWA: 15 mg/m³ 8 hour(s). Form: Total dust

TWA: 5 mg/m³ 8 hour(s). Form: Respirable fraction

Dimethyl sulfoxide

AIHA WEEL (United States, 1/2008).

TWA: 250 ppm 8 hour(s).

Consult local authorities for acceptable exposure limits.

Engineering measures

: If user operations generate dust, fumes, gas, vapor or mist, use process enclosures, local exhaust ventilation or other engineering controls to keep worker exposure to airborne contaminants below any recommended or statutory limits.

Personal protection

Eyes

: Safety eyewear complying with an approved standard should be used when a risk assessment indicates this is necessary to avoid exposure to liquid splashes, mists, gases or dusts.

8 . Exposure controls/personal protection

Skin	: Chemical resistant protective gloves and clothing are recommended. The choice of protective gloves or clothing must be based on chemical resistance and other use requirements. Generally, BUNA-N offers acceptable chemical resistance. Individuals who are acutely and specifically sensitive to this chemical may require additional protective clothing.
Respiratory	: Use a properly fitted, air-purifying or air-fed respirator complying with an approved standard if a risk assessment indicates this is necessary. Respirator selection must be based on known or anticipated exposure levels, the hazards of the product and the safe working limits of the selected respirator.
Hands	: Chemical-resistant, impervious gloves complying with an approved standard should be worn at all times when handling chemical products if a risk assessment indicates this is necessary.
Other protection	: Not available.
Hygiene measures	: Wash hands, forearms and face thoroughly after handling chemical products, before eating, smoking and using the lavatory and at the end of the working period. Appropriate techniques should be used to remove potentially contaminated clothing. Wash contaminated clothing before reusing. Ensure that eyewash stations and safety showers are close to the workstation location.

9 . Physical and chemical properties

Physical state	: pUC18 Control Plasmid Liquid. DNA
	: 1.42 M 2-Mercaptoethanol Liquid.
	: XL1-Blue Competent Liquid. Cells
Flash point	: pUC18 Control Plasmid Not applicable. DNA
	: 1.42 M 2-Mercaptoethanol Not applicable.
	: XL1-Blue Competent Not applicable. Cells
Color	: pUC18 Control Plasmid Not available. DNA
	: 1.42 M 2-Mercaptoethanol Not available.
	: XL1-Blue Competent Not available. Cells
Odor	: pUC18 Control Plasmid Not available. DNA
	: 1.42 M 2-Mercaptoethanol Not available.
	: XL1-Blue Competent Not available. Cells
pH	: pUC18 Control Plasmid Neutral. DNA
	: 1.42 M 2-Mercaptoethanol Neutral.
	: XL1-Blue Competent Neutral. Cells
Boiling/condensation point	: pUC18 Control Plasmid Lowest known value: 100°C (212°F) (Water). DNA
	: 1.42 M 2-Mercaptoethanol Lowest known value: 100°C (212°F) (Water). Weighted average: 105.7°C (222.3°F)
	: XL1-Blue Competent Lowest known value: 100°C (212°F) (Water). Weighted average: 122.01°C (251.6°F) Cells
Melting/freezing point	: pUC18 Control Plasmid May start to solidify at the following temperature: 0°C (32°F) DNA This is based on data for the following ingredient: Water.
	: 1.42 M 2-Mercaptoethanol May start to solidify at the following temperature: 0°C (32°F) This is based on data for the following ingredient: Water.
	: XL1-Blue Competent May start to solidify at the following temperature: 19.8°C (67.6°F) This is based on data for the following ingredient: Cells Glycerol. Weighted average: 3.02°C (37.4°F)

9 . Physical and chemical properties

Relative density	: pUC18 Control Plasmid DNA	Not available.
	: 1.42 M 2-Mercaptoethanol	Only known value: 1.1 (Water = 1) (2-Mercaptoethanol).
	: XL1-Blue Competent Cells	Weighted average: 1.29 (Water = 1)
Vapor pressure	: pUC18 Control Plasmid DNA	Highest known value: 2.3 kPa (17.5 mm Hg) (at 20°C) (Water).
	: 1.42 M 2-Mercaptoethanol	Highest known value: 2.3 kPa (17.5 mm Hg) (at 20°C) (Water). Weighted average: 2.08 kPa (15.6 mm Hg) (at 20°C)
	: XL1-Blue Competent Cells	Highest known value: 2.3 kPa (17.5 mm Hg) (at 20°C) (Water). Weighted average: 2.11 kPa (15.83 mm Hg) (at 20°C)
Vapor density	: pUC18 Control Plasmid DNA	Highest known value: 0.62 (Air = 1) (Water).
	: 1.42 M 2-Mercaptoethanol	Highest known value: 2.7 (Air = 1) (2-Mercaptoethanol). Weighted average: 0.83 (Air = 1)
	: XL1-Blue Competent Cells	Highest known value: 3.1 (Air = 1) (Glycerol). Weighted average: 0.98 (Air = 1)
Evaporation rate	: pUC18 Control Plasmid DNA	Not available.
	: 1.42 M 2-Mercaptoethanol	Not available.
	: XL1-Blue Competent Cells	0.026 (Dimethyl sulfoxide) compared with Butyl acetate.

10 . Stability and reactivity

Stability and reactivity	: The product is stable.
Incompatibility with various substances	: Highly reactive or incompatible with the following materials: oxidizing materials and organic materials. Reactive or incompatible with the following materials: acids.
Hazardous decomposition products	: pUC18 Control Plasmid DNA Under normal conditions of storage and use, hazardous decomposition products should not be produced. 1.42 M 2-Mercaptoethanol Under normal conditions of storage and use, hazardous decomposition products should not be produced. XL1-Blue Competent Cells Under normal conditions of storage and use, hazardous decomposition products should not be produced.
Conditions of reactivity - Flammability	: Flammable in the presence of the following materials or conditions: open flames, sparks and static discharge.

11 . Toxicological information

Acute toxicity

Product/ingredient name	Result	Species	Dose	Exposure
Dimethyl sulfoxide	LD50 Dermal	Rat	40 gm/kg	-
	LD50 Oral	Rat	14500 mg/kg	-
Sucrose	LD50 Oral	Rat	29700 mg/kg	-
Manganese dichloride	LD50 Oral	Rat	250 mg/kg	-
Glycerol	LD50 Dermal	Rabbit	>10 gm/kg	-
	LD50 Oral	Rat	12600 mg/kg	-
Potassium chloride	LD50 Oral	Rat	2600 mg/kg	-

Eyes	: pUC18 Control Plasmid DNA	No known significant effects or critical hazards.
	: 1.42 M 2-Mercaptoethanol	Irritating to eyes.
	: XL1-Blue Competent Cells	No known significant effects or critical hazards.

11 . Toxicological information

Skin	: pUC18 Control Plasmid	No known significant effects or critical hazards.
	DNA	
	1.42 M 2-Mercaptoethanol	Irritating to skin. May cause sensitization by skin contact.
	XL1-Blue Competent Cells	No known significant effects or critical hazards.
Inhalation	: pUC18 Control Plasmid	No known significant effects or critical hazards.
	DNA	
	1.42 M 2-Mercaptoethanol	No known significant effects or critical hazards.
	XL1-Blue Competent Cells	No known significant effects or critical hazards.
Ingestion	: pUC18 Control Plasmid	No known significant effects or critical hazards.
	DNA	
	1.42 M 2-Mercaptoethanol	Toxic if swallowed.
	XL1-Blue Competent Cells	Toxic if swallowed.

Classification

Product/ingredient name	ACGIH	IARC	EPA	NIOSH	NTP	OSHA
XL1-Blue Competent Cells						
Sucrose	A4	-	-	-	-	-

Potential chronic health effects

Chronic effects	: Contains material that may cause target organ damage, based on animal data.
Carcinogenicity	: No known significant effects or critical hazards.
Mutagenicity	: No known significant effects or critical hazards.
Teratogenicity	: No known significant effects or critical hazards.
Developmental effects	: No known significant effects or critical hazards.
Fertility effects	: No known significant effects or critical hazards.

Over-exposure signs/symptoms

Inhalation	: No specific data.
Ingestion	: No specific data.
Skin	: No specific data.
Eyes	: No specific data.
Target organs	: pUC18 Control Plasmid Not available. DNA 1.42 M 2-Mercaptoethanol Not available. XL1-Blue Competent Cells Contains material which may cause damage to the following organs: blood, kidneys, gastrointestinal tract, upper respiratory tract, skin, central nervous system (CNS), eye, lens or cornea.
Other adverse effects	: pUC18 Control Plasmid Not available. DNA 1.42 M 2-Mercaptoethanol Not available. XL1-Blue Competent Cells Not available.

12 . Ecological information

Environmental effects : No known significant effects or critical hazards.

12 . Ecological information

Aquatic ecotoxicity

Product/ingredient name	Test	Result	Species	Exposure
Dimethyl sulfoxide	-	Acute LC50 35 to 37 ml/L Fresh water	Fish	96 hours
	-	Acute LC50 34000000 ug/L Fresh water	Fish	96 hours
Manganese dichloride	-	Acute EC50 4700 ug/L Fresh water	Daphnia	48 hours
Glycerol	-	Acute LC50 54 to 57 ml/L Fresh water	Fish	96 hours
Potassium chloride	-	Acute EC50 83000 ug/L Fresh water	Daphnia	48 hours
	-	Acute LC50 337 mg/L Fresh water	Daphnia	48 hours
	-	Acute LC50 435000 ug/L Fresh water	Fish	96 hours

Other adverse effects : No known significant effects or critical hazards.

13 . Disposal considerations

Waste disposal : The generation of waste should be avoided or minimized wherever possible. Dispose of surplus and non-recyclable products via a licensed waste disposal contractor. Disposal of this product, solutions and any by-products should at all times comply with the requirements of environmental protection and waste disposal legislation and any regional local authority requirements. Avoid dispersal of spilled material and runoff and contact with soil, waterways, drains and sewers.

Disposal should be in accordance with applicable regional, national and local laws and regulations. Local regulations may be more stringent than regional or national requirements.

The information presented below only applies to the material as supplied. The identification based on characteristic(s) or listing may not apply if the material has been used or otherwise contaminated. It is the responsibility of the waste generator to determine the toxicity and physical properties of the material generated to determine the proper waste identification and disposal methods in compliance with applicable regulations.

Refer to Section 7: HANDLING AND STORAGE and Section 8: EXPOSURE CONTROLS/PERSONAL PROTECTION for additional handling information and protection of employees.

14 . Transport information

Regulatory information

DOT / IMDG / IATA : Not regulated.

15 . Regulatory information

HCS Classification	: pUC18 Control Plasmid DNA	Not regulated.
	1.42 M 2-Mercaptoethanol	Toxic material Irritating material Sensitizing material
	XL1-Blue Competent Cells	Toxic material Target organ effects

15 . Regulatory information

	pUC18 Control Plasmid DNA	Not available.
	1.42 M 2-Mercaptoethanol	Not available.
	XL1-Blue Competent Cells	Contains material which may cause damage to the following organs: blood, kidneys, gastrointestinal tract, upper respiratory tract, skin, central nervous system (CNS), eye, lens or cornea.
U.S. Federal regulations	: pUC18 Control Plasmid DNA	United States inventory (TSCA 8b): All components are listed or exempted.
	1.42 M 2-Mercaptoethanol	United States inventory (TSCA 8b): All components are listed or exempted.
	XL1-Blue Competent Cells	United States inventory (TSCA 8b): All components are listed or exempted.
	pUC18 Control Plasmid DNA	SARA 302/304/311/312 extremely hazardous substances: No products were found. SARA 302/304 emergency planning and notification: No products were found. SARA 302/304/311/312 hazardous chemicals: No products were found. SARA 311/312 MSDS distribution - chemical inventory - hazard identification: No products were found.
	1.42 M 2-Mercaptoethanol	SARA 302/304/311/312 extremely hazardous substances: No products were found. SARA 302/304 emergency planning and notification: No products were found. SARA 302/304/311/312 hazardous chemicals: 2-Mercaptoethanol SARA 311/312 MSDS distribution - chemical inventory - hazard identification: 2-Mercaptoethanol: Fire hazard, Immediate (acute) health hazard, Delayed (chronic) health hazard
	XL1-Blue Competent Cells	SARA 302/304/311/312 extremely hazardous substances: No products were found. SARA 302/304 emergency planning and notification: No products were found. SARA 302/304/311/312 hazardous chemicals: Potassium chloride; Glycerol; Manganese dichloride; Sucrose; Dimethyl sulfoxide SARA 311/312 MSDS distribution - chemical inventory - hazard identification: Potassium chloride: Immediate (acute) health hazard, Delayed (chronic) health hazard; Glycerol: Immediate (acute) health hazard, Delayed (chronic) health hazard; Manganese dichloride: Delayed (chronic) health hazard; Sucrose: Delayed (chronic) health hazard; Dimethyl sulfoxide: Immediate (acute) health hazard, Delayed (chronic) health hazard
	pUC18 Control Plasmid DNA	Clean Water Act (CWA) 307: No products were found.
	1.42 M 2-Mercaptoethanol	Clean Water Act (CWA) 307: No products were found.
	XL1-Blue Competent Cells	Clean Water Act (CWA) 307: No products were found.
	pUC18 Control Plasmid DNA	Clean Water Act (CWA) 311: Edetic acid
	1.42 M 2-Mercaptoethanol	Clean Water Act (CWA) 311: No products were found.
	XL1-Blue Competent Cells	Clean Water Act (CWA) 311: No products were found.

15 . Regulatory information

pUC18 Control Plasmid DNA	Clean Air Act (CAA) 112 accidental release prevention: No products were found.
1.42 M 2-Mercaptoethanol	Clean Air Act (CAA) 112 accidental release prevention: No products were found.
XL1-Blue Competent Cells	Clean Air Act (CAA) 112 accidental release prevention: No products were found.
pUC18 Control Plasmid DNA	Clean Air Act (CAA) 112 regulated flammable substances : No products were found.
1.42 M 2-Mercaptoethanol	Clean Air Act (CAA) 112 regulated flammable substances : No products were found.
XL1-Blue Competent Cells	Clean Air Act (CAA) 112 regulated flammable substances : No products were found.
pUC18 Control Plasmid DNA	Clean Air Act (CAA) 112 regulated toxic substances: No products were found.
1.42 M 2-Mercaptoethanol	Clean Air Act (CAA) 112 regulated toxic substances: No products were found.
XL1-Blue Competent Cells	Clean Air Act (CAA) 112 regulated toxic substances: No products were found.

SARA 313

Form R - Reporting requirements

<u>Product name</u>	<u>CAS number</u>	<u>Concentration</u>
XL1-Blue Competent Cells		
Manganese dichloride	7773-01-5	5 - 10
Hexaamminecobalt trichloride	10534-89-1	0.1 - 1

Supplier notification

XL1-Blue Competent Cells		
Manganese dichloride	7773-01-5	5 - 10
Hexaamminecobalt trichloride	10534-89-1	0.1 - 1

SARA 313 notifications must not be detached from the MSDS and any copying and redistribution of the MSDS shall include copying and redistribution of the notice attached to copies of the MSDS subsequently redistributed.

State regulations

pUC18 Control Plasmid DNA	<p>Connecticut Carcinogen Reporting: None of the components are listed.</p> <p>Connecticut Hazardous Material Survey: None of the components are listed.</p> <p>Florida substances: None of the components are listed.</p> <p>Illinois Chemical Safety Act: None of the components are listed.</p> <p>Illinois Toxic Substances Disclosure to Employee Act: None of the components are listed.</p> <p>Louisiana Reporting: None of the components are listed.</p> <p>Louisiana Spill: None of the components are listed.</p> <p>Massachusetts Spill: None of the components are listed.</p> <p>Massachusetts Substances: None of the components are listed.</p> <p>Michigan Critical Material: None of the components are listed.</p> <p>Minnesota Hazardous Substances: None of the components are listed.</p> <p>New Jersey Hazardous Substances: None of the components are listed.</p> <p>New Jersey Spill: None of the components are listed.</p> <p>New Jersey Toxic Catastrophe Prevention Act: None of the components are listed.</p> <p>New York Acutely Hazardous Substances: None of the components are listed.</p> <p>New York Toxic Chemical Release Reporting: None of the components are listed.</p> <p>Pennsylvania RTK Hazardous Substances: None of the</p>
------------------------------	--

15 . Regulatory information

components are listed.

Rhode Island Hazardous Substances: None of the components are listed.

- 1.42 M 2-Mercaptoethanol
- Connecticut Carcinogen Reporting:** None of the components are listed.
Connecticut Hazardous Material Survey: None of the components are listed.
Florida substances: None of the components are listed.
Illinois Chemical Safety Act: None of the components are listed.
Illinois Toxic Substances Disclosure to Employee Act: None of the components are listed.
Louisiana Reporting: None of the components are listed.
Louisiana Spill: None of the components are listed.
Massachusetts Spill: None of the components are listed.
Massachusetts Substances: The following components are listed: 2-Mercaptoethanol
Michigan Critical Material: None of the components are listed.
Minnesota Hazardous Substances: None of the components are listed.
New Jersey Hazardous Substances: None of the components are listed.
New Jersey Spill: None of the components are listed.
New Jersey Toxic Catastrophe Prevention Act: None of the components are listed.
New York Acutely Hazardous Substances: None of the components are listed.
New York Toxic Chemical Release Reporting: None of the components are listed.
Pennsylvania RTK Hazardous Substances: The following components are listed: 2-Mercaptoethanol
Rhode Island Hazardous Substances: None of the components are listed.

XL1-Blue Competent Cells

- Connecticut Carcinogen Reporting:** None of the components are listed.
Connecticut Hazardous Material Survey: None of the components are listed.
Florida substances: None of the components are listed.
Illinois Chemical Safety Act: None of the components are listed.
Illinois Toxic Substances Disclosure to Employee Act: None of the components are listed.
Louisiana Reporting: None of the components are listed.
Louisiana Spill: None of the components are listed.
Massachusetts Spill: None of the components are listed.
Massachusetts Substances: The following components are listed: Glycerol;Sucrose
Michigan Critical Material: None of the components are listed.
Minnesota Hazardous Substances: None of the components are listed.
New Jersey Hazardous Substances: The following components are listed: Manganese dichloride
New Jersey Spill: None of the components are listed.
New Jersey Toxic Catastrophe Prevention Act: None of the components are listed.
New York Acutely Hazardous Substances: None of the components are listed.
New York Toxic Chemical Release Reporting: None of the

15 . Regulatory information

components are listed.

Pennsylvania RTK Hazardous Substances: The following components are listed: Glycerol; Manganese dichloride; Sucrose

Rhode Island Hazardous Substances: None of the components are listed.

State regulations - California Prop. 65 : No products were found.

16 . Other information

Label requirements	: pUC18 Control Plasmid DNA	NOT EXPECTED TO PRODUCE SIGNIFICANT ADVERSE HEALTH EFFECTS WHEN THE RECOMMENDED INSTRUCTIONS FOR USE ARE FOLLOWED.
	: 1.42 M 2-Mercaptoethanol	HARMFUL IF SWALLOWED. CAUSES EYE AND SKIN IRRITATION. MAY CAUSE ALLERGIC SKIN REACTION.
	: XL1-Blue Competent Cells	HARMFUL IF SWALLOWED. CONTAINS MATERIAL THAT MAY CAUSE TARGET ORGAN DAMAGE, BASED ON ANIMAL DATA.

Date of issue : 01/09/2009

Version : 1

Notice to reader

DISCLAIMER: This Material Safety Data Sheet is offered without charge to the clients of Agilent Technologies. Data is the most current available to Agilent Technologies at the time of preparation and is issued as a matter of information only, no warranty as to its accuracy or completeness is expressed or implied.

▣ Indicates information that has changed from previously issued version.



Office of Biohazard Containment and Safety
Science Branch, CFIA
59 Camelot Drive, Ottawa, Ontario K1A 0Y9
Tel: (613) 221-7068 Fax: (613) 228-6129
Email: ImportZoopath@inspection.gc.ca

Bureau du confinement des biorisques et sécurité
Direction générale des sciences, ACIA
59 promenade Camelot, Ottawa, Ontario K1A 0Y9
Tél: (613) 221-7068 Téléc: (613) 228-6129
Courriel: ImportZoopath@inspection.gc.ca

October 20th, 2009

Ms. Shamila Survery / Mr. Michael Decosimo
Cedarlane Laboratories Ltd
4410 Paletta Court
Burlington, Ontario L7L 5R2

By Facsimile: (289) 288-0020

SUBJECT: Importation of *Escherichia coli* strains

Dear Ms. Survery / Mr. Decosimo:

Our office received your query about the importation of *Escherichia coli* from the American Type Culture Collection (ATCC) located in Manassas, Virginia, United States. The following *Escherichia coli* strains are considered to be level 1 animal pathogens:

• 5K	• CIE85	• J52	• MC4100 (MuLac)	• U5/41
• 58	• DH1	• J53	• MG1655	• W208
• 58-161	• DH10 GOLD	• JC3272	• MM294	• W945
• 679	• DH10B	• JC7661	• MS101	• W1485
• 1532	• DH5	• JC9387	• NC-7	• W3104
• AB284	• DH5-alpha	• JF1504	• Nissle 1917	• W3110
• AB311	• DP50	• JF1508	• One Shot STBL3	• WA704
• AB1157	• DY145	• JF1509	• OP50	• WP2
• AB1206	• DY380	• JJ055	• P678	• X1854
• AG1	• E11	• JM83	• PA309	• X2160T
• B	• EJ183	• JM101	• PK-5	• X2541
• BB4	• EL250	• JM109	• PMC103	• X2547T
• BD792	• EMG2	• K12	• PR13	• XL1-BLUE
• BL21	• EPI 300	• KC8	• Rri	• XL1-BLUE-MRF
• BL21 (DE3)	• EZ10	• KA802	• RV308	• XL0LR
• BM25.8	• FDA Seattle 1946	• KAM32	• S17-1λ -PIR	• Y10
• C	• Fusion-Blue	• KAM33	• SCS1	• Y1090 (1090)
• C-1a	• H1443	• KAM43	• SMR10	• YN2980
• C-3000	• HF4714	• LE450	• SOLR	• W3110
• C25	• HB101	• LE451	• SuperchargeEZ10	• WG1
• C41 (DE3)	• HS(PFAMP)R	• LE452	• SURE	• WG439
• C43 (DE3)	• Hfr3000	• MB408	• TOP10	• WG443
• C600	• Hfr3000 X74	• MBX1928	• TG1	• WG445
• Cavalli Hfr	• HMS174	• MC1061		

The Office of Biohazard Containment and Safety (BCS) of the Canadian Food Inspection Agency (CFIA) only issues import permits for microorganisms that are pathogenic to animals, or parts of microorganisms that are pathogenic to animals. As the products listed above are not considered pathogenic to animals, the Office of BCS does not have any regulatory requirements for their importation.

Please note that other legislation may apply. You may wish to contact the Public Health Agency of Canada's (PHAC) Office of Laboratory Security at (613) 957-1779.

Note: Microorganisms pathogenic to animals and veterinary biologics require an import permit from the CFIA.

Sincerely,

Cinthia Labrie
Head, Animal Pathogen Importation Program
Office of Biohazard Containment & Safety

Bacteria

ATCC® Number: **BAA-1025™** [Order this Item](#) Price: **\$205.00**

Related Links ▶

[NCBI Entrez Search](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[Related Products](#)

Organism: *Escherichia coli* (Migula) Castellani and Chalmers

Designations: BL21

Depositor: J Bull

History: J Bull I J Molineux

[Biosafety Level:](#) 1

Shipped: freeze-dried

Growth Conditions: [ATCC medium 129](#): Nutrient agar with 0.5% NaCl

Temperature: 37.0°C

Duration: aerobic

Permits/Forms: In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Applications: bacteriophage host (host for bacteriophages T3 and T7)

Related Products: bacteriophage:ATCC [BAA-1025-B1](#)

bacteriophage:ATCC [BAA-1025-B2](#)

[BioProducts](#)

[Cell, microbial and molecular genomics products for the life sciences](#)

- [sciences](#)

[BioServices](#)

[Bio-materials management; basic repository to complex partnership-level](#)

- [services](#)

[BioStandards](#)

[Biological Reference Material and Consensus Standards for the life science](#)

- [community](#)

[Return to Top](#)



[ATCC Advanced Catalog Search](#) » [Product Details](#)

Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's [Material Transfer Agreement](#) or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Canada, China, Hong Kong, India, Israel, Japan, Korea, Macao, Mexico, New Zealand, Singapore, and Taiwan, R.O.C. must contact a [local distributor](#) for pricing information and to place an order for ATCC cultures and products.

[Print this Page](#)

Patent Depository

ATCC® Number:	PTA-1798™	Order this Item	Price:	\$200.00
Designation / Description:	Escherichia coli DH5alpha			Related Links
U.S. Patent Number:	7,354,755			▶
Biosafety Level:	1			NCBI Entrez Search
Shipped:	frozen			Make a Deposit
Permits/Forms:	In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location.			Frequently Asked Questions
				Material Transfer Agreement
				Technical Support

This material is cited in a U.S. and/or other Patent or Patent Application, and may not be used to infringe on the patent claims. ATCC is required to inform the Patent Depositor of the party to which the material was furnished.

[Return to Top](#)

Notices and Disclaimers

ATCC products are intended for laboratory research purposes only, unless noted otherwise. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for non-profit institutions in the United States. Cultures that are ordered as fast tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

[Back to my Search](#)



ATCC THE ESSENTIALS OF LIFE SCIENCE RESEARCH GLOBALLY DELIVERED

ATCC Advanced Catalog Search » Product Details

Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's Material Transfer Agreement or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Alaska, Canada, China, Hong Kong, India, Israel, Japan, Korea, Macao, Mexico, New Zealand, Singapore, and Taiwan, ATCC must contact a local distributor for pricing information and to place an order for ATCC cultures and products.

Print this Page

Patent Depository

ATCC® Number: PTA-1798™ Order this Item Price: \$200.00

Designation / Description: Escherichia coli DH5alpha

U.S. Patent Number: 7,354,755

Biosafety Level: 1

Shipped: frozen

Permits/Forms: In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location.

Related Links

- NCBI Entrez Search
Make a Deposit
Frequently Asked Questions
Material Transfer Agreement
Technical Support

This material is cited in a U.S. and/or other Patent or Patent Application, and may not be used to infringe on the patent claims. ATCC is required to inform the Patent Depositor of the party to which the material was furnished.

Return to Top

Notices and Disclaimers

ATCC products are intended for laboratory research purposes only, unless noted otherwise. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for non-profit institutions in the United States. Cultures that are obtained as sub-cultures or feeds will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

Back to my Search

Fungi ,Yeasts and Yeast Genetic Stock

ATCC® Number: **201389™** [Order this Item](#) Price: **\$155.00**

Organism: *Saccharomyces cerevisiae* Meyen ex E.C. Hansen deposited as *Saccharomyces cerevisiae* Hansen, teleomorph

Alternate State: *Candida robusta* Diddens et Lodder

Designations: BY4742 [ATCC 4040004, YVC1]

Depositors: JD Boeke

Biosafety Level: 1

Shipped: frozen

Genotype/ORF/
Gene Name: MATalpha his3delta1 leu2delta0 lys2delta0 ura3delta0 [21662] [53392]

Growth Conditions: ATCC medium 1245: YEPD
Temperature: 25.0°C

Permits/Forms: In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Applications: transformation host [53392]

Mating Type: alpha

Karyotype: Ploidy: haploid

Comments: Derived from S288C, ATCC 204508. parental strain for the International Systematic Saccharomyces cerevisiae Gene Disruption Project [21662]

Related Products: genomic DNA: ATCC [201389D-5](#)

Subcollection: Yeasts

References: 21662: Brachmann CB, et al. Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* 14: 115-132, 1998. PubMed: [9483801](#)

53392: Palmer CP, et al. A TRP homolog in *Saccharomyces cerevisiae* forms an intracellular Ca²⁺-permeable channel in the yeast vacuolar membrane. *Proc. Natl. Acad. Sci. USA* 98: 7801-7805, 2001. PubMed: [11427713](#)

Related Links ▶

- [NCBI Entrez Search](#)
- [Make a Deposit](#)
- [Frequently Asked Questions](#)
- [Material Transfer Agreement](#)
- [Technical Support](#)
- [Related Products](#)

BioProducts

- [Cell, microbial and molecular genomics products for the life sciences](#)

BioServices

- [Bio-materials management; basic repository to complex partnership-level services](#)

BioStandards

- [Biological Reference Material and Consensus Standards for the life science community](#)

[Return to Top](#)

Info for Cell Line(s)

Cell Biology

ATCC® Number: **CRL-1555™** Order this Item Price: **\$279.00**

Designations: A-431
 Depositors: DJ Giard, SA Aaronson
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)

Morphology:



Organ: skin

Source: **Tissue:** epidermis
Disease: epidermoid carcinoma

Permits/Forms:

In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

Tumorigenic: Yes

DNA Profile (STR): Amelogenin: X
 CSF1PO: 11,12
 D13S317: 9,13
 D16S539: 12,14
 D5S818: 12,13
 D7S820: 10
 TH01: 9
 TPOX: 11
 vWA: 15,17

Cytogenetic Analysis: This is a hypertriploid human cell line. The modal chromosome number was 74 occurring in 36% of cells. The rate of cells with higher ploidies was 1.0%.

Isoenzymes: AK-1, 1
 ES-D, 1
 G6PD, B
 GLO-I, 2
 Me-2, 0
 PGM1, 1
 PGM3, 1

Age: 85 years

Related Links ▶

[NCBI Entrez Search](#)

[Cell Micrograph](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[Related Cell Culture Products](#)

Login Required ▶

[Product Information Sheet](#)

[BioProducts](#)

[Cell, microbial and molecular genomics products for the life sciences](#)

[BioServices](#)

[Bio-materials management; basic repository to complex partnership-level services](#)

Gender:	female
Comments:	The epidermoid carcinoma cell line A-431, derived from an 85-year-old female, is one of a series of cell lines established from solid tumors by D.J. Giard, et al. [23218]
Propagation:	ATCC complete growth medium: The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%. Temperature: 37.0°C

BioStandards

- [Biological Reference Material and Consensus Standards for the life science community](#)

Protocol:

- | | |
|---------------|---|
| Subculturing: | <ol style="list-style-type: none">1. Remove and discard culture medium.2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin-0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.5. Add appropriate aliquots of the cell suspension to new culture vessels.6. Incubate cultures at 37°C. |
|---------------|---|

Subcultivation Ratio: A subcultivation ratio of 1:3 to 1:8 is recommended

Medium Renewal: Every 2 to 3 days

Preservation:	Freeze medium: Complete growth medium supplemented with 5% (v/v) DMSO Storage temperature: liquid nitrogen vapor phase
Related Products:	Recommended medium (without the additional supplements or serum described under ATCC Medium):ATCC 30-2002 recommended serum:ATCC 30-2020 derivative:ATCC CRL-2592

References:

- 23093: Faust JB, Meeker TC. Amplification and expression of the bcl-1 gene in human solid tumor cell lines. *Cancer Res.* 52: 2460-2463, 1992. PubMed: [1568216](#)
- 23218: Giard DJ, et al. In vitro cultivation of human tumors: establishment of cell lines derived from a series of solid tumors. *J. Natl. Cancer Inst.* 51: 1417-1423, 1973. PubMed: [4357758](#)
- 32507: Kovelman R, et al. Enhanced transcriptional activation by E2 proteins from the oncogenic human papillomaviruses. *J. Virol.* 70: 7549-7560, 1996. PubMed: [8892874](#)
- 32568: Lee JH, et al. The proximal promoter of the human transglutaminase 3 gene. *J. Biol. Chem.* 271: 4561-4568, 1996. PubMed: [8626812](#)
- 32582: Chang K, Pastan I. Molecular cloning of mesothelin, a differentiation antigen present on mesothelium, mesotheliomas, and ovarian cancers. *Proc. Natl. Acad. Sci. USA* 93: 136-140, 1996. PubMed: [8552591](#)
- 32912: Wizemann TM, et al. Peptide methionine sulfoxide reductase contributes to the maintenance of adhesins in three major pathogens. *Proc. Natl. Acad. Sci. USA* 93: 7985-7990, 1996. PubMed: [8755589](#)

[Return to Top](#)

Cell Biology

ATCC® Number: **CCL-185™** [Order this Item](#) Price: **\$279.00**

Designations: A549
 Depositors: M Lieber
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)
 epithelial

Morphology: 

Source: **Organ:** lung
Disease: carcinoma

Cellular Products: keratin
 In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Permits/Forms:

Isolation: **Isolation date:** 1972

Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

DNA Profile (STR): Amelogenin: X,Y
 CSF1PO: 10,12
 D13S317: 11
 D16S539: 11,12
 D5S818: 11
 D7S820: 8,11
 THO1: 8,9.3
 TPOX: 8,11
 vWA: 14

Cytogenetic Analysis: This is a hypotriploid human cell line with the modal chromosome number of 66, occurring in 24% of cells. Cells with 64 (22%), 65, and 67 chromosome counts also occurred at relatively high frequencies; the rate with higher ploidies was low at 0.4%. There were 6 markers present in single copies in all cells. They include der(6)t(1;6) (q11;q27); ?del(6) (p23); del(11) (q21), del(2) (q11), M4 and M5. Most cells had two X and two Y chromosomes. However, one or both Y chromosomes were lost in 40% of 50 cells analyzed. Chromosomes N2 and N6 had single copies per cell; and N12 and N17 usually had 4 copies.

Related Links ▶

- [NCBI Entrez Search](#)
- [Cell Micrograph](#)
- [Make a Deposit](#)
- [Frequently Asked Questions](#)
- [Material Transfer Agreement](#)
- [Technical Support](#)
- [Related Cell Culture Products](#)

Login Required ▶

- [Product Information Sheet](#)

BioProducts

- [Cell, microbial and molecular genomics products for the](#)
- [life sciences](#)

BioServices

- [Bio-materials management; basic repository to complex partnership-level](#)
- [services](#)

Isoenzymes: G6PD, B
 Age: 58 years
 Gender: male
 Ethnicity: Caucasian

[BioStandards](#)

[Biological Reference Material and Consensus Standards for the life science community](#)

Comments: This line was initiated in 1972 by D.J. Giard, et al. through explant culture of lung carcinomatous tissue from a 58-year-old Caucasian male. [23218]
 Further studies by M. Lieber, et al. revealed that A549 cells could synthesize lecithin with a high percentage of desaturated fatty acids utilizing the cytidine diphosphocholine pathway. [58030]
 The cells are positive for keratin by immunoperoxidase staining.

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated F-12K Medium, Catalog No. 30-2004. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.
Atmosphere: air, 95%; carbon dioxide (CO₂), 5%
Temperature: 37.0°C

Subculturing: **Protocol:**

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin-0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
 Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessels.
 Cultures can be established between 2 X 10⁽³⁾ and 1 X 10⁽⁴⁾ viable cells/cm². Do not exceed 7 X 10⁽⁴⁾ cels/cm².
6. Incubate cultures at 37°C.

Interval: Maintain cultures at a cell concentration between 6 X 10⁽³⁾ and 6 X 10⁽⁴⁾ cell/cm².
Subcultivation Ratio: A subcultivation ratio of 1:3 to 1:8 is recommended
Medium Renewal: 2 to 3 times per week

Preservation: **Freeze medium:** Complete growth medium supplemented with 5% (v/v) DMSO
Storage temperature: liquid nitrogen vapor phase

Doubling Time: about 22 hours

Related Products: Recommended medium (without the additional supplements or serum described under ATCC Medium):[ATCC 30-2004](#)
recommended serum:[ATCC 30-2020](#)

References:


- 23218: Giard DJ, et al. In vitro cultivation of human tumors: establishment of cell lines derived from a series of solid tumors. *J. Natl. Cancer Inst.* 51: 1417-1423, 1973. PubMed: [4357758](#)
- 27669: Mayr GA, Freimuth P. A single locus on human chromosome 21 directs the expression of a receptor for adenovirus type 2 in mouse A9 cells. *J. Virol.* 71: 412-418, 1997. PubMed: [8985365](#)
- 27819: Goodrum FD, Ornelles DA. The early region 1B 55-kilodalton oncoprotein of adenovirus relieves growth restrictions imposed on viral replication by the cell cycle. *J. Virol.* 71: 548-561, 1997. PubMed: [8985383](#)
- 32299: St. Geme JW, et al. Characterization of the genetic locus encoding Haemophilus influenzae type b surface fibrils. *J. Bacteriol.* 178: 6281-6287, 1996. PubMed: [8892830](#)
- 32347: Horikami SM, et al. The Sendai virus V protein interacts with the NP protein to regulate viral genome RNA replication. *Virology* 222: 383-390, 1996. PubMed: [8806522](#)
- 32351: Huang S, et al. Adenovirus interaction with distinct integrins mediates separate events in cell entry and gene delivery to hematopoietic cells. *J. Virol.* 70: 4502-4508, 1996. PubMed: [8676475](#)
- 32373: Goodrum FD, et al. Adenovirus early region 4 34-kilodalton protein directs the nuclear localization of the early region 1B 55-kilodalton protein in primate cells. *J. Virol.* 70: 6323-6335, 1996. PubMed: [8709260](#)
- 32394: Fang R, Aust AE. Induction of ferritin synthesis in human lung epithelial cells treated with crocidolite asbestos. *Arch. Biochem. Biophys.* 340: 369-375, 1997. PubMed: [9143343](#)
- 32488: Geiger T, et al. Antitumor activity of a PKC-alpha antisense oligonucleotide in combination with standard chemotherapeutic agents against various human tumors transplanted into nude mice. *Anticancer Drug Des.* 13: 35-45, 1998. PubMed: [9474241](#)
- 32496: Evdokiou A, Cowled PA. Tumor-suppressive activity of the growth arrest-specific gene GAS1 in human tumor cell lines. *Int. J. Cancer* 75: 568-577, 1998. PubMed: [9466658](#)
- 32511: Giavedoni LD, Yilma T. Construction and characterization of replication-competent simian immunodeficiency virus vectors that express gamma interferon. *J. Virol.* 70: 2247-2251, 1996. PubMed: [8642649](#)
- 32514: Bartz SR, et al. Human immunodeficiency virus type 1 cell cycle control: Vpr is cytostatic and mediates G2 accumulation by a mechanism which differs from DNA damage checkpoint control. *J. Virol.* 70: 2324-2331, 1996. PubMed: [8642659](#)
- 32722: Garofalo R, et al. Transcriptional activation of the interleukin-8 gene by respiratory syncytial virus infection in alveolar epithelial cells: nuclear translocation of the RelA transcription factor as a mechanism producing airway mucosal inflammation. *J. Virol.* 70: 8773-8781, 1996. PubMed: [8971006](#)
- 32758: Jamaluddin M. et al. Inducible translational regulation of

[Return to Top](#)

Cell Biology

ATCC® Number: **CRL-1573™** Order this Item Price: **\$279.00**

Designations: 293 [HEK-293]
 Depositors: FL Graham
Biosafety Level: 2 [CELLS CONTAIN ADENOVIRUS]
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)
 epithelial



Source: **Organ:** embryonic kidney
Cell Type: transformed with adenovirus 5 DNA
 In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Permits/Forms:

Restrictions: These cells are distributed for research purposes only. 293 cells, their products, or their derivatives may not be distributed to third parties.

Applications: efficacy testing [92587]
 transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))
 viruscide testing [92579]

Receptors: vitronectin, expressed

Tumorigenic: YES

DNA Profile (STR): Amelogenin: X
 CSF1PO: 11,12
 D13S317: 12,14
 D16S539: 9,13
 D5S818: 8,9
 D7S820: 11,12
 THO1: 7,9.3
 TPOX: 11
 vWA: 16,19

Related Links ▶

- [NCBI Entrez Search](#)
- [Cell Micrograph](#)
- [Make a Deposit](#)
- [Frequently Asked Questions](#)
- [Material Transfer Agreement](#)
- [Technical Support](#)
- [Related Cell Culture Products](#)

Login Required ▶

[Product Information Sheet](#)

BioProducts

- [Cell, microbial and molecular genomics products for the life sciences](#)

BioServices

[Bio-materials management; basic repository to complex partnership-level services](#)

Cytogenetic Analysis: This is a hypotriploid human cell line. The modal chromosome number was 64, occurring in 30% of cells. The rate of cells with higher ploidies was 4.2 %. The der(1)t(1;15) (q42;q13), der(19)t(3;19) (q12;q13), der(12)t(8;12) (q22;p13), and four other marker chromosomes were common to most cells. Five other markers occurred in some cells only. The marker der(1) and M8 (or Xq+) were often paired. There were four copies of N17 and N22. Noticeably in addition to three copies of X chromosomes, there were paired Xq+, and a single Xp+ in most cells.

Age: fetus

Comments: Although an earlier report suggested that the cells contained Adenovirus 5 DNA from both the right and left ends of the viral genome [RF32764], it is now clear that only left end sequences are present. [39768]

The line is excellent for titrating human adenoviruses. The cells express an unusual cell surface receptor for vitronectin composed of the integrin beta-1 subunit and the vitronectin receptor alpha-v subunit. [23406] The Ad5 insert was cloned and sequenced, and it was determined that a colinear segment from nts 1 to 4344 is integrated into chromosome 19 (19q13.2). [39768]

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated Eagle's Minimum Essential Medium, Catalog No. 30-2003. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.
Atmosphere: air, 95%; carbon dioxide (CO2), 5%
Temperature: 37.0°C
The cell line does not adhere to the substrate when left at room temperature for any length of time, therefore, live cultures may be received with the cells detached. The cells will re-attach to the flask over a period of several days in culture at 37C.

[BioStandards](#)

- [Biological Reference Material and Consensus Standards for the life science community](#)

Protocol:

Subculturing:

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin-0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessels. An inoculum of 2×10^3 to 6×10^3 viable cells/cm² is recommended.
6. Incubate cultures at 37°C. 6. Subculture when cell concentration is between 6 and 7×10^4 cells/cm².

Subcultivation Ratio: 1:10 to 1:20 weekly.

Medium Renewal: Every 2 to 3 days

Preservation:

Freeze medium: Complete growth medium supplemented with 5% (v/v) DMSO

Storage temperature: liquid nitrogen vapor phase

Related Products:

Recommended medium (without the additional supplements or serum described under ATCC Medium): ATCC [30-2003](#)

derivative: ATCC [CRL-10852](#)

derivative: ATCC [CRL-12006](#)

derivative: ATCC [CRL-12007](#)

derivative: ATCC [CRL-12013](#)

derivative: ATCC [CRL-12479](#)

derivative: ATCC [CRL-2029](#)

derivative: ATCC [CRL-2368](#)

purified DNA: ATCC [CRL-1573D](#)

- 21624: Xie QW, et al. Complementation analysis of mutants of nitric oxide synthase reveals that the active site requires two hemes. *Proc. Natl. Acad. Sci. USA* 93: 4891-4896, 1996. PubMed: [8643499](#)
- 21631: Da Costa LT, et al. Converting cancer genes into killer genes. *Proc. Natl. Acad. Sci. USA* 93: 4192-4196, 1996. PubMed: [8633039](#)
- 22282: Graham FL, et al. Characteristics of a human cell line transformed by DNA from human adenovirus type 5. *J. Gen. Virol.* 36: 59-72, 1977. PubMed: [886304](#)
- 22319: Graham FL, et al. Defective transforming capacity of adenovirus type 5 host-range mutants. *Virology* 86: 10-21, 1978. PubMed: [664220](#)
- 22699: Harrison T, et al. Host-range mutants of adenovirus type 5 defective for growth in HeLa cells. *Virology* 77: 319-329, 1977. PubMed: [841862](#)
- 23406: Bodary SC, McLean JW. The integrin beta 1 subunit associates with the vitronectin receptor alpha v subunit to form a novel vitronectin receptor in a human embryonic kidney cell line. *J. Biol. Chem.* 265: 5938-5941, 1990. PubMed: [1690718](#)
- 27819: Goodrum FD, Ornelles DA. The early region 1B 55-kilodalton oncoprotein of adenovirus relieves growth restrictions imposed on viral replication by the cell cycle. *J. Virol.* 71: 548-561, 1997. PubMed: [8985383](#)
- 28301: Loffler S, et al. CD9, a tetraspan transmembrane protein, renders cells susceptible to canine distemper virus. *J. Virol.* 71: 42-49, 1997. PubMed: [8985321](#)
- 32283: Hu SX, et al. Development of an adenovirus vector with tetracycline-regulatable human tumor necrosis factor alpha gene expression. *Cancer Res.* 57: 3339-3343, 1997. PubMed: [9269991](#)
- 32396: Kolanus W, et al. alphaLbeta2 integrin/LFA-1 binding to ICAM-1 induced by cytohesin-1 a cytoplasmic regulatory molecule. *Cell* 86: 233-242, 1996. PubMed: [8706128](#)
- 32490: Stauderman KA, et al. Characterization of human recombinant neuronal nicotinic acetylcholine receptor subunit combinations alpha 2 beta 4, alpha 3 beta 4 and alpha 4 beta 4 stably expressed in HEK293 cells. *J. Pharmacol. Exp. Ther.* 284: 777-789, 1998. PubMed: [9454827](#)
- 32514: Bartz SR, et al. Human immunodeficiency virus type 1 cell cycle control: Vpr is cytostatic and mediates G2 accumulation by a mechanism which differs from DNA damage checkpoint control. *J. Virol.* 70: 2324-2331, 1996. PubMed: [8642659](#)
- 32726: Sandri-Goldin RM, Hibbard MK. The herpes simplex virus type 1 regulatory protein ICP27 coimmunoprecipitates with anti-sm antiserum, and the C terminus appears to be required for this interaction. *J. Virol.* 70: 108-118, 1996. PubMed: [8523514](#)
- 32829: Ansieau S, et al. Tumor necrosis factor receptor-associated factor (TRAF)-1, TRAF-2, and TRAF-3 interact in vivo with the CD30 cytoplasmic domain: TRAF-2 mediates

[Return to Top](#)

Cell Biology

ATCC® Number: **CCL-2™** Order this Item Price: **\$279.00**

Designations: HeLa
 Depositors: WF Scherer
Biosafety Level: 2 [Cells contain human papilloma virus]
 Shipped: frozen
 Medium & Serum: See Propagation
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)

Morphology:



Source: **Organ:** cervix
Disease: adenocarcinoma
Cell Type: epithelial
 keratin

Cellular Products: Lysophosphatidylcholine (lyso-PC) induces AP-1 activity and c-jun N-terminal kinase activity (JNK1) by a protein kinase C-independent pathway [26623]
 In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location.

Permits/Forms:

Applications: transfection host ([21491] Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents)
 screening for Escherichia coli strains with invasive potential [21447] [21491]

Virus Susceptibility: Human adenovirus 3
 Encephalomyocarditis virus
 Human poliovirus 1
 Human poliovirus 2
 Human poliovirus 3

DNA Profile (STR): Amelogenin: X
 CSF1PO: 9,10
 D13S317: 12,13.3
 D16S539: 9,10
 D5S818: 11,12
 D7S820: 8,12
 TH01: 7
 TPOX: 8,12
 vWA: 16,18

Related Links ▶

- [NCBI Entrez Search](#)
- [Cell Micrograph](#)
- [Make a Deposit](#)
- [Frequently Asked Questions](#)
- [Material Transfer Agreement](#)
- [Technical Support](#)
- [Related Cell Culture Products](#)

Login Required ▶

- [Product Information Sheet](#)

BioProducts

- [Cell, microbial and molecular genomics products for the life sciences](#)

BioServices

- [Bio-materials management; basic repository to complex partnership-level services](#)

Modal number = 82; range = 70 to 164.

There is a small telocentric chromosome in 98% of the cells. 100% aneuploidy in 1385 cells examined. Four typical HeLa marker chromosomes have been reported in the literature. HeLa Marker Chromosomes: One copy of M1, one copy of M2, four-five copies of M3, and two copies of M4 as revealed by G-banding patterns. M1 is a rearranged long arm and centromere of chromosome 1 and the long arm of chromosome 3. M2 is a combination of short arm of chromosome 3 and long arm of chromosome 5. M3 is an isochromosome of the short arm of chromosome 5. M4 consists of the long arm of chromosome 11 and an arm of chromosome 19.

[BioStandards](#)

[Biological Reference Material and Consensus Standards for the life science community](#)

Cytogenetic Analysis:

Isoenzymes:	G6PD, A
Age:	31 years adult
Gender:	female
Ethnicity:	Black
HeLa Markers:	Y

Comments:

The cells are positive for keratin by immunoperoxidase staining. HeLa cells have been reported to contain human papilloma virus 18 (HPV-18) sequences. P53 expression was reported to be low, and normal levels of pRB (retinoblastoma suppressor) were found.

Propagation:

ATCC complete growth medium: The base medium for this cell line is ATCC-formulated Eagle's Minimum Essential Medium, Catalog No. 30-2003. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.
Atmosphere: air, 95%; carbon dioxide (CO₂), 5%
Temperature: 37.0°C

Protocol:

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin-0.53 mM EDTA solution to remove all traces of serum which contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessels.
6. Incubate cultures at 37°C.

Subculturing:

Subcultivation Ratio: A subcultivation ratio of 1:2 to 1:6 is recommended

Medium Renewal: 2 to 3 times per week

Preservation:

Freeze medium: Complete growth medium supplemented with 5% (v/v) DMSO

Storage temperature: liquid nitrogen vapor phase

Related Products:

Recommended medium (without the additional supplements or serum described under ATCC Medium): [ATCC 30-2003](#)

recommended serum: [ATCC 30-2020](#)

also available as Certified Reference Material, ATCC

[CRM-CCL-2](#)

derivative: [ATCC CCL-2.1](#)

derivative: [ATCC CCL-2.2](#)

derivative: [ATCC CCL-2.3](#)

- 21447: American Public Health Association. Compendium of methods for the microbiological examination of foods. 3rd ed. Washington, DC: American Public Health Association; 1992.
- 21491: AOAC International Invasiveness by *Escherichia coli* of mammalian cells, microbiological method. Gaithersburg, MD:AOAC International;AOAC "Official Methods of Analysis of the AOAC International" 982.36.
- 21626: Baldi A, et al. Genomic structure of the human retinoblastoma-related Rb2/p130 gene. Proc. Natl. Acad. Sci. USA 93: 4629-4632, 1996. PubMed: [8643454](#)
- 22148: Gey GO, et al. Tissue culture studies of the proliferative capacity of cervical carcinoma and normal epithelium. Cancer Res. 12: 264-265, 1952.
- 22263: Chen TR. Re-evaluation of HeLa, HeLa S3, and HEP-2 karyotypes. Cytogenet. Cell Genet. 48: 19-24, 1988. PubMed: [3180844](#)
- 22766: Boshart M, et al. A new type of papillomavirus DNA, its presence in genital cancer biopsies and in cell lines derived from cervical cancer. EMBO J. 3: 1151-1157, 1984. PubMed: [6329740](#)
- 22767: Schneider-Gadicke A, Schwarz E. Different human cervical carcinoma cell lines show similar transcription patterns of human papillomavirus type 18 early genes. EMBO J. 5: 2285-2292, 1986. PubMed: [3023067](#)
- 22919: Schwarz E, et al. Structure and transcription of human papillomavirus sequences in cervical carcinoma cells. Nature 314: 111-114, 1985. PubMed: [2983228](#)
- 22995: Pater MM, Pater A. Human papillomavirus types 16 and 18 sequences in carcinoma cell lines of the cervix. Virology 145: 313-318, 1985. PubMed: [2992153](#)
- 23180: Yee C, et al. Presence and expression of human papillomavirus sequences in human cervical carcinoma cell lines. Am. J. Pathol. 119: 361-366, 1985. PubMed: [2990217](#)
- 23324: Scheffner M, et al. The state of the p53 and retinoblastoma genes in human cervical carcinoma cell lines. Proc. Natl. Acad. Sci. USA 88: 5523-5527, 1991. PubMed: [1648218](#)
- 25915: Jones HW Jr., et al. George Otto Gey. (1899-1970). The HeLa cell and a reappraisal of its origin. Obstet. Gynecol. 38: 945-949, 1971. PubMed: [4942173](#)
- 25919: Scherer WF, Hoogasian AF. Preservation at subzero temperatures of mouse fibroblasts (strain L) and human epithelial cells (strain HeLa). Proc. Soc. Exp. Biol. Med. 87: 480-487, 1954. PubMed: [13237281](#)
- 25921: Scherer WF, et al. Studies on the propagation in vitro of poliomyelitis viruses. IV. Viral multiplication in a stable strain of human malignant epithelial cells (strain HeLa) derived from an epidermoid carcinoma of the cervix. J. Exp. Med. 97: 695-710, 1953. PubMed: [13052828](#)
- 26623: Fang X, et al. Lysophosphatidylcholine stimulates activator protein 1 and the c-Jun N-terminal kinase activity. J.

[Return to Top](#)

PRODUCT: HeLa Tet-Off® Cell Line

CATALOG No.
630905

LOT NUMBER
Specified on product label.

STORAGE CONDITIONS
• Store cells in liquid nitrogen (−196°C) or in −150°C freezer.
• Store FBS in −20°C freezer.

STORAGE MEDIUM
70% Tet System Approved FBS, 20% DMEM & 10% DMSO

SHELF LIFE
1 year from date of receipt under proper storage conditions

SHIPPING CONDITIONS
Dry ice (−70°C)

FOR RESEARCH USE ONLY

DESCRIPTION:

Human cervical epithelioid carcinoma-derived cell line that expresses the tetracycline-controlled transactivator (tTA). An inducible, high-level gene expression system can be generated by stably transfecting these cells with a plasmid or infecting these cells with a virus that expresses the gene of interest under the control of a suitable Tet Response Element (TRE).

CELL TYPE INFORMATION & RECOMMENDED CELL CULTURE MEDIUM
See back page.

PACKAGE CONTENTS:

- 1 ml HeLa Tet-Off Cell Line (2.0 x 10⁶ cells/tube)
- 0.5 ml CHO-AA8-Luc Tet-Off Control Cell Line (1.0 x 10⁶ cells/tube)
- 50 ml Tet System Approved FBS

OTHER

- Tet Systems User Manual (PT3001-1)
- Tet Systems Protocol-at-a-Glance (PT3001-2)

QUALITY CONTROL DATA:

Functional Tests

HeLa Tet-Off Cells were transiently transfected by electroporation with pTRE-Luc. Luciferase activity in the presence and absence of 2 µg/ml doxycycline (Cat. No. 631311) was measured 48 hr later as described in the Tet Systems User Manual. Induction was observed to be at least 400-fold when cells were grown in media containing Tet System Approved FBS (Cat. No. 631101 or 631106).

Luciferase activity in the presence and absence of 2 µg/ml doxycycline was measured after 48 hr for the CHO-AA8-Luc Tet-Off Control Cell Line. Induction was found to be at least 1,000-fold when cells were grown in media containing Tet System Approved FBS (Cat. No. 631101 or 631106).

Mycoplasma Contamination Test

These lots of the two cell lines were tested and found to be free of *Mycoplasma* contamination.

REFERENCE:

1. Gossen, M. & Bujard, H. (1992) Tight control of gene expression in mammalian cells by tetracycline responsive promoters. *Proc. Natl. Acad. Sci. USA* **89**:5547-5551.

APPROVED BY: 

(PA641627)

CELL TYPE INFORMATION

HeLa Tet-Off Cell Line: Human cervical epithelioid carcinoma cell line transformed with pUHD15-1 and pSV2neo. This cell line is neomycin resistant.

CHO-AA8-Luc Tet-Off Control Cell Line: Chinese hamster ovary stably transformed with pUHD15-1, pSV2neo, pTRE-Luc, and pTK-Hyg. This cell line is neomycin and hygromycin resistant.

RECOMMENDED CELL CULTURE MEDIUM

HeLa Tet-Off Cell Line: 90% Dulbecco's Modified Eagle's Medium (DMEM), 10% Tet System Approved Fetal Bovine Serum (FBS), 4 mM L-glutamine, 100 µg/ml G418, 100 units/ml penicillin G sodium & 100 µg/ml streptomycin sulfate.

CHO-AA8-Luc Tet-Off Control Cell Line: 90% Eagle Minimum Essential Medium (alpha modification), 10% Tet System Approved Fetal Bovine Serum (FBS), 4 mM L-glutamine, 100 µg/ml G418, 100 units/ml penicillin G sodium, 100 µg/ml streptomycin sulfate & 100 µg/ml hygromycin B.

ADDITIONAL NOTES

We recommend that you titrate each new lot of hygromycin to determine its optimal concentration due to the lot-to-lot variation in antibiotic activity.

The doubling time for HeLa Tet-Off cells is approximately 20 hours during log phase.

The morphology of HeLa Tet-Off cells is elongated and adherent with 2-3 filopodia.

General Notice to Purchaser

Clontech products are to be used for research purposes only. They may not be used for any other purpose, including, but not limited to, use in drugs, in vitro diagnostic purposes, therapeutics, or in humans. Clontech products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without written approval of Clontech Laboratories, Inc.

Use of the Tetracycline controllable expression systems (the "Tet Technology") is covered by a series of patents including U.S. patents # 5,464,758 and #5,814,618, which are proprietary to TET Systems Holding GmbH & Co. KG. Academic research institutions are granted an automatic license with the purchase of this product to use the Tet Technology only for internal, academic research purposes, which license specifically excludes the right to sell, or otherwise transfer, the Tet Technology or its component parts to third parties. Notwithstanding the above, academic and not-for profit research institutions whose research using the Tet Technology is sponsored by for profit organizations, which shall receive ownership to all data and results stemming from the sponsored research, shall need a commercial license agreement from IP Merchandisers in order to use the Tet Technology. In accepting this license, all users acknowledge that the Tet Technology is experimental in nature. TET Systems Holding GmbH & Co. KG makes no warranties, express or implied or of any kind, and hereby disclaims any warranties, representations, or guarantees of any kind as to the Tet Technology, patents, or products. All others are invited to request a license from TET Systems Holding GmbH & Co. KG prior to purchasing these reagents or using them for any purpose. Clontech is required by its licensing agreement to submit a report of all purchasers of the Tet-controllable expression system to IP Merchandisers, Inc. For license information, please contact:

Hans Peter Kneubuehl
TET Systems Holding GmbH & Co. KG
Im Neuenheimer Feld 582
69120 Heidelberg
Germany

Tel +49 6221 588 04 00
Fax +49 6221 588 04 04
eMail: kneubuehl@tet-systems.de
or use our electronic licensing request form via
http://www.tetsystems.com/main_inquiry.htm

Clontech, Clontech logo and all other trademarks are the property of Clontech Laboratories, Inc.
Clontech is a Takara Bio Company. ©2006

Cell Biology

ATCC® Number: **CCL-2.2™** Order this Item Price: **\$279.00**

Designations: HeLa S3
 Depositors: TT Puck
Biosafety Level: 2 [Cells contain human papilloma virus (HPV-18)]
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)
 Morphology: epithelial

Source: **Organ:** cervix
Disease: adenocarcinoma

Cellular Products: keratin
 In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Isolation: **Isolation date:** 1955

Applications: transfection host ([Roche FuGENE® Transfection Reagents technology from amaxa](#))

DNA Profile (STR): Amelogenin: X
 CSF1PO: 9,10
 D13S317: 13.3
 D16S539: 9,10
 D5S818: 11,12
 D7S820: 8,12
 THO1: 7
 TPOX: 8,12
 vWA: 16,18

Cytogenetic Analysis: A medium-sized metacentric marker is present in 100% of the cells. HeLa Markers: One copy of M1, one copy of M2, two copies of M3, and one copy of M4.

Isoenzymes: G6PD, A

Age: 31 years

Gender: female

Ethnicity: Black

HeLa Markers: Y

- Related Links ▶**
[NCBI Entrez Search](#)
[Make a Deposit](#)
[Frequently Asked Questions](#)
[Material Transfer Agreement](#)
[Technical Support](#)
[Related Cell Culture Products](#)

- Login Required ▶**
[Product Information Sheet](#)

BioProducts

- [Cell, microbial and molecular genomics products for the](#)
 • [life sciences](#)

BioServices

- [Bio-materials management; basic repository to complex partnership-level](#)
 • [services](#)

HeLa S3 is a clonal derivative of the parent HeLa line (see ATCC [CCL-2](#)). S3 was cloned in 1955 by T.T. Puck, P.I. Marcus, and S.J. Cieciura. [[22814](#)]

The HeLa S3 clone has been very useful in the clonal analysis of mammalian cell populations relating to chromosomal variation, cell nutrition, and plaque-forming ability.

This line can be adapted to grow in suspension. [[25952](#)]

The cells are positive for keratin by immunoperoxidase staining. A culture at approximately passage 400 was submitted to the American Type Culture Collection in February, 1972.

HeLa cells have been reported to contain human papilloma virus 18 (HPV-18) sequences. [[23180](#)]

[BioStandards](#)

[Biological](#)

[Reference](#)

[Material and](#)

[Consensus](#)

[Standards for](#)

[the life science](#)

• [community](#)

Comments:

Propagation:

ATCC complete growth medium: The base medium for this cell line is ATCC-formulated F-12K Medium, Catalog No. 30-2004. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.

Atmosphere: air, 95%; carbon dioxide (CO₂), 5%

Temperature: 37.0°C

Protocol:

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin-0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessels.
6. Incubate cultures at 37°C.

Subculturing:

Subcultivation Ratio: A subcultivation ratio of 1:4 to 1:10 is recommended

Medium Renewal: 2 to 3 times per week

Preservation:

Freeze medium: Complete growth medium supplemented with 5% (v/v) DMSO

Storage temperature: liquid nitrogen vapor phase

Related Products:

Recommended medium (without the additional supplements or serum described under ATCC Medium): ATCC [30-2004](#)
recommended serum: ATCC [30-2020](#)

References:

- 22263: Chen TR. Re-evaluation of HeLa, HeLa S3, and HEP-2 karyotypes. *Cytogenet. Cell Genet.* 48: 19-24, 1988. PubMed: [3180844](#)
- 22766: Boshart M, et al. A new type of papillomavirus DNA, its presence in genital cancer biopsies and in cell lines derived from cervical cancer. *EMBO J.* 3: 1151-1157, 1984. PubMed: [6329740](#)
- 22814: Puck TT, et al. Clonal growth of mammalian cells in vitro; growth characteristics of colonies from single HeLa cells with and without a feeder layer. *J. Exp. Med.* 103: 273-283, 1956. PubMed: [13286432](#)
- 23180: Yee C, et al. Presence and expression of human papillomavirus sequences in human cervical carcinoma cell lines. *Am. J. Pathol.* 119: 361-366, 1985. PubMed: [2990217](#)
- 25929: Puck TT, Marcus PI. A rapid method for viable cell titration and clone production with HeLa cells in tissue culture: the use of x-irradiated cells to supply conditioning factors. *Proc. Natl. Acad. Sci. USA* 41: 432-437, 1955.
- 25932: . . *J. Exp. Med.* 104: 427-434, 1956.
- 25934: . . *Methods Enzymol.* 5: 90-119, 1962.
- 25952: Darnell JE Jr., et al. The effect of cell population density on the amino acid requirements for poliovirus synthesis in HeLa cells. *J. Exp. Med.* 110: 445-450, 1959. PubMed: [13814142](#)
- 25953: Cohen EP, Eagle H. A simplified chemostat for the growth of mammalian cells: characteristics of cell growth in continuous culture. *J. Exp. Med.* 113: 467-474, 1961.
- 25957: Darnell JE Jr., Sawyer TK. Variation in plaque-forming ability among parental and clonal strains of HeLa cells. *Virology* 8: 223-229, 1959. PubMed: [13669339](#)
- 25959: Sato G, et al. Molecular growth requirements of single mammalian cells. *Science* 126: 461-464, 1957. PubMed: [13486039](#)
- 32358: Soares K, et al. cis-Acting elements involved in transcriptional regulation of the herpes simplex virus type 1 latency-associated promoter 1 (LAP1) in vitro and in vivo. *J. Virol.* 70: 5384-5394, 1996. PubMed: [8764049](#)
- 32524: Chang YE, et al. Properties of the protein encoded by the UL32 open reading frame of herpes simplex virus 1. *J. Virol.* 70: 3938-3946, 1996. PubMed: [8648731](#)
- 32966: Jiang BH, et al. Hypoxia-inducible factor 1 levels vary exponentially over a physiologically relevant range of O₂ tension. *Am. J. Physiol.* 271: C1172-C1180, 1996. PubMed: [8897823](#)
- 33006: Genuario RR, Perry RP. The GA-binding protein can serve as both an activator and repressor of ribosomal protein gene transcription. *J. Biol. Chem.* 271: 4388-4395, 1996. PubMed: [8626789](#)
- 33032: Ladner RD, et al. Identification of a consensus cyclin-dependent kinase phosphorylation site unique to the nuclear form of human deoxyuridine triphosphate nucleotidohydrolase. *J. Biol. Chem.* 271: 7752-7757, 1996. PubMed: [8631817](#)
- 33033: Ladner RD, et al. Characterization of distinct nuclear

[Return to Top](#)

Cell Biology

ATCC® Number: **TIB-152™** Order this Item Price: **\$279.00**

Designations: Jurkat, Clone E6-1

Depositors: A Weiss

Biosafety Level: 1

Shipped: frozen

Medium & Serum: See Propagation

Growth Properties: suspension

Organism: *Homo sapiens* (human)
lymphoblast

Morphology: 

Source: **Disease:** acute T cell leukemia
Cell Type: T lymphocyte;

Cellular Products: interleukin-2 (interleukin 2, IL-2) [1609]

Permits/Forms: In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location.

Applications: transfection host (Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents)

Receptors: T cell antigen receptor, expressed

Antigen Expression: CD3; Homo sapiens, expressed

Amelogenin: X,Y

CSF1PO: 11,12

D13S317: 8,12

D16S539: 11

DNA Profile (STR): D5S818: 9

D7S820: 8,12

THO1: 6,9.3

TPOX: 8,10

vWA: 18

Cytogenetic Analysis: This is a pseudodiploid human cell line. The modal chromosome number is 46, occurring in 74% with polyploidy at 5.3%. The karyotype is 46,XY,-2,-18,del(2) (p21p23),del(18) (p11.2). Most cells had normal X and Y chromosomes.

Gender: male

Related Links ▶

[NCBI Entrez Search](#)

[Cell Micrograph](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[Related Cell Culture Products](#)

Login Required ▶

[Product Information Sheet](#)

BioProducts

[Cell, microbial and molecular genomics products for the life sciences](#)

BioServices

[Bio-materials management; basic repository to complex partnership-level services](#)

Comments: This is a clone of the Jurkat-FHCRC cell line, a derivative of the Jurkat cell line. [1609]
The Jurkat cell line was established from the peripheral blood of a 14 year old boy by Schneider et al., and was originally designated JM. [50685] [112530]
Clone E6-1 cells produce large amounts of IL-2 after stimulation with phorbol esters and either lectins or monoclonal antibodies against the T3 antigen (both types of stimulants are needed to induce IL-2 production. [1609]
The line was cloned from cells obtained from Dr. Kendall Smith and are mycoplasma free. [1609]

BioStandards

- [Biological Reference Material and Consensus Standards for the life science community](#)

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated RPMI-1640 Medium, Catalog No. 30-2001. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.
Atmosphere: air, 95%; carbon dioxide (CO₂), 5%
Temperature: 37.0°C

Subculturing: **Protocol:** Cultures can be maintained by the addition of fresh medium or replacement of medium. Alternatively, cultures can be established by centrifugation with subsequent resuspension at 1 X 10⁽⁵⁾ viable cells/ml. Do not allow the cell density to exceed 3 X 10⁽⁶⁾ cells/ml.
Interval: Maintain cultures at a cell concentraion between between 1 X 10⁽⁵⁾ and 1 X 10⁽⁶⁾ viable cells/ml.
Medium Renewal: Add fresh medium every 2 to 3 days (depending on cell density)

Preservation: **Freeze medium:** Complete growth medium supplemented with 5% (v/v) DMSO
Storage temperature: liquid nitrogen vapor phase

Doubling Time: 48 hrs

Related Products: Recommended medium (without the additional supplements or serum described under ATCC Medium):ATCC [30-2001](#)
recommended serum:ATCC [30-2020](#)
derivative:ATCC [CRL-1990](#)
derivative:ATCC [CRL-2063](#)
derivative:ATCC [TIB-153](#)

1609: Weiss A, et al. The role of T3 surface molecules in the activation of human T cells: a two-stimulus requirement for IL-2 production reflects events occurring at a pre-translational level. *J. Immunol.* 133: 123-128, 1984. PubMed: [6327821](#)

23430: Gillis S, Watson J. Biochemical and biological characterization of lymphocyte regulatory molecules. V. Identification of an interleukin 2-producing human leukemia T cell line. *J. Exp. Med.* 152: 1709-1719, 1980. PubMed: [6778951](#)

32253: Berninghausen O, Leippe M. Necrosis versus apoptosis as the mechanism of target cell death induced by *Entamoeba histolytica*. *Infect. Immun.* 65: 3615-3621, 1997. PubMed: [9284127](#)

32368: Churchill MJ, et al. The rev-responsive element negatively regulates human immunodeficiency virus type 1 env mRNA expression in primate cells. *J. Virol.* 70: 5786-5790, 1996. PubMed: [8709194](#)

32396: Kolanus W, et al. alphaLbeta2 integrin/LFA-1 binding to ICAM-1 induced by cytohesin-1 a cytoplasmic regulatory molecule. *Cell* 86: 233-242, 1996. PubMed: [8706128](#)

32446: Gan W, Rhoads RE. Internal initiation of translation directed by the 5'-untranslated region of the mRNA for eIF4G, a factor involved in the picornavirus-induced switch from cap-dependent to internal initiation. *J. Biol. Chem.* 271: 623-626, 1996. PubMed: [8557663](#)

32561: Tiffany HL, et al. Enhanced expression of the eosinophil-derived neurotoxin ribonuclease (RNS2) gene requires interaction between the promoter and intron. *J. Biol. Chem.* 271: 12387-12393, 1996. PubMed: [8647842](#)

32704: Chan YJ, et al. Synergistic interactions between overlapping binding sites for the serum response factor and ELK-1 proteins mediate both basal enhancement and phorbol ester responsiveness of primate cytomegalovirus. *J. Virol.* 70: 8590-8605, 1996. PubMed: [8970984](#)

32755: Kung SH, Medveczky PG. Identification of a herpesvirus saimiri cis-acting DNA fragment that permits stable replication of episomes in transformed T cells. *J. Virol.* 70: 1738-1744, 1996. PubMed: [8627695](#)

32796: Bloom TJ, Beavo JA. Identification and tissue-specific expression of PDE7 phosphodiesterase splice variants. *Proc. Natl. Acad. Sci. USA* 93: 14188-14192, 1996. PubMed: [8943082](#)

32901: Li YM, et al. Molecular identity and cellular distribution of advanced glycation endproduct receptors: relationship of p60 to OST-48 and p90 to 80K-H membrane proteins. *Proc. Natl. Acad. Sci. USA* 93: 11047-11052, 1996. PubMed: [8855306](#)

32904: Linette GP, et al. Cross talk between cell death and cell cycle progression: BCL-2 regulates NFAT-mediated activation. *Proc. Natl. Acad. Sci. USA* 93: 9545-9552, 1996. PubMed: [8790367](#)

32913: Miranda L, et al. Isolation of the human PC6 gene encoding the putative host protease for HIV-1 gp160 processing

References:

[Return to Top](#)

Cell Biology

ATCC® Number: **CRL-1740™** Order this Item Price: **\$279.00**

Designations: LNCaP clone FGC
 Depositors: JS Horoszewicz
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent, single cells and loosely attached clusters
 Organism: *Homo sapiens* (human)
 epithelial

Morphology: 

Organ: prostate
Disease: carcinoma

Source: **Derived from metastatic site:** left supraclavicular lymph node

Cellular Products: human prostatic acid phosphatase; prostate specific antigen [21889]

Permits/Forms: In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Restrictions: Distribution of this material for commercial purposes will require execution of a Non-exclusive License Agreement. At the time of placing an order, customers must send a request to licensing@ATCC.org. Orders will be shipped when Customer Service receives confirmation from our Licensing officer.

Isolation: **Isolation date:** 1977

Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

Receptors: androgen receptor, positive; estrogen receptor, positive [23045]

Tumorigenic: Yes

DNA Profile (STR): Amelogenin: X,Y
 CSF1PO: 10,11
 D13S317: 10,12
 D16S539: 11
 D5S818: 11,12
 D7S820: 9.1,10.3
 TH01: 9
 TPOX: 8,9
 vWA: 16,18

Related Links ▶

- [NCBI Entrez Search](#)
- [Cell Micrograph](#)
- [Make a Deposit](#)
- [Frequently Asked Questions](#)
- [Material Transfer Agreement](#)
- [Technical Support](#)
- [Related Cell Culture Products](#)

Login Required ▶

- [Product Information Sheet](#)

BioProducts

- [Cell, microbial and molecular genomics products for the life sciences](#)

BioServices

- [Bio-materials management; basic repository to complex partnership-level services](#)

Cytogenetic Analysis:	This is a hypotetraploid human cell line. The modal chromosome number was 84, occurring in 22% of cells. However, cells with chromosome counts of 86 (20%) and 87 (18%) also occurred at high frequencies. The rate of cells with higher ploidies was 6.0%.
Age:	50 years adult
Gender:	male
Ethnicity:	Caucasian
Comments:	<p>LNCaP clone FGC was isolated in 1977 by J.S. Horoszewicz, et al., from a needle aspiration biopsy of the left supraclavicular lymph node of a 50-year-old Caucasian male (blood type B+) with confirmed diagnosis of metastatic prostate carcinoma. [21889]</p> <p>These cells are responsive to 5-alpha-dihydrotestosterone (growth modulation and acid phosphatase production). [23045]</p> <p>The cells do not produce a uniform monolayer, but grow in clusters which should be broken apart by repeated pipetting when subcultures are prepared.</p> <p>They attach only lightly to the substrate, do not become confluent and rapidly acidify the medium.</p> <p>Growth is very slow.</p> <p>The cells should be allowed to incubate undisturbed for the first 48 hours after subculture.</p> <p>When flask cultures are shipped, the majority of the cells become detached from the flask and float in the medium. Upon receipt, incubate the flask (in the usual position for monolayer cultures) for 24 to 48 hours to allow the cells to re-attach.</p> <p>The medium can then be removed and replaced with fresh medium.</p> <p>If desired, the contents of the flask can be collected, centrifuged at 300 X g for 15 minutes, resuspended in 10 ml of medium and dispensed into a single flask.</p>
Propagation:	<p>ATCC complete growth medium: The base medium for this cell line is ATCC-formulated RPMI-1640 Medium, Catalog No. 30-2001. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.</p> <p>Atmosphere: air, 95%; carbon dioxide (CO₂), 5%</p> <p>Temperature: 37.0°C</p>

BioStandards

- Biological Reference Material and Consensus Standards for the life science community

Subculturing:	<p>Protocol:</p> <ol style="list-style-type: none">1. Remove and discard culture medium.2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin-0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.5. Add appropriate aliquots of the cell suspension to new culture vessels. Maintain cultures at a cell concentration between 1×10^4 and 2×10^5 cells/cm².6. Incubate cultures at 37°C. <p>Subcultivation Ratio: A subcultivation ratio of 1:3 to 1:6 is recommended</p> <p>Medium Renewal: Twice per week</p>
Preservation:	<p>Freeze medium: Complete growth medium supplemented with 5% (v/v) DMSO</p> <p>Storage temperature: liquid nitrogen vapor phase</p>
Doubling Time:	about 34 hours
Related Products:	<p>Recommended medium (without the additional supplements or serum described under ATCC Medium):ATCC 30-2001</p> <p>recommended serum:ATCC 30-2020</p> <p>derivative:ATCC CRL-10995</p> <p>purified DNA:ATCC CRL-1740D</p>

References:


- 21889: . Models for prostate cancer. 37New York: Liss; 1980.
- 22410: Gibas Z, et al. A high-resolution study of chromosome changes in a human prostatic carcinoma cell line (LNCaP). Cancer Genet. Cytogenet. 11: 399-404, 1984. PubMed: [6584201](#)
- 23045: Horoszewicz JS, et al. LNCaP model of human prostatic carcinoma. Cancer Res. 43: 1809-1818, 1983. PubMed: [6831420](#)
- 32283: Hu SX, et al. Development of an adenovirus vector with tetracycline-regulatable human tumor necrosis factor alpha gene expression. Cancer Res. 57: 3339-3343, 1997. PubMed: [9269991](#)
- 33090: Boffa LC, et al. Invasion of the CAG triplet repeats by a complementary peptide nucleic acid inhibits transcription of the androgen receptor and TATA-binding protein genes and correlates with refolding of an active nucleosome containing a unique AR gene sequence. J. Biol. Chem. 271: 13228-13233, 1996. PubMed: [8662737](#)

[Return to Top](#)

Cell Biology

ATCC® Number: **HTB-22™** Order this Item Price: **\$279.00**

Designations: MCF7
 Depositors: CM McGrath
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)
 epithelial



Organ: mammary gland; breast

Disease: adenocarcinoma

Source: **Derived from metastatic site:** pleural effusion

Cell Type: epithelial

Cellular Products: insulin-like growth factor binding proteins (IGFBP) BP-2; BP-4; BP-5

Permits/Forms: In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

Receptors: estrogen receptor, expressed

Antigen Expression: Blood Type O; Rh+

DNA Profile (STR):
 Amelogenin: X
 CSF1PO: 10
 D13S317: 11
 D16S539: 11,12
 D5S818: 11,12
 D7S820: 8,9
 TH01: 6
 TPOX: 9,12
 vWA: 14,15

Related Links ▶

[NCBI Entrez Search](#)

[Cell Micrograph](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[Related Cell Culture Products](#)

Login Required ▶

[Product Information Sheet](#)

[BioProducts](#)

- [Cell, microbial and molecular genomics products for the life sciences](#)

[BioServices](#)

- [Bio-materials management; basic repository to complex partnership-level services](#)

Cytogenetic Analysis: modal number = 82; range = 66 to 87. The stemline chromosome numbers ranged from hypertriploidy to hypotetraploidy, with the 2S component occurring at 1%. There were 29 to 34 marker chromosomes per S metaphase; 24 to 28 markers occurred in at least 30% of cells, and generally one large submetacentric (M1) and 3 large subtelocentric (M2, M3, and M4) markers were recognizable in over 80% of metaphases. No DM were detected. Chromosome 20 was nullisomic and X was disomic.

Isoenzymes: AK-1, 1
ES-D, 1-2
G6PD, B
GLO-I, 1-2
PGM1, 1-2
PGM3, 1

Age: 69 years adult

Gender: female

Ethnicity: Caucasian

Comments: The MCF7 line retains several characteristics of differentiated mammary epithelium including ability to process estradiol via cytoplasmic estrogen receptors and the capability of forming domes. The cells express the WNT7B oncogene [PubMed: 8168088]. Growth of MCF7 cells is inhibited by tumor necrosis factor alpha (TNF alpha). Secretion of IGFBP's can be modulated by treatment with anti-estrogens.

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated Eagle's Minimum Essential Medium, Catalog No. 30-2003. To make the complete growth medium, add the following components to the base medium: 0.01 mg/ml bovine insulin; fetal bovine serum to a final concentration of 10% .
Atmosphere: air, 95%; carbon dioxide (CO2), 5%
Temperature: 37.0°C

BioStandards

Biological Reference Material and Consensus Standards for the life science

- community

Protocol: Volumes used in this protocol are for 75 sq cm flasks; proportionally reduce or increase amount of dissociation medium for culture vessels of other sizes.

Note: if floating cells are present, it is recommended that they be transferred at the first two (2) subcultures as described below. It is not necessary to transfer floating cells for subsequent subcultures.

Subculturing:

1. Remove culture medium to a centrifuge tube.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin - 0.53 mM EDTA solution to remove all traces of serum which contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37C to facilitate dispersal.
4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Transfer the cell suspension to the centrifuge tube with the medium and cells from step 1, and centrifuge at approximately 125 xg for 5 to 10 minutes. Discard the supernatant.
6. Resuspend the cell pellet in fresh growth medium. Add appropriate aliquots of the cell suspension to new culture vessels.
7. Incubate cultures at 37C.

Subcultivation Ratio: A subcultivation ratio of 1:3 to 1:6 is recommended

Medium Renewal: 2 to 3 times per week

Preservation:	Freeze medium: Complete growth medium supplemented with 5% (v/v) DMSO Storage temperature: liquid nitrogen vapor phase
Doubling Time:	29 hrs
Related Products:	Recommended medium (without the additional supplements or serum described under ATCC Medium): ATCC 30-2003 recommended serum: ATCC 30-2020 purified DNA: ATCC HTB-22D purified RNA: ATCC HTB-22R 0.25% (w/v) Trypsin - 0.53 mM EDTA in Hank' BSS (w/o Ca++, Mg++): ATCC 30-2101 Cell culture tested DMSO: ATCC 4-X

21405: Sugarman BJ, et al. Recombinant human tumor necrosis factor-alpha: effects on proliferation of normal and transformed cells in vitro. *Science* 230: 943-945, 1985. PubMed: [3933111](#)

22871: Takahashi K, Suzuki K. Association of insulin-like growth-factor-I-induced DNA synthesis with phosphorylation and nuclear exclusion of p53 in human breast cancer MCF-7 cells. *Int. J. Cancer* 55: 453-458, 1993. PubMed: [8375929](#)

23046: Brandes LJ, Hermonat MW. Receptor status and subsequent sensitivity of subclones of MCF-7 human breast cancer cells surviving exposure to diethylstilbestrol. *Cancer Res.* 43: 2831-2835, 1983. PubMed: [6850594](#)

23079: Lan MS, et al. Polypeptide core of a human pancreatic tumor mucin antigen. *Cancer Res.* 50: 2997-3001, 1990. PubMed: [2334903](#)

23107: Pratt SE, Pollak MN. Estrogen and antiestrogen modulation of MCF7 human breast cancer cell proliferation is associated with specific alterations in accumulation of insulin-like growth factor-binding proteins in conditioned media. *Cancer Res.* 53: 5193-5198, 1993. PubMed: [7693333](#)

23113: Huguet EL, et al. Differential expression of human Wnt genes 2, 3, 4, and 7B in human breast cell lines and normal and disease states of human breast tissue. *Cancer Res.* 54: 2615-2621, 1994. PubMed: [8168088](#)

23217: Soule HD, et al. A human cell line from a pleural effusion derived from a breast carcinoma. *J. Natl. Cancer Inst.* 51: 1409-1416, 1973. PubMed: [4357757](#)

25065: Bellet D, et al. Malignant transformation of nontrophoblastic cells is associated with the expression of chorionic gonadotropin beta genes normally transcribed in trophoblastic cells. *Cancer Res.* 57: 516-523, 1997. PubMed: [9012484](#)

32275: Littlewood-Evans AJ, et al. The osteoclast-associated protease cathepsin K is expressed in human breast carcinoma. *Cancer Res.* 57: 5386-5390, 1997. PubMed: [9393764](#)

32278: Komarova EA, et al. Intracellular localization of p53 tumor suppressor protein in gamma-irradiated cells is cell cycle regulated and determined by the nucleus. *Cancer Res.* 57: 5217-5220, 1997. PubMed: [9393737](#)

32285: van Dijk MA, et al. A functional assay in yeas for the human estrogen receptor displays wild-type and variant estrogen receptor messenger RNAs present in breast carcinoma. *Cancer Res.* 57: 3478-3485, 1997. PubMed: [9270016](#)

32288: Landers JE, et al. Translational enhancement of mdm2 oncogene expression in human tumor cells containing a stabilized wild-type p53 protein. *Cancer Res.* 57: 3562-3568, 1997. PubMed: [9270029](#)

32344: Umekita Y, et al. Human prostate tumor growth in athymic mice: inhibition by androgens and stimulation by finasteride. *Proc. Natl. Acad. Sci. USA* 93: 11802-11807, 1996. PubMed: [8876218](#)

32467: Zamora-Leon SP, et al. Expression of the fructose transporter GLUT5 in human breast cancer. *Proc. Natl. Acad.*

References:

[Return to Top](#)

Cell Biology

ATCC® Number: **CRL-1469™** Order this Item Price: **\$279.00**

Designations: PANC-1
 Depositors: M Lieber
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)
 Morphology: epithelial

Related Links ▶
[NCBI Entrez Search](#)
[Make a Deposit](#)
[Frequently Asked Questions](#)
[Material Transfer Agreement](#)
[Technical Support](#)
[Related Cell Culture Products](#)

Source: **Organ:** pancreas
Tissue: duct
Disease: epithelioid carcinoma
 In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Permits/Forms:

Login Required ▶
[Product Information Sheet](#)

Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

[BioProducts](#)
[Cell, microbial and molecular genomics products for the life sciences](#)

Amelogenin: X
 CSF1PO: 10,12
 D13S317: 11
 D16S539: 11
 DNA Profile (STR): D5S818: 11,13
 D7S820: 8,10
 TH01: 7,8
 TPOX: 8,11
 vWA: 15

[BioServices](#)
[Bio-materials management; basic repository to complex partnership-level services](#)

Cytogenetic Analysis: Chromosome studies indicate a modal number of 63 with 3 distinct marker chromosomes and a small ring chromosome. This is a hypertriploid human cell line. The modal chromosome number was 61, occurring in 32% of cells., However, cells with 63 chromosomes also occurred at a high frequency (22%). The rate of cells with higher ploidies was 8.5%.

Isoenzymes: G6PD, B
 Age: 56 years
 Gender: male
 Ethnicity: Caucasian
 Comments: Growth is inhibited by 1 unit/ml L-asparaginase. The cells will grow in soft agar.

Propagation:	<p>ATCC complete growth medium: The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.</p> <p>Temperature: 37.0°C</p> <p>Atmosphere: air, 95%; carbon dioxide (CO₂), 5%</p>
Subculturing:	<p>Subcultivation Ratio: A subcultivation ratio of 1:2 to 1:4 is recommended</p> <p>Medium Renewal: 2 to 3 times per week</p> <p>Remove medium, and rinse with 0.25% trypsin, 0.53mM EDTA solution. Remove the solution and add an additional 1 to 2 ml of trypsin-EDTA solution. Allow the flask to sit at room temperature (or at 37C) until the cells detach. Add fresh culture medium, aspirate and dispense into new culture flasks.</p>
Preservation:	culture medium 95%; DMSO, 5%
Doubling Time:	52 hrs
Related Products:	<p>Recommended medium (without the additional supplements or serum described under ATCC Medium):ATCC 30-2002</p> <p>recommended serum:ATCC 30-2020</p>
References:	<p>22850: Lieber M, et al. Establishment of a continuous tumor-cell line (panc-1) from a human carcinoma of the exocrine pancreas. Int. J. Cancer 15: 741-747, 1975. PubMed: 1140870</p> <p>22859: Wu MC, et al. Mechanism of sensitivity of cultured pancreatic carcinoma to asparaginase. Int. J. Cancer 22: 728-733, 1978. PubMed: 363626</p> <p>23079: Lan MS, et al. Polypeptide core of a human pancreatic tumor mucin antigen. Cancer Res. 50: 2997-3001, 1990. PubMed: 2334903</p>

[BioStandards](#)

[Biological Reference Material and Consensus Standards for the life science community](#)

[Return to Top](#)

In the interest of conserving resources, we are no longer shipping manuals with products. Please visit www.clontech.com/manuals to obtain an electronic version.

PRODUCT: Saos-2 Tet-Off[®] Cell Line

CATALOG No. 630911

(Not sold separately)

LOT NUMBER

Specified on product label

STORAGE CONDITIONS

Liquid nitrogen vapor phase

STORAGE MEDIUM

70% Tet System Approved FBS,
20% DMEM, and 10% DMSO

SHELF LIFE

1 year from date of receipt under
proper storage conditions

SHIPPING CONDITIONS

Dry ice (-70°C)

DESCRIPTION

Saos-2 Tet-Off is a human osteosarcoma-derived cell line that expresses the tetracycline-regulated transactivator Tet-Off (1). Inducible expression of any gene can be achieved by transfecting or transducing this cell line with a vector containing your gene of interest under the control of a tetracycline-responsive promoter. Expression is induced by the withdrawal of doxycycline (Dox) from the culture medium.

**CELL TYPE INFORMATION &
RECOMMENDED CELL CULTURE MEDIUM**

See back of page.

PACKAGE CONTENTS

- 1 ml Saos-2 Tet-Off Cell Line
(2.0 x 10⁶ cells/tube)

OTHER

- Tet-Off and Tet-On Gene Expression Systems User Manual (PT3001-1)
- Tet Cell Lines Protocol-at-a-Glance (PT3001-2)

FOR RESEARCH USE ONLY

QUALITY CONTROL DATA

Functional Test

Saos-2 Tet-Off Cells were transiently transfected by electroporation with pTRE2-Luc. Luciferase activity in the presence and absence of 1 µg/ml doxycycline (Cat No. 631311) was measured 48 hr later as described in the Tet Systems User Manual. Induction was observed to be at least 30-fold when cells were grown in medium containing Clontech's Tet System Approved FBS.

Mycoplasma Contamination Test

This lot of cells has been tested and found to be free of mycoplasma contamination.



Clontech

United States/Canada
800.662.2566

Asia Pacific
+1.650.919.7300

Europe
+33.(0)1.3904.6880

Japan
+81.(0)77543.6116

Clontech Laboratories, Inc.
A Takara Bio Company
1290 Terra Bella Ave.
Mountain View, CA 94043
Technical Support (US)
E-mail: tech@clontech.com
www.clontech.com

APPROVED BY: _____

(PA993392)

CELL TYPE INFORMATION

Saos-2 Tet-Off is a human osteosarcoma-derived cell line stably transfected with pTet-Off (pUHD15-1neo). This cell line is G418 resistant.

RECOMMENDED CELL CULTURE MEDIUM

Grow the cells in 90% Dulbecco's Modified Eagle's Medium (DMEM), 10% Tet System Approved Fetal Bovine Serum (FBS), 4 mM L-glutamine, 100 µg/ml G418, 100 units/ml penicillin G sodium, and 100 µg/ml streptomycin sulfate, in the presence of 5% CO₂.

ADDITIONAL NOTES

The Saos-2 Tet-Off Cell Line exhibits hygromycin resistance. We therefore recommend use of puromycin resistance vectors (e.g., pTRE2 with pPUR or pTRE2-pur) for secondary transfections in this cell line.

REFERENCE

1. Gossen, M. & Bujard, H. (1992) *Proc. Natl. Acad. Sci. USA* **89**(12):5547-5551.

Notice to Purchaser

Clontech products are to be used for research purposes only. They may not be used for any other purpose, including, but not limited to, use in drugs, *in vitro* diagnostic purposes, therapeutics, or in humans. Clontech products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without written approval of Clontech Laboratories, Inc.

Use of the Tetracycline controllable expression systems (the Tet Technology) is covered by a series of patents including U.S. patents # 5,464,758 and #5,814,618, which are proprietary to TET Systems Holding GmbH & Co. KG. Academic research institutions are granted an automatic license with the purchase of this product to use the Tet Technology only for internal, academic research purposes, which license specifically excludes the right to sell, or otherwise transfer, the Tet Technology or its component parts to third parties. Notwithstanding the above, academic and not-for profit research institutions whose research using the Tet Technology is sponsored by for profit organizations, which shall receive ownership to all data and results stemming from the sponsored research, shall need a commercial license agreement from IP Merchandisers in order to use the Tet Technology. In accepting this license, all users acknowledge that the Tet Technology is experimental in nature. TET Systems Holding GmbH & Co. KG makes no warranties, express or implied or of any kind, and hereby disclaims any warranties, representations, or guarantees of any kind as to the Tet Technology, patents, or products. All others are invited to request a license from TET Systems Holding GmbH & Co. KG prior to purchasing these reagents or using them for any purpose. Clontech is required by its licensing agreement to submit a report of all purchasers of the Tet-controllable expression system to IP Merchandisers, Inc. For license information, please contact:

Attn: GSF/CEO
TET Systems Holding GmbH & Co. KG
Im Neuenheimer Feld 582
69120 Heidelberg
Germany
Tel +49 6221 588 04 00
Fax +49 6221 588 04 04
eMail: info@tetsystems.com
or use our electronic licensing request form via
http://www.tetsystems.com/main_inquiry.htm

Clontech, the Clontech logo and all other trademarks are the property of Clontech Laboratories, Inc., unless noted otherwise. Clontech is a Takara Bio Company. ©2009 Clontech Laboratories, Inc.

Cell Biology

ATCC® Number: **HTB-96™** Order this Item Price: **\$279.00**

Designations: U-2 OS
 Depositors: Hellstrom
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Homo sapiens* (human)
 Morphology: epithelial

Source: **Organ:** bone
Disease: osteosarcoma

Cellular Products: osteosarcoma derived growth factor (ODGF)
 In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

Receptors: insulin-like growth factor I (IGF-I); insulin-like growth factor II (IGF II)

Antigen Expression: Blood Type A; Rh+; HLA A2, Aw30, B12, Bw35, B40(+/-)

Amelogenin: X
 CSF1PO: 13
 D13S317: 13
 D16S539: 11,12
 DNA Profile (STR): D5S818: 11
 D7S820: 11,12
 THO1: 6,9.3
 TPOX: 11,12
 vWA: 14,18

Cytogenetic Analysis: Cell line U-2 OS is chromosomally highly altered, with chromosome counts in the hypertriploid range. We did not find the hypodiploid cell population described by J. Ponten, et al.,. Instead, most of the population has slightly higher counts than first described. Very few normal chromosomes are present, but a high number of stable marker chromosomes are identified., Different chromosomal rearrangements involving the same chromosomes (N1, N7, N9, and N11 particularly), are seen. Twenty-two markers are found including: t(9qter--->9q21::1p36--->1p::?), 7p+, iso(17q), t(15q;?), 4q+, del(3)(q21), 5q(aberrant) and others. [\[22509\]](#)

Related Links ▶

- [NCBI Entrez Search](#)
- [Make a Deposit](#)
- [Frequently Asked Questions](#)
- [Material Transfer Agreement](#)
- [Technical Support](#)
- [Related Cell Culture Products](#)

Login Required ▶

- [Product Information Sheet](#)

BioProducts

- [Cell, microbial and molecular genomics products for the life sciences](#)

BioServices

- [Bio-materials management; basic repository to complex partnership-level services](#)

Isoenzymes:	AK-1, 1 ES-D, 1 G6PD, B GLO-I, 2 PGM1, 2 PGM3, 1
Age:	15 years
Gender:	female
Ethnicity:	Caucasian
Comments:	J. Ponten and E. Saksela derived this line (originally 2T) in 1964 from a moderately differentiated sarcoma of the tibia of a 15 year old girl. Viruses were not detected during co-cultivation with WI-38 cells or in CF tests against SV40, RSV or adenoviruses. Mycoplasma contamination was detected and eliminated in 1972.
Propagation:	ATCC complete growth medium: The base medium for this cell line is ATCC-formulated McCoy's 5a Medium Modified, Catalog No. 30-2007. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%. Temperature: 37.0°C
Subculturing:	Subcultivation Ratio: A subcultivation ratio of 1:3 to 1:6 is recommended Medium Renewal: 2 to 3 times per week Remove medium, and rinse with 0.25% trypsin, 0.03% EDTA solution. Remove the solution and add an additional 1 to 2 ml of trypsin-EDTA solution. Allow the flask to sit at room temperature (or at 37C) until the cells detach. Add fresh culture medium, aspirate and dispense into new culture flasks.
Preservation:	Culture medium, 95%; DMSO, 5%
Related Products:	recommended serum: ATCC 30-2020

BioStandards

- [Biological Reference Material and Consensus Standards for the life science community](#)

References:

- 22237: Heldin CH, et al. A human osteosarcoma cell line secretes a growth factor structurally related to a homodimer of PDGF A-chains. *Nature* 319: 511-514, 1986. PubMed: [3456080](#)
- 22509: Ponten J, Saksela E. Two established in vitro cell lines from human mesenchymal tumours. *Int. J. Cancer* 2: 434-447, 1967. PubMed: [6081590](#)
- 23011: Raile K, et al. Human osteosarcoma (U-2 OS) cells express both insulin-like growth factor-I (IGF-I) receptors and insulin-like growth factor-II/mannose-6-phosphate (IGF-II/M6P) receptors and synthesize IGF-II: autocrine growth stimulation by IGF-II via the IGF-I receptor. *J. Cell. Physiol.* 159: 531-541, 1994. PubMed: [8188767](#)
- 32288: Landers JE, et al. Translational enhancement of mdm2 oncogene expression in human tumor cells containing a stabilized wild-type p53 protein. *Cancer Res.* 57: 3562-3568, 1997. PubMed: [9270029](#)
- 32308: Moradpour D, et al. Characterization of cell lines allowing tightly regulated expression of hepatitis C virus core protein. *Virology* 222: 51-63, 1996. PubMed: [8806487](#)

[Return to Top](#)

Cell Biology

ATCC® Number: **CL-173™** [Order this Item](#) Price: **\$279.00**

Designations: 3T3-L1
 Depositors: Massachusetts Institute of Technology
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: [See Propagation](#)
 Growth Properties: adherent
 Organism: *Mus musculus* (mouse)
 fibroblast

Morphology: 

Source: **Organ:** embryo
Cell Type: fibroblast

Cellular Products: triglycerides [\[3491\]](#)
 In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Permits/Forms:

Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

Receptors: insulin, expressed

Age: embryo

L1 is a continuous substrain of 3T3 (Swiss albino) developed through clonal isolation. The cells undergo a pre-adipose to adipose like conversion as they progress from a rapidly dividing to a confluent and contact inhibited state. A high serum content in the medium enhances fat accumulation [PubMed ID: 4426090].

Comments: Tested and found negative for ectromelia virus (mousepox). This line is also designated as ATCC CCL-92.1. ATCC CL-173 was deposited in 1974 without passage number information from the depositor. At the time of submission, ATCC prepared approximately 30 vials of seed stock at about 4 passages beyond the original depositor material (passage number: unknown +4).

Related Links ▶

- [NCBI Entrez Search](#)
- [Cell Micrograph](#)
- [Make a Deposit](#)
- [Frequently Asked Questions](#)
- [Material Transfer Agreement](#)
- [Technical Support](#)
- [Related Cell Culture Products](#)

Login Required ▶

- [Product Information Sheet](#)

BioProducts

- [Cell, microbial and molecular genomics products for the life sciences](#)

BioServices

- [Bio-materials management; basic repository to complex partnership-level services](#)

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium: bovine calf serum to a final concentration of 10%.
Atmosphere: air, 95%; carbon dioxide (CO₂), 5%
Temperature: 37.0°C
Growth Conditions: The serum used is important in culturing this line. Calf serum is recommended and not fetal bovine serum.

BioStandards

Biological Reference Material and Consensus Standards for the life science

- community

Protocol: Never allow culture to become completely confluent.

- Subculturing:**
1. Remove and discard culture medium.
 2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin - 0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.
 3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37C to facilitate dispersal.
 4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
 5. Add appropriate aliquots of the cell suspension to new culture vessels.
The recommended inoculum is 2 to 3 X 10⁽³⁾ cells/sq. cm. Subculture before cultures become 70 to 80% confluent or when cells reach 5 to 6 X10⁽⁴⁾ viable cells/sq. cm.
 6. Incubate cultures at 37C.

Interval: Every three days

Medium Renewal: 2 to 3 times per week

Preservation: **Freeze medium:** Complete growth medium supplemented with 5% (v/v) DMSO
Storage temperature: liquid nitrogen vapor phase

Doubling Time: 14 hrs

Related Products: Recommended medium (without the additional supplements or serum described under ATCC Medium):ATCC [30-2002](#) formerly distributed as:ATCC CCL-92.1
0.25% (w/v) Trypsin - 0.53 mM EDTA in Hank' BSS (w/o Ca⁺⁺, Mg⁺⁺):ATCC [30-2101](#)
Cell culture tested DMSO:ATCC [4-X](#)
Recommended serum: ATCC [30-2030](#)

886: Green H, Meuth M. An established pre-adipose cell line and its differentiation in culture. *Cell* 3: 127-133, 1974.

PubMed: [4426090](#)

3491: Green H. Triglyceride-accumulating clonal cell line. US Patent 4,003,789 dated Jan 18 1977

32373: Goodrum FD, et al. Adenovirus early region 4 34-kilodalton protein directs the nuclear localization of the early region 1B 55-kilodalton protein in primate cells. *J. Virol.*

70: 6323-6335, 1996. PubMed: [8709260](#)

32455: Scherer PE, et al. Identification, sequence, and expression of caveolin-2 defines a caveolin gene family. *Proc. Natl. Acad. Sci. USA* 93: 131-135, 1996. PubMed: [8552590](#)

32787: Kallen CB, Lazar MA. Antidiabetic thiazolidinediones inhibit leptin (ob) gene expression in 3T3-L1 adipocytes. *Proc. Natl. Acad. Sci. USA* 93: 5793-5796, 1996. PubMed: [8650171](#)

References:

[Return to Top](#)

Cell Biology

ATCC® Number: **CRL-1772™** Order this Item Price: **\$279.00**

Designations: C2C12

Biosafety Level: 1

Shipped: frozen

Medium & Serum: [See Propagation](#)

Growth Properties: adherent

Organism: *Mus musculus* (mouse)

myoblast

Morphology:



Tissue: muscle

Source: **Strain:** C3H

Cell Type: myoblast;

Permits/Forms:

In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Applications:

transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

This is a subclone (produced by H. Blau, et al) of the mouse myoblast cell line established by D. Yaffe and O. Saxel. [\[22903\]](#)

Comments:

The C2C12 cell line differentiates rapidly, forming contractile myotubes and producing characteristic muscle proteins. [\[22953\]](#) Treatment with bone morphogenic protein 2 (BMP-2) cause a shift in the differentiation pathway from myoblastic to osteoblastic. [\[23427\]](#)

Tested and found negative for ectromelia virus (mousepox).

Propagation:

ATCC complete growth medium: The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.

Temperature: 37.0°C

Related Links ▶

[NCBI Entrez Search](#)

[Cell Micrograph](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[Related Cell Culture Products](#)

Login Required ▶

[Product Information Sheet](#)

[BioProducts](#)

[Cell, microbial and molecular genomics products for the](#)

- [life sciences](#)

[BioServices](#)

[Bio-materials management; basic repository to complex partnership-level](#)

- [services](#)

Protocol: IMPORTANT - DO NOT ALLOW CULTURES TO BECOME CONFLUENT.

Cultures must not be allowed to become confluent as this will deplete the myoblastic population in the culture. Myotube formation is enhanced when the medium is supplemented with 10% horse serum instead of fetal bovine serum.

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin-0.53 mM EDTA solution to remove all traces of serum which contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessels.
Inoculate at a cell concentration between 1.5×10^5 and 1.0×10^6 viable cells/75 cm².
6. Incubate cultures at 37°C.

Subculturing:

[BioStandards](#)

[Biological Reference Material and Consensus Standards for the life science community](#)

Medium Renewal: Every two to three days

Preservation: **Freeze medium:** Complete growth medium supplemented with 5% (v/v) DMSO

Storage temperature: liquid nitrogen vapor phase

Related Products: Recommended medium (without the additional supplements or serum described under ATCC Medium):[ATCC 30-2002](#)
recommended serum:[ATCC 30-2020](#)

References:

- 22903: Yaffe D, Saxel O. Serial passaging and differentiation of myogenic cells isolated from dystrophic mouse muscle. *Nature* 270: 725-727, 1977. PubMed: [563524](#)
- 22953: Blau HM, et al. Plasticity of the differentiated state. *Science* 230: 758-766, 1985. PubMed: [2414846](#)
- 23427: Katagiri T, et al. Bone morphogenetic protein-2 converts the differentiation pathway of C2C12 myoblasts into the osteoblast lineage [published erratum appears in *J Cell Biol* 1995 Feb;128(4):following 713]. *J. Cell Biol.* 127: 1755-1766, 1994. PubMed: [7798324](#)
- 28236: Chow YH, et al. Improvement of hepatitis B virus DNA vaccines by plasmids coexpressing hepatitis B surface antigen and interleukin-2. *J. Virol.* 71: 169-178, 1997. PubMed: [8985336](#)
- 32828: Kessler PD, et al. Gene delivery to skeletal muscle results in sustained expression and systemic delivery of a therapeutic protein. *Proc. Natl. Acad. Sci. USA* 93: 14082-14087, 1996. PubMed: [8943064](#)
- 33069: Hsu DK, et al. Identification of a murine TEF-1-related gene expressed after mitogenic stimulation of quiescent fibroblasts and during myogenic differentiation. *J. Biol. Chem.* 271: 13786-13795, 1996. PubMed: [8662936](#)

[Return to Top](#)

Cell Biology

ATCC® Number: **CRL-1720™** Order this Item Price: **\$379.00**

Designations: F9
 Depositors: S Strickland
Biosafety Level: 1
 Shipped: frozen
 Medium & Serum: See Propagation
 Growth Properties: adherent
 Organism: *Mus musculus* (mouse)
 Morphology: epithelial

Related Links ▶
[NCBI Entrez Search](#)
[Make a Deposit](#)
[Frequently Asked Questions](#)
[Material Transfer Agreement](#)
[Technical Support](#)
[Related Cell Culture Products](#)

Source: **Organ:** testis
Strain: 129
Disease: embryonal carcinoma; testicular teratoma

Cellular Products: plasminogen activator; laminin; type IV collagen
 In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

BioProducts

[Cell, microbial and molecular genomics products for the life sciences](#)

Permits/Forms:

Applications: transfection host ([Roche FuGENE® Transfection Reagents](#))

Age: embryo

BioServices

[Bio-materials management; basic repository to complex partnership-level services](#)

Comments: F9 cells can be stimulated to differentiate into parietal endoderm in the presence of retinoic acid and dibutyryl cyclic AMP (cAMP). Differentiating cells synthesize plasminogen activator, laminin and type IV collagen. cAMP is active only on cells that have been treated with retinoic acid. The cells maintain three copies of the beta 1 integrin gene. Tested and found negative for ectromelia virus (mousepox).

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.
Temperature: 37.0°C

BioStandards

[Biological Reference Material and Consensus Standards for the life science community](#)

Protocol: NOTE: culture vessels must be coated with 0.1% gelatin prior to use. To do so, cover the surface of the vessel with 0.1% gelatin (Difco) in sterile distilled water for 2 hours at 4C, then wash three times with sterile distilled water. Treated flasks and dishes can be stored at room temperature.

Subculturing:

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin-0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new coated culture vessels.
6. Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:10 is recommended

Medium Renewal: Every 2 to 3 days

Related Products:

Recommended medium (without the additional supplements or serum described under ATCC Medium): [ATCC 30-2002](#)
recommended serum: [ATCC 30-2020](#)

References:

- 1160: Strickland S, et al. Hormonal induction of differentiation in teratocarcinoma stem cells: generation of parietal endoderm by retinoic acid and dibutyryl cAMP. *Cell* 21: 347-355, 1980. PubMed: [6250719](#)
- 1161: Strickland S, Mahdavi V. The induction of differentiation in teratocarcinoma stem cells by retinoic acid. *Cell* 15: 393-403, 1978. PubMed: [214238](#)
- 23426: Stephens LE, et al. Targeted deletion of beta 1 integrins in F9 embryonal carcinoma cells affects morphological differentiation but not tissue-specific gene expression. *J. Cell Biol.* 123: 1607-1620, 1993. PubMed: [7504677](#)
- 26151: Berstine EG, et al. Alkaline phosphatase activity in mouse teratoma. *Proc. Natl. Acad. Sci. USA* 70: 3899-3903, 1973. PubMed: [4521215](#)
- 32547: Jang SI, et al. Activator protein 1 activity is involved in the regulation of the cell type-specific expression from the proximal promoter of the human profilaggrin gene. *J. Biol. Chem.* 271: 24105-24114, 1996. PubMed: [8798649](#)

[Return to Top](#)

Cell Biology

ATCC® Number: **CRL-1658™** [Order this Item](#) Price: **\$279.00**

Designations: NIH/3T3

Biosafety Level: 1

Shipped: frozen

Medium & Serum: [See Propagation](#)

Growth Properties: adherent

Organism: *Mus musculus* (mouse)
fibroblast

Morphology: 

Source: **Organ:** embryo

Strain: NIH/Swiss

Cell Type: fibroblast fibroblast;

Permits/Forms: In addition to the [MTA](#) mentioned above, other [ATCC](#) and/or [regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

Virus Susceptibility: Murine leukemia virus

Age: embryo

Comments: The NIH/3T3 is highly sensitive to sarcoma virus focus formation and leukemia virus propagation and has proven to be very useful in DNA transfection studies [PubMed ID: 222457]. Tested and found negative for ectromelia virus (mousepox).

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium: bovine calf serum to a final concentration of 10%.

Atmosphere: air, 95%; carbon dioxide (CO₂), 5%

Temperature: 37.0°C

Growth Conditions: The serum used is important in culturing this line. Calf serum is recommended and not fetal bovine serum. The calf serum initially employed and found to be satisfactory was from the Colorado Serum Co. Denver.

Related Links ▶

[NCBI Entrez Search](#)

[Cell Micrograph](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[Related Cell Culture Products](#)

Login Required ▶

[Product Information Sheet](#)

BioProducts

[Cell, microbial and molecular genomics products for the life sciences](#)

- [life sciences](#)

BioServices

[Bio-materials management; basic repository to complex partnership-level services](#)

- [services](#)

[Return to Top](#)

Cell Biology

ATCC® Number: **CCL-92™** Order this Item Price: **\$279.00**

Designations: 3T3-Swiss albino

Depositors: H Green

Biosafety Level: 1

Shipped: frozen

Medium & Serum: See Propagation

Growth Properties: adherent

Organism: *Mus musculus* (mouse)
fibroblast

Morphology: 

Source: **Organ:** embryo
Cell Type: fibroblast

Cellular Products: Lysophosphatidylcholine (lyso-PC) induces AP-1 activity and c-jun N-terminal kinase activity (JNK1) by a protein kinase C-independent pathway [26623]

Permits/Forms: In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location.

Isolation: **Isolation date:** 1962

Cytogenetic Analysis: This is a hypertriploid mouse cell line. The modal chromosome number was 68 occurring in 30% of cells. The rate of cells with higher ploidies was 2.4%.

Age: embryo

Comments: The 3T3 cell line was established by G. Todaro and H. Green in 1962 from disaggregated Swiss mouse embryos. [5732]
The cells are contact inhibited.
A confluent monolayer yields 40000 cells/sq cm.
Tested and found negative for ectromelia virus (mousepox).
The cells should be grown in plastic flasks, they do not grow well on some types of glass surfaces.
A saturation density of approximately 50000 cells per sq cm can be reached.

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium: bovine calf serum to a final concentration of 10%.
Temperature: 37.0°C

Related Links ▶

[NCBI Entrez Search](#)

[Cell Micrograph](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[Related Cell Culture Products](#)

Login Required ▶

[Product Information Sheet](#)

BioProducts

[Cell, microbial and molecular genomics products for the life sciences](#)

BioServices

[Bio-materials management; basic repository to complex partnership-level services](#)

References:

- 5732: Todaro GJ, Green H. Quantitative studies of the growth of mouse embryo cells in culture and their development into established lines. *J. Cell Biol.* 17: 299-313, 1963. PubMed: [13985244](#)
- 21632: Bennicelli JL, et al. Mechanism for transcriptional gain of function resulting from chromosomal translocation in alveolar rhabdomyosarcoma. *Proc. Natl. Acad. Sci. USA* 93: 5455-5459, 1996. PubMed: [8643596](#)
- 26261: Vogt M, Dulbecco R. Studies on cells rendered neoplastic by polyoma virus: the problem of the presence of virus-related materials. *Virology* 16: 41-51, 1962. PubMed: [13926482](#)
- 26262: Todaro GJ, et al. Antigenic and cultural properties of cells doubly transformed by polyoma virus and SV40. *Virology* 27: 179-185, 1965. PubMed: [4284655](#)
- 26263: Todaro GJ, et al. Transformation of properties of an established cell line by SV40 and polyoma virus. *Proc. Natl. Acad. Sci. USA* 51: 66-73, 1964. PubMed: [14104605](#)
- 26623: Fang X, et al. Lysophosphatidylcholine stimulates activator protein 1 and the c-Jun N-terminal kinase activity. *J. Biol. Chem.* 272: 13683-13689, 1997. PubMed: [9153219](#)
- 32307: Chen ST, et al. Generation of packaging cell lines for pseudotyped retroviral vectors of the G protein of vesicular stomatitis virus by using a modified tetracycline inducible system. *Proc. Natl. Acad. Sci. USA* 93: 10057-10062, 1996. PubMed: [8816750](#)
- 32500: Campbell M, et al. The simian foamy virus type 1 transcriptional transactivator (Tas) binds and activates an enhancer element in the gag gene. *J. Virol.* 70: 6847-6855, 1996. PubMed: [8794326](#)
- 33069: Hsu DK, et al. Identification of a murine TEF-1-related gene expressed after mitogenic stimulation of quiescent fibroblasts and during myogenic differentiation. *J. Biol. Chem.* 271: 13786-13795, 1996. PubMed: [8662936](#)

[Return to Top](#)

Cell Biology

ATCC® Number: **CRL-1651™** Order this Item Price: **\$279.00**

Designations: COS-7
 Depositors: Y Gluzman
Biosafety Level: 2 [Cells Contain SV-40 viral DNA sequences]
 Shipped: frozen
 Medium & Serum: See Propagation
 Growth Properties: adherent
 Organism: *Cercopithecus aethiops*
 fibroblast

Morphology: 

Source: **Organ:** kidney
Cell Type: SV40 transformed

Cellular Products: T antigen
 In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

Applications: transfection host ([Nucleofection technology from Lonza Roche FuGENE® Transfection Reagents](#))

Comments: This is an African green monkey kidney fibroblast-like cell line suitable for transfection by vectors requiring expression of SV40 T antigen. This line contains T antigen, retains complete permissiveness for lytic growth of SV40, supports the replication of ts A209 virus at 40C, and supports the replication of pure populations of SV40 mutants with deletions in the early region. The line was derived from the CV-1 cell line (ATCC® CCL-70?) by transformation with an origin defective mutant of SV40 which codes for wild type T antigen.

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.
Atmosphere: air, 95%; carbon dioxide (CO2), 5%
Temperature: 37.0°C

- Related Links ▶**
[NCBI Entrez Search](#)
[Cell Micrograph](#)
[Make a Deposit](#)
[Frequently Asked Questions](#)
[Material Transfer Agreement](#)
[Technical Support](#)
[Related Cell Culture Products](#)

- Login Required ▶**
[Product Information Sheet](#)

BioProducts

- [Cell, microbial and molecular genomics products for the](#)
 • [life sciences](#)

BioServices

- [Bio-materials management; basic repository to complex partnership-level](#)
 • [services](#)

Protocol:

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin-0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessels.
6. Incubate cultures at 37°C.

Subculturing:

BioStandards

- [Biological Reference Material and Consensus Standards for the life science community](#)

Subcultivation Ratio: A subcultivation ratio of 1:4 to 1:8 is recommended

Medium Renewal: 2 to 3 times per week

Preservation:

Freeze medium: Complete growth medium supplemented with 5% (v/v) DMSO

Storage temperature: liquid nitrogen vapor phase

Recommended medium (without the additional supplements or serum described under ATCC Medium): [ATCC 30-2002](#)

recommended serum: [ATCC 30-2020](#)

Related Products:

parental cell line: [ATCC CCL-70](#)

0.25% (w/v) Trypsin - 0.53 mM EDTA in Hank' BSS (w/o Ca⁺⁺, Mg⁺⁺): [ATCC 30-2101](#)

Cell culture tested DMSO: [ATCC 4-X](#)

- 1822: Gluzman Y. SV40-transformed simian cells support the replication of early SV40 mutants. *Cell* 23: 175-182, 1981. PubMed: [6260373](#)
- 32447: Fernandez LM, Puett D. Lys583 in the third extracellular loop of the lutropin/choriogonadotropin receptor is critical for signaling. *J. Biol. Chem.* 271: 925-930, 1996. PubMed: [8557706](#)
- 32459: Maestrini E, et al. A family of transmembrane proteins with homology to the MET-hepatocyte growth factor receptor. *Proc. Natl. Acad. Sci. USA* 93: 674-678, 1996. PubMed: [8570614](#)
- 32500: Campbell M, et al. The simian foamy virus type 1 transcriptional transactivator (Tas) binds and activates an enhancer element in the gag gene. *J. Virol.* 70: 6847-6855, 1996. PubMed: [8794326](#)
- 32502: Gonzalez Armas JC, et al. DNA immunization confers protection against murine cytomegalovirus infection. *J. Virol.* 70: 7921-7928, 1996. PubMed: [8892915](#)
- 32547: Jang SI, et al. Activator protein 1 activity is involved in the regulation of the cell type-specific expression from the proximal promoter of the human profilaggrin gene. *J. Biol. Chem.* 271: 24105-24114, 1996. PubMed: [8798649](#)
- 32566: Dittrich E, et al. A di-leucine motif and an upstream serine in the interleukin-6 (IL-6) signal transducer gp130 mediate ligand-induced endocytosis and down-regulation of the IL-6 receptor. *J. Biol. Chem.* 271: 5487-5494, 1996. PubMed: [8621406](#)
- 32568: Lee JH, et al. The proximal promoter of the human transglutaminase 3 gene. *J. Biol. Chem.* 271: 4561-4568, 1996. PubMed: [8626812](#)
- 32720: Chen Y, et al. Demonstration of binding of dengue virus envelope protein to target cells. *J. Virol.* 70: 8765-8772, 1996. PubMed: [8971005](#)
- 32728: Russell DW, Miller AD. Foamy virus vectors. *J. Virol.* 70: 217-222, 1996. PubMed: [8523528](#)
- 32861: Wright DA, et al. Association of human fas (CD95) with a ubiquitin-conjugating enzyme (UBC-FAP). *J. Biol. Chem.* 49: 31037-31043, 1996. PubMed: [8940097](#)
- 32893: Zhang J, et al. Dynamin and beta-arrestin reveal distinct mechanisms for G protein-coupled receptor internalization. *J. Biol. Chem.* 271: 18302-18305, 1996. PubMed: [8702465](#)
- 33011: Ozcelebi F, et al. Phosphorylation of cholecystokinin receptors expressed on chinese hamster ovary cells. *J. Biol. Chem.* 271: 3750-3755, 1996. PubMed: [8631990](#)
- 33013: Gibson S, et al. Functional LCK is required for optimal CD28-mediated activation of the TEC family tyrosine kinase EMT/ITK. *J. Biol. Chem.* 271: 7079-7083, 1996. PubMed: [8636141](#)
- 33016: Shaul PW, et al. Acylation targets endothelial nitric-oxide synthase to plasmalemmal caveolae. *J. Biol. Chem.* 271: 6518-6522, 1996. PubMed: [8626455](#)
- 33032: Ladner RD, et al. Identification of a consensus cyclin-

References:

[Return to Top](#)

----- Original Message -----

Subject:[Fwd: Fwd: updated Biological Agent Registry Form (Litchfield)]

Date:Mon, 21 Mar 2011 16:25:25 -0400

From:Laszlo Gyenis <lgyenis@uwo.ca>

To:Jennifer Stanley <jstanle2@uwo.ca>

CC:David Litchfield <litchfi@uwo.ca>

Hi Jennifer,

I am sending you the plasmid maps of pBI, pRc/CMV, pTRE, pEGFP and pcDNA3.1 vectors.
We are not using oncogenes.

Thank you, Laszlo



Info on Plasmid(s)



Find this plasmid at: www.addgene.org
Enter "25536" in the search box

Plasmid 25536: NME3

Gene/insert name: NME3
 Alternative names: PDB: 1ZS6
 Nucleoside-diphosphate Kinase 3
 c371H6.2, DR-nm23, KIAA0516
 Insert size (bp): 507
 Gene/insert aliases: NME3, NDPKC, NDPK-C, NM23H3, DR-nm23, NM23-
 H3, KIAA0516, c371H6.2
 Species of gene(s): H. sapiens (human)
 Fusion proteins or tags: His
 Terminal: N terminal on insert
 Vector backbone: pET28a-LIC
[\(Search Vector Database\)](#)
 Backbone manufacturer: SGC
 Type of vector: Bacterial expression
 Backbone size (bp): 7328
 5' Sequencing primer: T7 [\(List of Sequencing Primers\)](#)
 3' Sequencing primer: T7-term
 Bacteria resistance: Kanamycin
 High or low copy: High Copy
 Grow in standard E. coli @ 37C: Yes
 Sequence: Visit www.addgene.org/25536
 Plasmid Provided In: DH5a
 Principal Investigator: Cheryl Arrowsmith

Comments: PDB: 1ZS6 <http://www.thesgconline.org/SGC-WebPages/StructureDescription/1ZS6.php>

Please acknowledge the principal investigator if you use this plasmid in a publication.

Also, please include the text "Addgene plasmid 25536" in your Materials and Methods section. This information allows Addgene to create a link from the plasmid page to your publication.


Please check www.addgene.org/25536 for updated plasmid information and related links.

Page 1 of 2 - Date: 03/08/2011

Information on this datasheet is provided pursuant to Addgene's Terms of Use at www.addgene.org.



Find this plasmid at: www.addgene.org
Enter "25536" in the search box



Please check www.addgene.org/25536 for updated plasmid information and related links.

Page 2 of 2 - Date: 03/08/2011

Information on this datasheet is provided pursuant to Addgene's Terms of Use at www.addgene.org.



Find this plasmid at: www.addgene.org
Enter "25536" in the search box

Plasmid 25536: NME3

Gene insert name	NME3
Alternative names	PDB: 1ZS6 Nucleoside-diphosphate Kinase 3 c371H6.2, DR-nm23, KIAA0516
Insert size (bp)	507
Gene name aliases	NME3, NDPKC, NDPK-C, NM23H3, DR-nm23, NM23-H3, KIAA0516, c371H6.2
Species of gene(s)	H. sapiens (human)
Fusion proteins or tags	His
Terminal	N terminal on insert
Vector backbone	pET28a-LIC (Search Vector Database)
Backbone manufacturer	SGC
Type of vector	Bacterial expression
Backbone size (bp)	7328
5' Sequencing primer	T7 (List of Sequencing Primers)
3' Sequencing primer	T7-term
Bacteria resistance	Kanamycin
High or low copy	High Copy
Growth standard E. coli @ 37C	Yes
Sequence	Visit www.addgene.org/25536
Plasmid Provided in	DH5a
Principal Investigator	Cheryl Arrowsmith

Comments: PDB: 1ZS6 <http://www.thesgconline.org/SGC-WebPages/StructureDescription/1ZS6.php>

Please acknowledge the principal investigator if you use this plasmid in a publication.

Also, please include the text "Addgene plasmid 25536" in your Materials and Methods section. This information allows Addgene to create a link from the plasmid page to your publication.


Please check www.addgene.org/25536 for updated plasmid information and related links.

Page 1 of 2 - Date: 03/21/2011

Information on this datasheet is provided pursuant to Addgene's Terms of Use at www.addgene.org.



Find this plasmid at: www.addgene.org
Enter "25536" in the search box



Please check www.addgene.org/25536 for updated plasmid information and related links.

Page 2 of 2 - Date: 03/21/2011

Information on this datasheet is provided pursuant to Addgene's Terms of Use at www.addgene.org.

Patent Depository

ATCC® Number: **PTA-4822™** Order this Item Price: **\$200.00**

Designation / Description: Plasmid, 819PH59 in E.coli XL1-Blue

U.S. Patent Number: [6,855,365](#)
[7,432,091](#)

[Biosafety Level:](#) 1

Shipped: frozen

Permits/Forms: In addition to the [MTA](#) mentioned above, other [ATCC](#) and/or [regulatory permits](#) may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please [click here](#) for information regarding the specific requirements for shipment to your location.

This material is cited in a U.S. and/or other Patent or Patent Application, and may not be used to infringe on the patent claims. ATCC is required to inform the Patent Depositor of the party to which the material was furnished.

Related Links ▶

[NCBI Entrez Search](#)

[Make a Deposit](#)

[Frequently Asked Questions](#)

[Material Transfer Agreement](#)

[Technical Support](#)

[BioProducts](#)

[Cell, microbial and molecular genomics products for the life sciences](#)

- [sciences](#)

[BioServices](#)

[Bio-materials management; basic repository to complex partnership-level](#)

- [services](#)

[BioStandards](#)

[Biological Reference Material and Consensus Standards for the life science](#)

- [community](#)

[Return to Top](#)

1. PRODUCT AND COMPANY IDENTIFICATION

Product name : Okadaic acid, from *Prorocentrum concavum*

Product Number : O9381

Brand : Sigma

Product Use : For laboratory research purposes.

Supplier : Sigma-Aldrich Canada, Ltd
2149 Winston Park Drive
OAKVILLE ON L6H 6J8
CANADA

Manufacturer : Sigma-Aldrich Corporation
3050 Spruce St.
St. Louis, Missouri 63103
USA

Telephone : +19058299500

Fax : +19058299292

Emergency Phone # (For both supplier and manufacturer) : 1-800-424-9300

Preparation Information : Sigma-Aldrich Corporation
Product Safety - Americas Region
1-800-521-8956

2. HAZARDS IDENTIFICATION

Emergency Overview

Target Organs

Smooth muscle.

WHMIS Classification

D1A	Very Toxic Material Causing Immediate and	Highly Toxic
D2B	Serious Toxic Effects	Moderate skin irritant

GHS Classification

Acute toxicity, Oral (Category 3)
Acute toxicity, Inhalation (Category 3)
Acute toxicity, Dermal (Category 3)
Skin irritation (Category 2)

GHS Label elements, including precautionary statements

Pictogram



Signal word : Danger

Hazard statement(s)

H301 + H311	Toxic if swallowed or in contact with skin.
H315	Causes skin irritation.
H331	Toxic if inhaled.

Precautionary statement(s)

P261	Avoid breathing dust/ fume/ gas/ mist/ vapours/ spray.
P280	Wear protective gloves/ protective clothing.
P301 + P310	IF SWALLOWED: Immediately call a POISON CENTER or doctor/ physician.
P311	Call a POISON CENTER or doctor/ physician.

HMIS Classification

Health hazard: 2

Toxin Info

Flammability: 0
Physical hazards: 0

Potential Health Effects

Inhalation Toxic if inhaled. Causes respiratory tract irritation.
Skin Toxic if absorbed through skin. Causes skin irritation.
Eyes Causes eye irritation.
Ingestion Toxic if swallowed.

3. COMPOSITION/INFORMATION ON INGREDIENTS

Synonyms : OA
Formula : C₄₄H₆₈O₁₃ C₄₄H₆₈O₁₃
Molecular Weight : 805.00 g/mol

CAS-No.	EC-No.	Index-No.	Concentration
Okadaic acid			
78111-17-8	-	-	-

4. FIRST AID MEASURES

General advice

Consult a physician. Show this safety data sheet to the doctor in attendance. Move out of dangerous area.

If inhaled

If breathed in, move person into fresh air. If not breathing, give artificial respiration. Consult a physician.

In case of skin contact

Wash off with soap and plenty of water. Take victim immediately to hospital. Consult a physician.

In case of eye contact

Flush eyes with water as a precaution.

If swallowed

Never give anything by mouth to an unconscious person. Rinse mouth with water. Consult a physician.

5. FIRE-FIGHTING MEASURES

Conditions of flammability

Not flammable or combustible.

Suitable extinguishing media

Use water spray, alcohol-resistant foam, dry chemical or carbon dioxide.

Special protective equipment for fire-fighters

Wear self contained breathing apparatus for fire fighting if necessary.

Hazardous combustion products

Hazardous decomposition products formed under fire conditions. - Carbon oxides

Explosion data - sensitivity to mechanical impact

no data available

Explosion data - sensitivity to static discharge

no data available

6. ACCIDENTAL RELEASE MEASURES

Personal precautions

Wear respiratory protection. Avoid dust formation. Avoid breathing vapors, mist or gas. Ensure adequate ventilation. Evacuate personnel to safe areas. Avoid breathing dust.

Environmental precautions

Prevent further leakage or spillage if safe to do so. Do not let product enter drains.

Methods and materials for containment and cleaning up

Pick up and arrange disposal without creating dust. Sweep up and shovel. Keep in suitable, closed containers for disposal.

7. HANDLING AND STORAGE

Precautions for safe handling

Avoid contact with skin and eyes. Avoid formation of dust and aerosols. Provide appropriate exhaust ventilation at places where dust is formed. Normal measures for preventive fire protection.

Conditions for safe storage

Keep container tightly closed in a dry and well-ventilated place.

Recommended storage temperature: -20 °C

Keep in a dry place.

8. EXPOSURE CONTROLS/PERSONAL PROTECTION

Contains no substances with occupational exposure limit values.

Personal protective equipment

Respiratory protection

Where risk assessment shows air-purifying respirators are appropriate use a full-face particle respirator type N99 (US) or type P2 (EN 143) respirator cartridges as a backup to engineering controls. If the respirator is the sole means of protection, use a full-face supplied air respirator. Use respirators and components tested and approved under appropriate government standards such as NIOSH (US) or CEN (EU).

Hand protection

Handle with gloves. Gloves must be inspected prior to use. Use proper glove removal technique (without touching glove's outer surface) to avoid skin contact with this product. Dispose of contaminated gloves after use in accordance with applicable laws and good laboratory practices. Wash and dry hands.

Eye protection

Face shield and safety glasses Use equipment for eye protection tested and approved under appropriate government standards such as NIOSH (US) or EN 166(EU).

Skin and body protection

Complete suit protecting against chemicals, The type of protective equipment must be selected according to the concentration and amount of the dangerous substance at the specific workplace.

Hygiene measures

Avoid contact with skin, eyes and clothing. Wash hands before breaks and immediately after handling the product.

Specific engineering controls

Use mechanical exhaust or laboratory fumehood to avoid exposure.

9. PHYSICAL AND CHEMICAL PROPERTIES

Appearance

Form	powder
Colour	white

Safety data

pH	no data available
Melting/freezing point	no data available
Boiling point	no data available
Flash point	no data available
Ignition temperature	no data available
Autoignition	no data available

temperature	
Lower explosion limit	no data available
Upper explosion limit	no data available
Vapour pressure	no data available
Density	no data available
Water solubility	no data available
Partition coefficient: n-octanol/water	no data available
Relative vapour density	no data available
Odour	no data available
Odour Threshold	no data available
Evaporation rate	no data available

10. STABILITY AND REACTIVITY

Chemical stability

Stable under recommended storage conditions.

Possibility of hazardous reactions

no data available

Conditions to avoid

no data available

Materials to avoid

Strong oxidizing agents

Hazardous decomposition products

Hazardous decomposition products formed under fire conditions. - Carbon oxides

Other decomposition products - no data available

11. TOXICOLOGICAL INFORMATION

Acute toxicity

Oral LD50

TDL_o Oral - mouse - 0.15 mg/kg

Inhalation LC50

Dermal LD50

Other information on acute toxicity

no data available

Skin corrosion/irritation

no data available

Serious eye damage/eye irritation

no data available

Respiratory or skin sensitization

no data available

Germ cell mutagenicity

Genotoxicity in vitro - Human - lymphocyte

Sister chromatid exchange

Genotoxicity in vitro - Hamster - Lungs

Mutation in mammalian somatic cells.

Carcinogenicity

Carcinogenicity - mouse - Skin

Tumorigenic: Equivocal tumorigenic agent by RTECS criteria. Skin and Appendages: Other: Tumors.

IARC: No component of this product present at levels greater than or equal to 0.1% is identified as probable, possible or confirmed human carcinogen by IARC.

ACGIH: No component of this product present at levels greater than or equal to 0.1% is identified as a carcinogen or potential carcinogen by ACGIH.

Reproductive toxicity

no data available

Teratogenicity

no data available

Specific target organ toxicity - single exposure (Globally Harmonized System)

no data available

Specific target organ toxicity - repeated exposure (Globally Harmonized System)

no data available

Aspiration hazard

no data available

Potential health effects

Inhalation	Toxic if inhaled. Causes respiratory tract irritation.
Ingestion	Toxic if swallowed.
Skin	Toxic if absorbed through skin. Causes skin irritation.
Eyes	Causes eye irritation.

Signs and Symptoms of Exposure

Tumor promoter., Gastrointestinal disturbance

Synergistic effects

no data available

Additional Information

RTECS: AA8227800

12. ECOLOGICAL INFORMATION

Toxicity

no data available

Persistence and degradability

no data available

Bioaccumulative potential

no data available

Mobility in soil

no data available

PBT and vPvB assessment

no data available

Other adverse effects

no data available

13. DISPOSAL CONSIDERATIONS

Product

Offer surplus and non-recyclable solutions to a licensed disposal company. Contact a licensed professional waste disposal service to dispose of this material. Dissolve or mix the material with a combustible solvent and burn in a chemical incinerator equipped with an afterburner and scrubber.

Contaminated packaging

Dispose of as unused product.

14. TRANSPORT INFORMATION**DOT (US)**

UN-Number: 3462 Class: 6.1 Packing group: I
Proper shipping name: Toxins, extracted from living sources, solid, n.o.s. (Okadaic acid)
Marine pollutant: No
Poison Inhalation Hazard: No

IMDG

UN-Number: 3462 Class: 6.1 Packing group: I EMS-No: F-A, S-A
Proper shipping name: TOXINS, EXTRACTED FROM LIVING SOURCES, SOLID, N.O.S. (Okadaic acid)
Marine pollutant: No

IATA

UN-Number: 3462 Class: 6.1 Packing group: I
Proper shipping name: Toxins, extracted from living sources, solid, n.o.s. (Okadaic acid)

15. REGULATORY INFORMATION**DSL Status**

This product contains the following components that are not on the Canadian DSL nor NDSL lists.

Okadaic acid

CAS-No.
78111-17-8

WHMIS Classification

D1A	Very Toxic Material Causing Immediate and	Highly Toxic
D2B	Serious Toxic Effects	Moderate skin irritant

This product has been classified in accordance with the hazard criteria of the Controlled Products Regulations and the MSDS contains all the information required by the Controlled Products Regulations.

16. OTHER INFORMATION**Further information**

Copyright 2011 Sigma-Aldrich Co. License granted to make unlimited paper copies for internal use only.
The above information is believed to be correct but does not purport to be all inclusive and shall be used only as a guide. The information in this document is based on the present state of our knowledge and is applicable to the product with regard to appropriate safety precautions. It does not represent any guarantee of the properties of the product. Sigma-Aldrich Co., shall not be held liable for any damage resulting from handling or from contact with the above product. See reverse side of invoice or packing slip for additional terms and conditions of sale.



TOXIN USE RISK ASSESSMENT

Name of Toxin:	Okadaic Acid
Proposed Use Dose:	5 µg
Proposed Storage Dose:	100 µg
LD₅₀ (species):	150 µg

Calculation:	
150 µg/kg	x 50 kg/person
Dose per person based on LD ₅₀ in µg = 7500	
LD₅₀ per person with safety factor of 10 based on LD₅₀ in µg =	750

Comments/Recommendations: