

THE UNIVERSITY OF WESTERN ONTARIO
BIOHAZARDOUS AGENTS REGISTRY FORM
Approved Biohazards Subcommittee: November 21, 2008
Biosafety Website: www.uwo.ca/humanresources/biosafety/

This form must be completed by each Principal Investigator holding a grant administered by the University of Western Ontario or in charge of a laboratory/facility where the use of Level 1, 2 or 3 biohazardous agents is described in the laboratory or animal work proposed. The form must also be completed if any work is proposed involving animals carrying zoonotic agents infectious to humans or involving plants, fungi, or insects that require Health Canada (HC) or Canadian Food Inspection Agency (CFIA) permits.

This form must also be updated at least every 3 years or when there are changes to the biohazards being used.

Containment Levels will be established in accordance with Laboratory Biosafety Guidelines, 3rd edition, 2004, Health Canada (HC) or Containment Standards for Veterinary Facilities, 1st edition 1996, Canadian Food Inspection Agency (CFIA).

Completed forms are to be returned to Occupational Health and Safety, (OHS), (Support Services Building, Room 4190) for distribution to the Biohazard Subcommittee. For questions regarding this form, please contact the Biosafety Officer at extension 81135. If there are changes to the information on this form (excluding grant title and funding agencies), modifications must be submitted to Occupational Health and Safety. See website: www.uwo.ca/humanresources/biosafety/

PRINCIPAL INVESTIGATOR Dr. Ting-Ying Lee
SIGNATURE Ting-Ying Lee
DEPARTMENT Imaging
ADDRESS The Lawson Health Research Institute
PHONE NUMBER (x) 24131
EMAIL tleee@lawsonimaging.ca

Location of experimental work to be carried out: Building(s) LHR1 Room(s) F4-127a

*For work being performed at Institutions affiliated with the University of Western Ontario, the Safety Officer for the Institution where experiments will take place must sign the form prior to its being sent to Occupational Health and Safety (See Section 12.0, Approvals). For research being done at Lawson Health Research Institute, London Regional Cancer Program, Child and Parent Research Institute, or Robarts Research Institute, a University Biosafety Committee member can also sign as the Safety Officer for the Institution.

FUNDING AGENCY/AGENCIES: _____
GRANT TITLE(S): _____

PLEASE ATTACH A BRIEF DESCRIPTION OF YOUR WORK THAT EXPLAINS THE BIOHAZARDS USED AND HOW THEY WILL BE USED. PROJECTS SUBMITTED WITHOUT A SUMMARY WILL NOT BE REVIEWED.

Names of all personnel working under Principal Investigators supervision in this location:
Lisa Hoffman

1.0 Microorganisms

1.1 Does your work involve the use of microorganisms or biological agents of plant or animal origin (including but not limited to viruses, prions, parasites, bacteria)? YES NO
 If no, please proceed to Section 2.0

Do you use microorganisms that require a permit from the CFIA? YES NO
 If YES, please give the name of the species. lentiviruses

What is the origin of the microorganism(s)? (see attached info from Invitrogen)

Please describe the risk (if any) of escape and how this will be mitigated:
(see attached). Rooms in which lentiviruses are Level 2+ (ie, Hepa-certified BSCs, portable autoclave, gloves, labcoats, bleach, etc.

Please attach the CFIA permit.

Please describe any CFIA permit conditions:

1.2 Please complete the table below:

| Name of Biological agent(s)* | Is it known to be a human pathogen? YES/NO | Is it known to be an animal pathogen? YES/NO | Is it known to be a zoonotic agent? YES/NO | Maximum quantity to be cultured at one time? (in Litres) | Source/Supplier | Health Canada or CFIA Containment Level |
|------------------------------|---|---|---|---|-----------------|---|
| | <input type="radio"/> Yes <input type="radio"/> No | <input type="radio"/> Yes <input type="radio"/> No | <input type="radio"/> Yes <input type="radio"/> No | | | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 |
| | <input type="radio"/> Yes <input type="radio"/> No | <input type="radio"/> Yes <input type="radio"/> No | <input type="radio"/> Yes <input type="radio"/> No | | | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 |
| | <input type="radio"/> Yes <input type="radio"/> No | <input type="radio"/> Yes <input type="radio"/> No | <input type="radio"/> Yes <input type="radio"/> No | | | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 |
| | <input type="radio"/> Yes <input type="radio"/> No | <input type="radio"/> Yes <input type="radio"/> No | <input type="radio"/> Yes <input type="radio"/> No | | | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 |

*Please attach a Material Safety Data Sheet or equivalent from the supplier.

2.0 Cell Culture

2.1 Does your work involve the use of cell cultures? YES NO
 If no, please proceed to Section 3.0

2.2 Please indicate the type of primary cells (i.e. derived from fresh tissue) that will be grown in culture in the table below

| Cell Type | Is this cell type used in your work? | Source of Primary Cell Culture Tissue | AUS Protocol Number |
|-------------------|---|---|---------------------|
| Human | <input checked="" type="radio"/> Yes <input type="radio"/> No | | Not applicable |
| Rodent | <input checked="" type="radio"/> Yes <input type="radio"/> No | 1) muscle satellite cells (SCs) harvested from transgenic mice (not a biohazard, level 1, housed at | |
| Non-human primate | <input type="radio"/> Yes <input type="radio"/> No | | |
| Other (specify) | <input type="radio"/> Yes <input type="radio"/> No | 2) LHe1 primary myoblasts (obtained from Dr. Michael Rudnicki, see attached reference) | |

4) HEK (also called 293) received from Dr. Greg DeLaban; available from Invitrogen
 - human kidney (see attached notes)
 5) 293T (ATCC/human kidney) (see attached notes)

Level 2

2.3 Please indicate the type of established cells that will be grown in culture in the table below.

| Cell Type | Is this cell type used in your work? | Specific cell line(s)* | Supplier / Source |
|-------------------|---|--------------------------------------|---|
| Human | <input checked="" type="radio"/> Yes <input type="radio"/> No | 1) LoVo 2) NCI - H1299 3) PC-3 | ATCC (colorectal adenocarcinoma) ATCC / lung carcinoma ATCC / prostate adenocarcinoma |
| Rodent | <input checked="" type="radio"/> Yes <input type="radio"/> No | 1) C2C12 (mouse) 2) C6 (rat) | ATCC (muscle carcinoma) ATCC / glioma |
| Non-human primate | <input type="radio"/> Yes <input type="radio"/> No | | |
| Other (specify) | <input type="radio"/> Yes <input type="radio"/> No | | |

Level 1
Level 1
Level 1
Level 1
Level 1

*Please attach a Material Safety Data Sheet or equivalent from the supplier. (For more information, see www.atcc.org)

2.4 For above named cell types(s) indicate HC or CFIA containment level required 1 2 3

3.0 Use of Human Source Materials

3.1 Does your work involve the use of human source materials? YES NO
 If no, please proceed to Section 4.0

HEK/293T

3.2 Indicate in the table below the Human Source Material to be used.

| Human Source Material | Source/Supplier /Company Name | Is Human Source Material Known to Be Infected With An Infectious Agent? YES/NO | Name of Infectious Agent (If applicable) | HC or CFIA Containment Level (Select one) |
|--|-------------------------------|--|--|---|
| Human Blood (whole) or other Body Fluid | | <input type="radio"/> Yes <input type="radio"/> No | | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 |
| Human Blood (fraction) or other Body Fluid | | <input type="radio"/> Yes <input type="radio"/> No | | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 |
| Human Organs or Tissues (unpreserved) | | <input type="radio"/> Yes <input type="radio"/> No | | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 |
| Human Organs or Tissues (preserved) | | <input type="radio"/> Yes <input type="radio"/> No | | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 |

4.0 Genetically Modified Organisms and Cell lines

4.1 Will genetic modifications be made to the microorganisms, biological agents, or cells described in Sections 1.0 and 2.0? YES NO If no, please proceed to Section 5.0

4.2 Will genetic modification(s) involving plasmids be done? YES, complete table below NO

| Bacteria Used for Cloning * | Plasmid(s) * | Source of Plasmid | Gene Transfected | Describe the change that results |
|-----------------------------|--|-------------------|---|----------------------------------|
| Stb13 competent E. coli | ViraPower Promoteres Lentiviral vector | Invitrogen | Myogenin Promoter Tribusian Reporter | Expression of reporter genes |

* Please attach a Material Data Sheet or equivalent if available. (Firefly Luciferase/ monomeric red fluorescent protein/ truncated s39-thymidine Kinase) for optical & PET imaging

→ Lipofectamine cationic lipid-based transfection reagent to be used (Invitrogen)

* DESCRIPTION MUST BE ATTACHED TO THIS FORM OR PROJECT WILL NOT BE REVIEWED*

4.3 Will genetic modification(s) involving viral vectors be done? YES, complete table below NO

| Virus Used for Transduction * | Vector(s) * | Source of Vector | Gene Transfected | Describe the change that results |
|-------------------------------|-------------|------------------|------------------|----------------------------------|
| Lentivirus | See 4.2 | See 4.2 | See 4.2 | See 4.2 |

* Please attach a Material Safety Data Sheet or equivalent.

4.4 Will genetic sequences from the following be involved?

- ◆ HIV YES, please specify see attached sheet NO
- ◆ HTLV 1 or 2 or genes from any Level 1 or Level 2 pathogens YES, specify _____ NO
- ◆ SV 40 Large T antigen YES NO
- ◆ E1A oncogene YES NO
- ◆ Known oncogenes YES, please specify _____ NO
- ◆ Other human or animal pathogen and or their toxins YES, please specify _____ NO

4.5 Will virus be replication defective? YES NO

4.6 Will virus be infectious to humans or animals? YES NO

4.7 Will this be expected to increase the containment level required? YES NO

5.0 Human Gene Therapy Trials

5.1 Will human clinical trials be conducted using the viral vector in 4.0? YES NO
 If no, please proceed to Section 6.0 If YES attach a full description of the make-up of the virus.

5.2 Will virus be able to replicate in the host? YES NO

5.3 How will the virus be administered? _____

5.4 Please give the Health Care Facility where the clinical trial will be conducted: _____

5.5 Has human ethics approval been obtained? YES, number: _____ NO PENDING

6.0 Animal Experiments

6.1 Will live animals be used? YES NO If no, please proceed to section 7.0

6.2 Name of animal species to be used mdx mouse; mdx ltrn7 mouse

6.3 AUS protocol # 2008-067

6.4 Will any of the agents listed be used in live animals YES, specify: _____ NO

10.0 Plants Requiring CFIA Permits

10.1 Do you use plants that require a permit from the CFIA? YES NO
If no, please proceed to Section 11.0

10.2 If YES, please give the name of the species. _____

10.3 What is the origin of the plant? _____

10.4 What is the form of the plant (seed, seedling, plant, tree...)? _____

10.5 What is your intention? Grow and maintain a crop "One-time" use

10.6 Do you do any modifications to the plant? YES NO
If yes, please describe: _____

10.7 Please describe the risk (if any) of loss of the material from the lab and how this will be mitigated:

10.8 Is the CFIA permit attached? YES NO

10.9 Please describe any CFIA permit conditions:

11.0 Import Requirements

11.1 Will any of the above agents be imported? YES, please give country of origin USA
If no, please proceed to Section 10.0 NO

11.2 Has an Import Permit been obtained from HC for human pathogens? YES NO

11.3 Has an import permit been obtained from CFIA for animal or plant pathogens? YES NO

11.4 Has the import permit been sent to OHS? YES, please provide permit # P-13043 NO
A-2007-00178-4

12.0 Training Requirements for Personnel Named on Form

All personnel named on the above form who will be using any of the above named agents are required to attend the following training courses given by OHS:

- ◆ Biosafety
- ◆ Laboratory and Environmental/Waste Management Safety
- ◆ WHMIS (Western or equivalent)
- ◆ Employee Health and Safety Orientation

As the Principal Investigator, I have ensured that all of the personnel named on the form who will be using any of the biohazardous agents in Sections 1.0 to 9.0 have been trained.

SIGNATURE Feng Zhao Lee

13.0 Containment Levels

11.1 For the work described in sections 1.0 to 9.0, please indicate the highest HC or CFIA Containment Level required. O 1 2 O 3

13.2 Has the facility been certified by OHS for this level of containment?
 YES, permit # if on-campus B10-L1H1-028
 NO
 NOT REQUIRED

14.0 Procedures to be Followed

14.1 As the Principal Investigator, I will ensure that this project will follow the Western Biosafety Guidelines and Procedures Manual for Containment Level 1 & 2 Laboratories (and the Level 3 Facilities Manual for Level 3 projects). I will ensure that UWO faculty, staff and students working in my laboratory have an up-to-date Hazard Communication Form, found at <http://www.wph.uwo.ca/>

SIGNATURE Tony Lee Date: May 5/2009

15.0 Approvals

UWO Biohazard Subcommittee: SIGNATURE: _____
Date: _____

Safety Officer for Institution where experiments will take place: SIGNATURE: [Signature]
Date: May 5/2009

Safety Officer for University of Western Ontario (if different from above): SIGNATURE: _____
Date: _____

Approval Number: _____ Expiry Date (3 years from Approval): _____

Special Conditions of Approval:

----- Original Message -----

Subject:Re: Dr. Lee Biohazardous Agents Registry Form

Date:Thu, 14 May 2009 13:39:39 -0400

From:Jeff Tucker <Jeff.Tucker@sjhc.london.on.ca>

To:Jennifer Stanley <jstanle2@uwo.ca>

References:<4A0B3427.3000900@uwo.ca>

Hi Jennifer:

He is an imaging scientist and I believe that the documentation submitted was for all of the agents his lab group would be using, partly as requested by yourself. All of the agents work is done by Savita's group since she has the culture room and expertise working with these agents.

I think part of the problem is that the form only has three choices 1,2 and 3. The level 2+ work would be related to the lentiviral kits and I believe this submission is just for culturing and not specifically for administration to animals. The administration part, as I understand from Jennifer Hadway is still being worked out in regard to the HEPA containment device.

Savita has a level 2+ approved lab with portable autoclave.

Jeff

----- Original Message -----

Subject:Re: Biohazardous Agents Registry Form: Dr. T. Lee

Date:Tue, 19 May 2009 13:31:33 -0400 (EDT)

From:Lisa Hoffman <lhoffman@lawsonimaging.ca>

To:Jennifer Stanley <jstanle2@uwo.ca>

References:<4A0DDC7A.8040009@uwo.ca>

Hi Jennifer,

As per your request, here is a brief description of the research to be conducted with the following cell lines:

1) C2C12 cells, muscle satellite cells (SCs) and primary myoblasts (all engineered to express our PET reporter gene): for transplantation into dystrophic mice; muscle repair/regeneration will be assessed post cell therapy.

2) HEK, 293T cells: our PET reporter construct will be introduced into these producer cell lines. The resulting supernatant will be used to infect cultured myoblasts.

3) C6 cells: used to create gliomas in nude rats

4) LoVo cells: used to create subcutaneous tumors in nude rats

5) NCI-H1299: used to create a lung carcinoma in nude mice

6) PC-3: used to create prostate tumors in nude mice

I hope this will suffice. Please let me know if you require any additional information.

Thanks,

Lisa

Please Remit To:

University of Western Ontario
Financial Services
Accounts Receivable Office
Support Services Building, Suite 6100
London, ON N6A 3K7

Page: 1
Invoice No: T342356
Invoice Date: 12/19/2008
Customer Number: WES001379
Payment Terms: On Receipt
Due Date: 12/19/2008

Bill To:

Healthcare Materials Mgmt Serv-SJHC
Anna Ivanisevic
Accounts Payable
LHSC-UC
London ON N6A 4V2
Canada

Ship To:

AMOUNT DUE: \$ 120.00 CAD

REMITTANCE ADVICE

PLEASE RETURN WITH PAYMENT

\$ _____
Amount Remitted

INVOICE

| Line | Description | Quantity | Unit of Measure | Unit Price | Amount | GST/ HST | PST |
|---------------------------|-------------|----------|-----------------|------------|---------------|------------|-----|
| | Dhanvantari | | | | | | |
| 1 | Inv 12855 | 1.00 | | 120.00 | 120.00 | | |
| TOTAL AMOUNT DUE : | | | | | 120.00 | CAD | |

Billing Inquiries - 519-661-2111 EXT82656

Payment Inquiries - contact Accounts Receivable - Phone: 519/661-3870 Fax: 519/661-3829 Email: fin-aroffice@uwo.ca

| | | |
|---|--|-------------------------------------|
| Invoice Number: T342356 | Issued By: Kathleen Perry | Page: 1 |
| Invoice Date: 12/19/2008 | Issuing Unit: Animal Care & Vet Services | Payment Terms: On Receipt |
| Customer Number: WES001379 | Phone Number: 519-661-2111 EXT82656 | 1.50% per month on overdue accounts |
| Purchase Order Number: | Fax Number: 519/661-2028 | |
| Contract Number: | Bill Type: 003 | |
| GST/Business Number: 10816 2587 RT 0001 | | Amount Remitted: \$ _____ |



University of Western Ontario
Financial Services
Accounts Receivable Office
Support Services Building, Suite 6100
London, ON N6A 3K7

Permit to import human pathogen(s)

Permis d'importation d'agent(s) anthropopathogène(s)

P-1304A

Under the authority of the Human Pathogens Importation Regulations.

Sous le régime du Règlement sur l'importation des agents anthropopathogènes.

Importer-Name, address and postal code - Importateur-Nom, adresse et code postal

Facsimile-Télécopieur

Telephone no.- No. de téléphone

Lawson Health Research Institute
268 Grosvenor Street
London, ON N6A 4V2

(519) 646-6110

(519) 646-6100
ext.: 65738

Attn.: Dr. Savita Dhanvantari

Supplier-Name and address - Fournisseur-Nom et adresse

Name(s) of Port(s) of Entry- To Clear Customs at Port(s) of entry
Nom(s) de(s) point(s) d'entrée -Dédouanement au(x) point(s) d'entrée

Invitrogen Corporation Inc.
1600 Faraday Ave., Carlsbad, CA 92008, USA

Various ports

Description of Pathogen(s)-For the importation of- Description de(s) agent(s) anthropopathogène(s)-Pour l'importation de

ViraPower Promoterless Lentiviral Gateway Expression System (cat# K5910-00)*

*Pathogen(s) indicated on this permit also require an accompanying valid CFIA permit for importation -
*Les agents anthropopathogènes indiqués sur ce permis doivent aussi être accompagnés d'un permis d'importation de l'ACIA.

On the following terms and conditions as marked:-Selon les conditions indiquées:

- 1. Work involving any of the imported material shall be limited to *in vitro* laboratory studies. Les travaux auxquels la matière importée est destinée doivent se limiter à des études de laboratoire *in vitro*.
- 2. Domestic animals, including poultry, cattle, sheep, swine and horses, shall not be directly or indirectly exposed to infection by any of the imported material. Les animaux domestiques, y compris les volailles, bovins, ovins, porcins et chevaux, ne doivent pas être exposés, directement ou indirectement, à l'infection par la matière importée.
- 3. All animals exposed to infection by any of the imported material shall be so exposed and held only in isolated insect-and rodent-proof facilities. Les animaux exposés à l'infection par la matière importée doivent y être exposés et être gardés uniquement dans des installations isolées à l'abri des insectes et des rongeurs.
- 4. All equipment, animal pens, cages, bedding, waste and other articles under the importer's control, that come in direct or indirect contact with any of the imported material, shall be sterilized by autoclaving or incinerated. L'équipement, les enclos pour animaux, les cages, les litières, les déchets et tout autre article sous la responsabilité de l'importateur qui viennent en contact direct ou indirect avec la matière importée doivent être stérilisés par autoclavage ou incinérés.
- 5. Packaging materials, containers and all unused portions of the imported material shall be sterilized by autoclaving or incinerated. Le matériel d'emballage, les récipients et toute partie inutilisée de la matière importée doivent être stérilisés par autoclavage ou incinérés.
- 6. No work on the imported material shall be done, except work conducted or directed by the importer in the facilities described in the application for this permit. NO HUMAN PATHOGEN BELONGING TO RISK GROUP 3 OR 4 MAY BE REMOVED TO ANOTHER LOCATION, OR TRANSFERRED INTO THE POSSESSION OF A PERSON OTHER THAN THE IMPORTER, WITHOUT THE PERMISSION OF THE DIRECTOR. La matière importée ne peut servir qu'aux travaux effectués ou dirigés par l'importateur dans les installations décrites dans la demande de permis. AUCUN AGENT ANTHROPOPATHOGENE DU GROUPE DE RISQUE 3 OU 4 NE PEUT ÊTRE TRANSPORTÉ, SANS LA PERMISSION DU DIRECTEUR, VERS UN AUTRE LIEU OU ÊTRE MIS EN LA POSSESSION D'UNE AUTRE PERSONNE QUE L'IMPORTATEUR.
- 7. On completion of the importer's work involving the imported human pathogen, the pathogen and all its derivatives shall be destroyed. Au terme des travaux de l'importateur auxquels a servi l'agent anthropopathogène importé, celui-ci et tous ses dérivés doivent être détruits.
- 8. Primary isolation, identification and/or manipulation may be done in level 2 containment (physical requirements) using containment level 3 operational requirements. No culturing of Risk Group 3 pathogens shall be done. On peut accomplir l'isolation, l'identification primaire, et/ou la manipulation au niveau de confinement 2 (exigences physiques) en utilisant les exigences opérationnelles de niveau de confinement 3. Aucune culture d'agent anthropopathogène du Groupe de risque 3 ne sera entreprise.
- 9. NO IMPORTED MATERIAL MAY BE REMOVED TO ANOTHER LOCATION, OR TRANSFERRED INTO THE POSSESSION OF A PERSON OTHER THAN THE IMPORTER, WITHOUT THE PERMISSION OF THE DIRECTOR. AUCUNE MATIÈRE IMPORTÉE NE PEUT ÊTRE TRANSPORTÉE, SANS LA PERMISSION DU DIRECTEUR, VERS UN AUTRE LIEU OU ÊTRE MISE EN LA POSSESSION D'UNE AUTRE PERSONNE QUE L'IMPORTATEUR.
- 10. The Director must approve all new work with the imported material involving construction of recombinants that requires an increase of containment from level 2. Tous nouveaux travaux de manipulation génétique (recombiné) avec la matière importée qui demandera que le niveau 2 de confinement soit augmenté exigera l'approbation du Directeur.

11. This permit is valid only for:
Le présent permis n'est valide que pour:

a) a single entry into Canada or
une seule entrée au Canada ou

b) importations at intervals of
les importations effectuées à intervalles de

during the period beginning on
au cours de la période commençant le

and ending on
et se terminant le

September 27, 2006

September 30, 2007

Authorization-Signature of Director
Autorisation-Signature du Directeur

Paul J. Payette, Ph.D.

Date September 27, 2006

Note: Transporting and otherwise dealing with imported material are subject to federal, provincial and municipal laws (if any), to the extent that, those laws apply in respect of that material.

Remarque: Les opérations relatives à la matière importée, y compris le transport, sont assujetties aux lois fédérales, provinciales et aux règlements municipaux applicables.



Permit to import human pathogen(s)

Permis d'importation d'agent(s) anthropopathogène(s)

Under the authority of the Human Pathogens Importation Regulations.

Sous le régime du Règlement sur l'importation des agents anthropopathogènes.

Importer-Name, address and postal code - Importateur-Nom, adresse et code postal

Facsimile-Télécopieur

Telephone no. - No. de téléphone

Lawson Health Research Institute
268 Grosvenor Street
London, ON N6A 4V2

(519) 646-6110

(519) 646-6100
ext.: 65738

Attn.: Dr. Savita Dhanvantari

Supplier-Name and address - Fournisseur-Nom et adresse

Name(s) of Port(s) of Entry - To Clear Customs at Port(s) of entry
Nom(s) de(s) point(s) d'entrée -Dédouanement au(x) point(s) d'entrée

Invitrogen Corporation Inc.
1600 Faraday Ave., Carlsbad, CA 92008, USA

Various ports

Description of Pathogen(s)-For the importation of- Description de(s) agent(s) anthropopathogène(s)-Pour l'importation de

ViraPower Promoterless Lentiviral Gateway Expression System (cat# K5910-00)*.

*Pathogen(s) indicated on this permit also require an accompanying valid CFIA permit for importation -
*Les agents anthropopathogènes indiqués sur ce permis doivent aussi être accompagnés d'un permis d'importation de l'ACIA.

On the following terms and conditions as marked:-Selon les conditions indiquées:

- | | | |
|--|---|--|
| <p>1. Work involving any of the imported material shall be limited to <i>in vitro</i> laboratory studies.</p> <p>2. Domestic animals, including poultry, cattle, sheep, swine and horses, shall not be directly or indirectly exposed to infection by any of the imported material.</p> <p>3. All animals exposed to infection by any of the imported material shall be so exposed and held only in isolated insect-and rodent-proof facilities.</p> <p>4. All equipment, animal pens, cages, bedding, waste and other articles under the importer's control, that come in direct or indirect contact with any of the imported material, shall be sterilized by autoclaving or incinerated.</p> <p>5. Packaging materials, containers and all unused portions of the imported material shall be sterilized by autoclaving or incinerated.</p> <p>6. No work on the imported material shall be done, except work conducted or directed by the importer in the facilities described in the application for this permit. NO HUMAN PATHOGEN BELONGING TO RISK GROUP 3 OR 4 MAY BE REMOVED TO ANOTHER LOCATION, OR TRANSFERRED INTO THE POSSESSION OF A PERSON OTHER THAN THE IMPORTER, WITHOUT THE PERMISSION OF THE DIRECTOR.</p> <p>7. On completion of the importer's work involving the imported human pathogen, the pathogen and all its derivatives shall be destroyed.</p> <p>8. Primary isolation, identification and/or manipulation may be done in level 2 containment (physical requirements) using containment level 3 operational requirements. No culturing of Risk Group 3 pathogens shall be done.</p> <p>9. NO IMPORTED MATERIAL MAY BE REMOVED TO ANOTHER LOCATION, OR TRANSFERRED INTO THE POSSESSION OF A PERSON OTHER THAN THE IMPORTER, WITHOUT THE PERMISSION OF THE DIRECTOR.</p> <p>10. The Director must approve all new work with the imported material involving construction of recombinants that requires an increase of containment from level 2.</p> | <p><input checked="" type="checkbox"/></p> <p><input checked="" type="checkbox"/></p> <p><input checked="" type="checkbox"/></p> <p><input checked="" type="checkbox"/></p> <p><input checked="" type="checkbox"/></p> | <p>Les travaux auxquels la matière importée est destinée doivent se limiter à des études de laboratoire <i>in vitro</i>.</p> <p>Les animaux domestiques, y compris les volailles, bovins, ovins, porcins et chevaux, ne doivent pas être exposés, directement ou indirectement, à l'infection par la matière importée.</p> <p>Les animaux exposés à l'infection par la matière importée doivent y être exposés et être gardés uniquement dans des installations isolées à l'abri des insectes et des rongeurs.</p> <p>L'équipement, les enclos pour animaux, les cages, les litières, les déchets et tout autre article sous la responsabilité de l'importateur qui viennent en contact direct ou indirect avec la matière importée doivent être stérilisés par autoclavage ou incinérés.</p> <p>Le matériel d'emballage, les récipients et toute partie inutilisée de la matière importée doivent être stérilisés par autoclavage ou incinérés.</p> <p>La matière importée ne peut servir qu'aux travaux effectués ou dirigés par l'importateur dans les installations décrites dans la demande de permis. AUCUNE AGENT ANTHROPOPATHOGÈNE DU GROUPE DE RISQUE 3 OU 4 NE PEUT ÊTRE TRANSPORTÉ, SANS LA PERMISSION DU DIRECTEUR, VERS UN AUTRE LIEU OU ÊTRE MIS EN LA POSSESSION D'UNE AUTRE PERSONNE QUE L'IMPORTATEUR.</p> <p>Au terme des travaux de l'importateur auxquels a servi l'agent anthropopathogène importé, celui-ci et tous ses dérivés doivent être détruits.</p> <p>On peut accomplir l'isolation, l'identification primaire, et/ou la manipulation au niveau de confinement 2 (exigences physiques) en utilisant les exigences opérationnelles de niveau de confinement 3. Aucune culture d'agent anthropopathogène du Groupe de risque 3 ne sera entreprise.</p> <p>AUCUNE MATIÈRE IMPORTÉE NE PEUT ÊTRE TRANSPORTÉE, SANS LA PERMISSION DU DIRECTEUR, VERS UN AUTRE LIEU OU ÊTRE MISE EN LA POSSESSION D'UNE AUTRE PERSONNE QUE L'IMPORTATEUR.</p> <p>Tous nouveaux travaux de manipulation génétique (recombiné) avec la matière importée qui demandera que le niveau 2 de confinement soit augmenté exigera l'approbation du Directeur.</p> |
|--|---|--|

11. This permit is valid only for:
Le présent permis n'est valide que pour:

a) a single entry into Canada or
une seule entrée au Canada ou

during the period beginning on
au cours de la période commençant le

and ending on
et se terminant le

September 27, 2006

September 30, 2007

Authorization-Signature of Director
Autorisation-Signature du Directeur

Paul J. Payette, Ph.D.

Date September 27, 2006

Note: Transporting and otherwise dealing with imported material are subject to federal, provincial and municipal laws (if any), to the extent that, those laws apply in respect of that material.

Remarque: Les opérations relatives à la matière importée, y compris le transport, sont assujetties aux lois fédérales, provinciales et aux règlements municipaux applicables.

Canadian Food Inspection Agency
Government of Canada

Agence canadienne d'inspection des aliments
Gouvernement du Canada

Permit No./N° de permis:
A-2007-00178-4
ORIGINAL
2007/01/12
year/mo/day
année/mois/jour

IMPORT PERMIT

PERMIS D'IMPORTATION

Page 1 of/de 3

THIS PERMIT IS ISSUED PURSUANT TO:/CE PERMIS EST DÉLIVRÉ CONFORMÉMENT A:

THE HEALTH OF ANIMALS ACT AND REGULATIONS/LOI ET RÈGLEMENT SUR LA SANTÉ DES ANIMAUX

Importer/Importateur

LAWSON RESEARCH INSTITUTE

268 GROSVENOR STREET, ROOM H417
LONDON, ONTARIO
N6A4V2

Contact: Dr. Savita Dhanvantari Applicant Name: DR. SAVITA
DHANVANTARI
Phone: (519) 646-6100 ext. 65738 Fax: (519) 646-6110

Exporter/Exportateur

INVITROGEN CORPORATION INC.

1600 FARADAY AVENUE
CARLSBAD CALIFORNIA
UNITED STATES
440190

Contact: Mike Galleno

Phone: (760) 603-7219 Fax: (760) 602-6519

Quarantine/Destination/QuarantaineProducer/Producteur

Valid/Valide from/du 2007/01/12 to/au 2008/01/31
year/month/day year/month/day
année/mois/jour année/mois/jour

Country of Origin/
Pays d'Origine

UNITED STATES

For the entry of/ Pour l'entrée de: _____ Single shipment/Chargement simple Multiple shipments/Chargements multiples

Place of entry into Canada/Lieu d'entrée au Canada:
Various Ports of Entry

FOR THE IMPORTATION OF:/POUR L'IMPORTATION DE:

(Description of things(s)/Description de la ou des choses)

1. Product Description: ONE OR MORE OF THE INVITROGEN LENTIVIRAL PRODUCTS LISTED ON THE ATTACHMENT TITLED
"ATTACHMENT TO ANIMAL PATHOGEN(S) IMPORT PERMIT # A-2007-00178-4.

(TO BE USED IN ROOM 4-508, CULTURE ROOM F4-127A, LAWSON HEALTH RESEARCH INSTITUTE, LONDON, ON.) Proposed End Use: "In
Vitro" Scientific Name: Biocontainment Level: 2

**A PERSON WHO IMPORTS A THING UNDER THIS PERMIT SHALL COMPLY WITH ALL THE CONDITIONS SET OUT
HEREIN/TOUTE PERSONNE QUI IMPORTE UNE CHOSE EN VERTU DE CE PERMIS DEVRA RESPECTER TOUTES LES
CONDITIONS DÉCRITES CI-DESSOUS**

Selected Conditions / Conditions Choisies

**ONE OR MORE OF THE INVITROGEN LENTIVIRAL PRODUCTS LISTED ON THE ATTACHMENT TITLED
"ATTACHMENT TO ANIMAL PATHOGEN(S) IMPORT PERMIT # A-2007-00178-4.**

(TO BE USED IN ROOM 4-508, CULTURE ROOM F4-127A, LAWSON HEALTH RESEARCH INSTITUTE, LONDON, ON.)

1. The original or a copy of the signed original of this permit and any other necessary import / export documentation pertaining to the shipment of animal(s)
or thing(s) must be provided for inspection at the first port of entry or to a Canadian Food Inspection Agency Import Service Center.

2. The conditions in this permit can only be changed or amended by a CFIA inspector. Any change to the permit by an unauthorized person will render the
permit invalid.

Canadian Food Inspection Agency
Government of Canada

Agence canadienne d'inspection des aliments
Gouvernement du Canada

Permit No./N° de permis:
A-2007-00178-4
ORIGINAL
2007/01/12
year/mo/day
année/mois/jour

IMPORT PERMIT

PERMIS D'IMPORTATION

Page 2 of/de 3

THIS PERMIT IS ISSUED PURSUANT TO /CE PERMIS EST DÉLIVRÉ CONFORMÉMENT A:

THE HEALTH OF ANIMALS ACT AND REGULATIONS/LOI ET RÈGLEMENT SUR LA SANTÉ DES ANIMAUX

Importer/Importateur

LAWSON RESEARCH INSTITUTE

268 GROSVENOR STREET, ROOM H417
LONDON, ONTARIO
N6A4V2

Contact: Dr. Savita Dhanvantari Applicant Name: DR. SAVITA
DHANVANTARI
Phone: (519) 646-6100 ext. 65738 Fax: (519) 646-6110

Exporter/Exportateur

INVITROGEN CORPORATION INC.

1600 FARADAY AVENUE
CARLSBAD CALIFORNIA
UNITED STATES
440190

Contact: Mike Galleno

Phone: (760) 603-7219 Fax: (760) 602-6519

Selected Conditions / Conditions Choies (Continued/Suite)

3. The imported material must be packaged in appropriate shipping containers to prevent accidental spillage of contents during shipping. Importers should be aware of their obligations under Transport Canada's regulations concerning transportation of dangerous goods.
4. All infectious material must be handled in appropriate animal pathogen containment level 2 facilities as described in Containment Standards for Veterinary Facilities, 1996, AAFC publication no. 1921.
5. The material authorized for importation by this permit is to be used in in vitro studies ONLY and must not to be introduced into laboratory, domestic or wild animals (including birds or fish) unless written authorization is obtained from the Canadian Food Inspection Agency.
6. The animal(s) or thing(s) imported under this permit must not be removed from the premises of destination listed on this permit, unless written authorization is obtained from the Canadian Food Inspection Agency.
7. Upon completion of the tests or experiments, the imported material as described on this permit and any derivatives thereof must be autoclaved, incinerated or alternatively disposed of in a manner approved by an inspector of the Canadian Food Inspection Agency.
8. Records pertaining to the imported product's use, storage and disposal must be maintained for two (2) years following importation. These records must be made available for inspection by the Canadian Food Inspection Agency upon request.
9. The importer is responsible for all costs incurred or associated with any testing or treatment of the animal(s) or thing(s) that may be required under the import permit or under the authority of the Health of Animals Act or the Health of Animals Regulations. The importer shall pay all fees for services required in respect of the importation under the National Animal Health Program Cost Recovery Fees Regulations in place at the time of importation.
10. Consideration of an application necessary for issuance of a permit to import the described animal or thing is subject to Class 1 fees.
11. The issuance of this permit does not relieve the owner or the importer of the obligation to comply with any other relevant federal, provincial or municipal legislation or requirement.
12. Failure to comply with the conditions contained in this permit or with the provisions of the Health of Animals Act and Regulations may result in the cancellation of this permit and will result in the forfeiture to the Crown of the imported thing(s) or in the removal of the thing(s) from Canada, all without compensation to, and at the expense of the importer. The importer(s) are responsible for the imported thing(s), their freedom from extraneous disease, active or latent, and genetic or other defects. The importer, his heirs, executors, successors and assigns release and discharges Her Majesty the Queen in right of Canada and the CFIA of and from all claims and demands, damages, actions or causes of action arising or to arise by reason of the importation of the thing(s) and agrees to indemnify and save harmless Her Majesty the Queen in right of Canada and the CFIA from and against all actions, damages, claims

Canadian Food Inspection Agency
Government of Canada

Agence canadienne d'inspection des aliments
Gouvernement du Canada

Permit No./N° de permis:

A-2007-00178-4

ORIGINAL

2007/01/12

year/mo/day

année/mois/jour

IMPORT PERMIT

PERMIS D'IMPORTATION

Page 3 of/de 3

THIS PERMIT IS ISSUED PURSUANT TO/CE PERMIS EST DÉLIVRÉ CONFORMÉMENT A:

THE HEALTH OF ANIMALS ACT AND REGULATIONS/LOI ET RÈGLEMENT SUR LA SANTÉ DES ANIMAUX

Importer/Importateur

LAWSON RESEARCH INSTITUTE

68 GROSVENOR STREET, ROOM H417
LONDON, ONTARIO
N6A4V2

Contact: Dr. Savita Dhanvantari Applicant Name: DR. SAVITA
DHANVANTARI
Phone: (519) 646-6100 ext. 65738 Fax: (519) 646-6110

Exporter/Exportateur

INVITROGEN CORPORATION INC.

1600 FARADAY AVENUE
CARLSBAD CALIFORNIA
UNITED STATES
440190

Contact: Mike Galleno

Phone: (760) 603-7219 Fax: (760) 602-6519

Selected Conditions / Conditions Choies (Continued/Suite)

and demands which may be brought in respect of or arising out of the importation of such thing(s), any contamination with extraneous disease or other effects.


3. This permit is conditional upon a permit being obtained under the Human Pathogens Importation Regulations to import the pathogenic material and upon that import permit being produced and valid when the above pathogenic material is presented to an inspector for inspection at the time of importation.

Additional Conditions Additionnelles

ONE OR MORE OF THE INVITROGEN LENTIVIRAL PRODUCTS LISTED ON THE ATTACHMENT TITLED "ATTACHMENT TO ANIMAL PATHOGEN(S) IMPORT PERMIT # A-2007-00178-4.

TO BE USED IN ROOM 4-508, CULTURE ROOM F4-127A, LAWSON HEALTH RESEARCH INSTITUTE, LONDON, ON.)

No culturing of containment level 3 or 4 pathogens shall be done.


Authorized By:/Approuvé par:
CINTHIA LABRIE

For the Minister of Agriculture and Agri-Food
Pour le ministre d'agriculture et agroalimentaire



Canadian Food
Inspection Agency

Agence canadienne
d'inspection des aliments



Office of Biohazard Containment and Safety
Science Advice and Biohazards Division
Science Strategies Directorate, CFIA
159 Cleopatra Drive, Ottawa, Ontario K1A 0Y9
Tel: (613) 221-7068 Fax: (613) 228-6129
Email: ImportZoopath@inspection.gc.ca

Bureau du confinement des biorisques et sécurité
Division des avis scientifiques et contrôle des biorisques
Direction des stratégies scientifiques, ACIA
159 promenade Cleopatra, Ottawa, Ontario K1A 0Y9
Tél: (613) 221-7068 Téléc: (613) 228-6129
Courriel: ImportZoopath@inspection.gc.ca

**ATTACHMENT TO ANIMAL PATHOGEN(S) IMPORTATION PERMIT
ATTACHEMENT AU PERMIS D'IMPORTATION D'AGENTS ZOOPATHOGÈNES**

#A-2007-00178-4

Issued to/ Délivré à: Dr. Savita Dhanvantari, Lawson Health Research Institute,
268 Grosvenor Street, London ON N6A 4V2.

Includes the following animal pathogen containment Level 2 microorganisms:
Inclut les agents zoopathogènes de niveau de confinement 2 suivant:

Invitrogen Lentiviral Products / Produits Lentiviral d'Invitrogen:

- PCDNA6.2/C-EMGFP-GW/TOPO (K35920)
- PCDNA6.2/N-EMGFP-GW/TOPO (K36020)
- PCDNA6.2/C-YFP-GW/TOPO (K36120)
- PCDNA6.2/N-YFP-GW/TOPO (K36620)
- Virapower II Lenti GW System (K36720)
- Virapower II Lenti C-Lumio system (K37020)
- Virapower II Lenti N-Lumio system (K37120)
- POL III MIR Rnai Vector (K493500)
- POL II MIR Rnai GFP Vector (K493600)
- Lenti POL II MIRE Rnai Vector (K493700)
- Lenti POL II MIRE Rnai w/GFP (K493800)
- Block it Lenti RNAi Expression system (K494400)
- Virapower Lentiviral directional (K495000)
- Virapower Lentiviral Gateway (K496000)
- Lentiviral T Rex Expression system (K496500)
- Virapower packaging mix (K497500)
- Virapower Zeo Lenti Expression (K498000)
- Virapower Zeo Lentiviral Support Kit (K498500)
- Virapower UBC Lenti expression (K499000)
- Virapower Lentiviral support (K497000)
- Plenti6/Block it RNAi vector (K494300)
- Plenti 6/V5 Directional TOPO (K495510)
- VP TR GW Vector kit (K496700)
- PCDNA6.2/EMGFP-BSD/V5 Dest (V36620)
- Plenti6.2/V5-DEST GW vector (V36820)
- Plenti6.2-GW/EMGFP Exp vector (V36920)
- Plenti6/TR vector (V48020)
- Block-iT Lenti RNAi ZW GW Vector (V48820)
- Plenti6/V5 Gtwy vector pack (V49610)
- Plenti4/V5 -Dest Gateway vector (V49810)
- Plenti6/UBC/V5 Dest vector (V49910)
- Block it Lentiviral Inducible RNAi (K492500)
- Promotorless Lenti Exp kit (K591000)

The above products may contain one or more of the following components / Les produits ci-dessus peuvent contenir un ou plusieurs des composants suivants:

Plenti6/Block it Dest RNAi , PLP1, PLP2, PLP3/VSVG, Plenti6/V5-Dtopo, Plenti6/V5-GW/LacZ, plenti6/V5 Dest vector, plenti6/TR, plenti4/TO/V5 Dest, plenti4/TO/V5-GW/LacZ, plenti 4/V5 Dest, plenti4/V5 -GW/LacZ vector, plenti4/Blockit Dest, plenti6/UBC/ V5 Dest vector, plenti6/UBC/V5-GW/LacZ vector, plenti6/R4R2/V5-Dest, 293 FT cells, PCDNA6.2-GW/MIR Neg TB, PCDNA6.2-GW-EMGFP-MIR Neg, Plenti6.2/C-Lumio/V5 DEST, Plenti 6.2/C-Lumio-V5-GW/LA, Plenti6.2/N-Lumio/V5 Dest, Plenti6.2/N-Lumio/V5-GW/LA, Plenti6.2-GW/EMGFP Kit, Plenti 6.2.V5 Dest Kit, Plenti6.2/V5-GW LacZ, PCDNA6.2/EMGFP-BSD/V5 Dest, PCDNA6.2/EMGFP-BSD/V5-GW/C, PCDNA6.2/C-EMGFP-GW, PCDNA6.2/C-EMGFP-GW/CAT, PCDNA6.2/N-EMGFP-GW, PCDNA6.2/N-EMGFP-GW, PCDNA6.2/C-YFP-GW, PCDNA6.2 C-YFP-GW/CAT, PCDNA6.2/N-YFP-GW, PCDNA6.2/N-YFP-GW-CAT.

REVISED: May 01, 2006.

Cynthia Labrie
Cynthia Labrie

A/Chief, Animal Pathogen Importation Program/
Chef intérimaire, Programme d'importation des agents zoopathogènes

Jan 12/07
Date

Canada



* IMPORTANT NOTICE *

Your file Votre référence

Our file Notre référence

1) ZONOTIC IMPORTS: Please check the “Description of Pathogen(s)” section of your attached permit, and if the following message (in red print) has been included: “***Pathogen(s) indicated on this permit also require an accompanying valid CFIA permit for importation.**”, then the material is of a **zoonotic** nature and a valid permit from the Canadian Food Inspection Agency (CFIA) is required for this importation in addition to your attached human pathogens import permit. If you do not have a valid permit from the Canadian Food Inspection Agency, please contact them directly for assistance at: (613) 221-7068.

2) INSTRUCTIONS FOR USE OF YOUR PERMIT:
[as per the *Human Pathogen(s) Importation Regulations (SOR/94-558)*]

Prior to shipment of the human pathogen described in the Import Permit the importer **must:**

- a) provide a copy of the importation permit to the supplier and notify the supplier that **a copy of the importation permit must be attached to each shipment;**
- b) **notify the supplier** that the outer shipping container in which the human pathogen is transported must display clearly, on the outside surface of the container, the importation permit number and the following statement immediately preceding that number:

“Human Pathogen – Importation Permit Number:/Agent anthropopathogène – Numéro du permis d’importation:”

If the permit holder who arranges to import a human pathogen that belongs to Risk Group 3 or 4, does not receive the human pathogen on, or within three (3) days after, such date of receipt as may reasonably be expected in the circumstances, he shall forthwith give to the Director, Office of Laboratory Security a notice that the human pathogen has not been received and provide the Director with the importation permit number.

To facilitate Customs clearance, a copy of the importation permit should be kept by the importer and presented to Customs or sent to the importer’s customs broker.

3) Please note that importation of this material may also be subject to the requirements of the *New Substances Notification Regulations (Organisms)* of the *Canadian Environmental Protection Act, 1999*, administered by Environment Canada and Health Canada. Please contact the New Substances Information Line at 1-800-567-1999 or nsn-infoline@ec.gc.ca for assistance.

Direct inquiries to:

Office of Laboratory Security
Public Health Agency Canada
Centre for Emergency Preparedness and Response
100 Colonnade Road, Loc.: 6201A
Ottawa, Ontario K1A 0K9

Tel.: (613) 957-1779
Fax: (613) 941-0596



Canadian Food
Inspection Agency

Agence canadienne
d'inspection des aliments



Office of Biohazard Containment and Safety
Science Advice and Biohazards Division
Science Strategies Directorate, CFIA
159 Cleopatra Drive, Ottawa, Ontario K1A 0Y9

Bureau du confinement des biorisques et sécurité
Division des avis scientifiques et contrôle des biorisques
Direction des stratégies scientifiques, ACIA
159 promenade Cleopatra, Ottawa, Ontario K1A 0Y9

FACSIMILE TRANSMITTAL NOTICE / TRANSMISSION PAR TÉLÉCOPIEUR

| | | | |
|---|----------------------------------|---|-----------------------------|
| To / À: Dr. Sativa Dhanvantari Lawson Research Institute | | From / De: Andrew Halliday Animal Pathogen Import Program / Programme d'importation des agents zoopathogènes | |
| Facsimile/télécopieur. | 519-646-6110 | Facsimile/télécopieur. | 613-228-6129 |
| Subject/Objet: Importation of animal pathogens / Importation d'agents zoopathogènes | | | |
| Message: | | | |
| Please find attached / Veuillez trouver ci-joint : | | | |
| <input type="checkbox"/> A copy of a Non-pathogenic letter for the product(s) you requested. / Une copie de la lettre de non-pathogénéicité pour le(s) produit(s) demandé(s). | | | |
| <input checked="" type="checkbox"/> A copy of the import permit for which you applied. Please review the conditions appearing on your permit. / Une copie de votre permis d'importation. Veuillez s'il-vous-plaît prendre note des conditions apparaissant sur votre permis. | | | |
| <input checked="" type="checkbox"/> Condition # 13: The product(s) requested is(are) also regulated by the Public Health Agency of Canada (PHAC). Please contact PHAC at (613) 957-1779. / Le(s) produit(s) demandé(s) sont également réglementés par l'Agence de santé publique du Canada (ASPC). Veuillez contacter l'ASPC au (613)-957-1779. | | | |
| Andrew Halliday importzoopath@inspection.gc.ca | | | |
| Please visit our website at: http://www.inspection.gc.ca/english/sci/bio/bioe.shtml . Veuillez visiter notre site internet au: http://www.inspection.gc.ca/francais/sci/bio/biof.shtml . | | | |
| Signature: | Date: January 12, 2007 | Telephone/Téléphone: 613-221-7068 | No./Nbre Pages: 5 |



QUOTATION

IN RESPONSE TO YOUR INQUIRY

TO ORDER:
 Invitrogen Canada Inc
 2270 Industrial Street, Burlington, ON L7P 1A1
 To Order: (800) 263-6236
 Fax No.: (800) 387-1007
 E-mail: caorders@invitrogen.com

TO:
 LAWSON RESEARCH INSTITUTE
 FOR: Dr. Dhanvantari/Lisa Hoffman

 LONDON
 ON N6A 4V2 Canada
 ATTN: Lisa Hoffman

QUOTATION NO.: S6912311 _ B
 To ensure correct pricing and terms, the above quote number must appear on all orders and correspondence.
FROM: 07/21/2006 **THROUGH:** 07/20/2007
 EXCEPT WHERE NOTED BELOW
TERMS: NET 30 DAYS
ESTIMATED DELIVERY: DAYS, A.R.O.
FOB: Shipping Point

To place an order please call Customer Service
 1-800-263-6236

WE ARE PLEASED TO QUOTE ON YOUR REQUIREMENTS AS FOLLOWS:
NOUS AVONS LE PLAISIR DE VOUS ENVOYER LA SOUMISSION CORRESPONDANT À VOTRE REQUÊTE :

Natalie Shier Territory Sales Manager 661

| ITEM NO | CATALOG NO | DESCRIPTION | QUALIFYING LIMIT | PRICE OR % DISCOUNT | |
|---------|------------|---|------------------|---------------------|---------------|
| | | | | DISCOUNT/UNIT | EXTENDED/UNIT |
| 1 | K591000 | ViraPower™ Promoterless Lentiviral Gateway® Expression System with MultiSite™ Gateway® Technology 1 kit | 1+ | \$1,575.00 | \$1,575.00 |

TERMS AND CONDITIONS

Telephone Number:

(Continue)

THESE GOODS ARE FOR RESEARCH ONLY, UNLESS OTHERWISE SPECIFIED. SEE "AUTHORISED USERS" IN GENERAL TERMS AND CONDITIONS.
 À moins d'indications contraires, ces produits sont destinés à la recherche. Voir la section "Utilisations autorisées" dans les conditions générales.



QUOTATION

IN RESPONSE TO YOUR INQUIRY

TO ORDER:

Invitrogen Canada Inc
2270 Industrial Street, Burlington, ON L7P 1A1
To Order: (800) 263-6236
Fax No: (800) 387-1007
E-mail: caorders@invitrogen.com

TO:

LAWSON RESEARCH INSTITUTE
FOR: Dr. Dhanvantari/Lisa Hoffman

LONDON
ON N6A 4V2 Canada
ATTN: Lisa Hoffman

QUOTATION NO.: S6912311 _ B

To ensure correct pricing and terms, the above quote number must appear on all orders and correspondence.

FROM: 07/21/2006 **THROUGH:** 07/20/2007

EXCEPT WHERE NOTED BELOW

TERMS: NET 30 DAYS

ESTIMATED DELIVERY: DAYS, A.R.O.

FOB: Shipping Point

To place an order please call Customer Service
1-800-263-6236

WE ARE PLEASED TO QUOTE ON YOUR REQUIREMENTS AS FOLLOWS:

NOUS AVONS LE PLAISIR DE VOUS ENVOYER LA SOUMISSION CORRESPONDANT À VOTRE REQUÊTE :

TERMS AND CONDITIONS

ADDITIONAL TERMS AND CONDITIONS OF QUOTATION

1. General Terms and Conditions listed on the customer copy of packing lists and invoices from Invitrogen Corporation will apply except where otherwise agreed in writing by an authorized representative of Invitrogen Corporation.
2. In order to receive quoted prices, the quotation number must be referenced at time of order. Credits will not be issued for orders not referencing quotation numbers.
3. The effective dates of this quotation appear in the upper right corner of each page unless otherwise noted. Exceptions are noted within the body of this quotation.
4. The quantities noted on this quotation reflect minimum order requirements necessary to receive quoted prices.
5. Percentage discounts will be calculated from current list price.
6. This quotation may be terminated by Invitrogen Corporation upon written notice.
7. This quotation contains confidential Invitrogen Corporation pricing information which if disclosed to third parties could cause competitive harm to Invitrogen Corporation. Subject to overriding obligations to third party funding agencies or governmental entities, the customer agrees to keep all pricing information contained herein confidential...

IF OUR SUPPLIER COSTS CHANGE DURING THE DURATION OF THIS QUOTE, YOUR PRICES MAY BE ADJUSTED.

Telephone Number:

(Continue)

THESE GOODS ARE FOR RESEARCH ONLY, UNLESS OTHERWISE SPECIFIED. SEE "AUTHORISED USERS" IN GENERAL TERMS AND CONDITIONS.
À moins d'indications contraires, ces produits sont destinés à la recherche. Voir la section "Utilisations autorisées" dans les conditions générales.



Public Health
Agency of Canada

Agence de santé
publique du Canada

Date: September 27, 2006

Your file Votre référence

Importer address: Lawson Health Research Institute
268 Grosvenor St.
London, ON
N6A 4V2

Our file Notre référence

Dear Dr. Savita Dhanvantari,

Enclosed you will find your Public Health Agency of Canada permit to import human pathogen(s), **P-13043**.

Due to the nature of the material requested for import, some additional conditions apply. Please review and note the conditions of import, in particular conditions #8, #9 and #10. Condition #8 states that "Primary isolation, identification and/or manipulation may be done in level 2 containment (physical requirements) using containment level 3 operational requirements. No culturing of Risk Group 3 pathogens shall be done". Condition #9 states that "No imported material may be removed to another location, or transferred into the possession of a person other than the importer, without the permission of the director [of the Office of Laboratory Security, Public Health Agency of Canada]". Condition #10 states that "The Director [of the Office of Laboratory Security, Public Health Agency of Canada] must approve all new work with the imported material involving construction of recombinants that require an increase of containment from level 2".

If you have any questions or comments regarding this matter, please do not hesitate to contact our office.

Sincerely,

Paul J. Payette, Ph.D.
Director, Office of Laboratory Security
Centre for Emergency Preparedness and Response
100 Colonnade Road, Loc.: 6201A
Ottawa, Ontario, Canada K1A 0K9
Phone: (613) 957-1779
Fax: (613) 941-0596

Encl.



* IMPORTANT NOTICE *

Your file Votre référence

Our file Notre référence

1) ZONOTIC IMPORTS: Please check the “**Description of Pathogen(s)**” section of your attached permit, and **if** the following message (in red print) has been included: “***Pathogen(s) indicated on this permit also require an accompanying valid CFIA permit for importation.**”, then the material is of a **zoonotic** nature and a valid permit from the Canadian Food Inspection Agency (CFIA) is required for this importation in addition to your attached human pathogens import permit. If you do not have a valid permit from the Canadian Food Inspection Agency, please contact them directly for assistance at: **(613) 221-7068.**

7088 ANDREW

2) INSTRUCTIONS FOR USE OF YOUR PERMIT:
[as per the *Human Pathogen(s) Importation Regulations (SOR/94-558)*]

Prior to shipment of the human pathogen described in the Import Permit the importer **must:**

- a) provide a copy of the importation permit to the supplier and notify the supplier that **a copy of the importation permit must be attached to each shipment;**
- b) **notify the supplier** that the outer shipping container in which the human pathogen is transported must display clearly, on the outside surface of the container, the importation permit number and the following statement immediately preceding that number:

“Human Pathogen – Importation Permit Number:/Agent anthropopathogène – Numéro du permis d’importation:”

If the permit holder who arranges to import a human pathogen that belongs to Risk Group 3 or 4, does not receive the human pathogen on, or within three (3) days after, such date of receipt as may reasonably be expected in the circumstances, he shall forthwith give to the Director, Office of Laboratory Security a notice that the human pathogen has not been received and provide the Director with the importation permit number.

To facilitate Customs clearance, a copy of the importation permit should be kept by the importer and presented to Customs or sent to the importer’s customs broker.

3) Please note that importation of this material may also be subject to the requirements of the *New Substances Notification Regulations (Organisms)* of the *Canadian Environmental Protection Act, 1999*, administered by Environment Canada and Health Canada. Please contact the New Substances Information Line at 1-800-567-1999 or nsn-infoline@ec.gc.ca for assistance.

Direct inquiries to:

Office of Laboratory Security

Public Health Agency Canada

Centre for Emergency Preparedness and Response

100 Colonnade Road, Loc.: 6201A

Ottawa, Ontario K1A 0K9

Tel.: (613) 957-1779

Fax: (613) 941-0596



Public Health
Agency of Canada

Agence de santé
publique du Canada

Dear Sir/Madam:

Please find attached, for your convenience and future use, an application form for a Public Health Agency of Canada permit to import human pathogens. When filling out this form, please note the following directives:

- If ordering from a commercial supplier (e.g. ATCC), please provide the product name, catalogue number and any relevant descriptive information. If the product is coming from another researcher, please provide background information (references, etc.).

- If your work objectives (Box 10) include *in vivo* activities, please describe in full, including animal species used.

When completed, please forward the **original application** to our office at the following address:

Office of Laboratory Security
Public Health Agency of Canada
Centre for Emergency Preparedness
and Response
100 Colonnade Road, Loc.: 6201A
Ottawa, Ontario
K1A 0K9

Tel.: (613) 957-1779
Fax: (613) 941-0596

Upon receipt of your application, a permit will be issued and faxed back to you (usually within 5 working days) and the original will follow through regular mail.

If you have any questions regarding this matter, please do not hesitate to contact our office.

Thank you for your collaboration.

Your file Votre référence

Our file Notre référence

Monsieur, Madame,

Vous trouverez sous pli, pour utilisation future, un formulaire de demande de permis de l'Agence de santé publique du Canada pour l'importation d'agent(s) anthropopathogène(s). Nous vous prions de tenir compte des directives suivantes lorsque vous complétez le formulaire :

- Si vous commandez d'un fournisseur commercial (p.ex. l'ATCC), prière de nous fournir le nom du produit, le numéro de catalogue et toute l'information et/ou description qui s'y rattache. Si le produit doit vous parvenir d'un autre chercheur, prière de nous fournir toute l'information pertinente (références, etc.).

- Si l'objectif de votre travail (Section 10) comprend des activités *in vivo*, prière de nous fournir une description complète, incluant les espèces d'animaux utilisés.

Une fois complété, veuillez nous faire parvenir la **copie originale du formulaire de demande de permis** à l'adresse suivante :

Bureau de la sécurité des laboratoires
Agence de santé publique du Canada
Centre de mesures et d'interventions d'urgence
100 chemin Colonnade, Loc.: 6201A
Ottawa, Ontario
K1A 0K9

Tél.: (613) 957-1779
Fax: (613) 941-0596

Nous pouvons émettre un permis et vous le faire parvenir par télécopieur, habituellement dans les cinq jours suivant la réception du formulaire de demande. La copie originale du permis vous parviendra ensuite par la poste.

Pour de plus amples renseignements, n'hésitez pas à entrer en contact avec notre bureau.

Merci de votre collaboration.

Kit Contents and Storage

Types of Kits

This manual is supplied with the following products.

| Product | Catalog no. |
|--------------------------------|-------------|
| 293FT Cell Line | R700-07 |
| BioModule™ Lentiviral 293 Unit | WFGE08-S |

Kit Components

The 293FT Cell Line and BioModule™ Lentiviral 293 Unit include the following components. For detailed contents, see the following pages.

The 293FT Cell Line and BioModule™ Lentiviral 293 Unit are shipped as described below. Upon receipt, store each item as detailed below.

| Component | Catalog no. | | Shipping | Storage |
|---|-------------|----------|------------------|-------------------------------|
| | R700-07 | WFGE08-S | | |
| 293FT Cell Line | √ | | Dry ice | Liquid nitrogen |
| Dulbecco's Modified Eagle Medium (D-MEM) | | √ | Room Temperature | 2°C to 8°C |
| 10 mM MEM Non-Essential Amino Acids Solution (100X) | | √ | Room Temperature | 2°C to 8°C |
| MEM Sodium Pyruvate Solution (100X) | | √ | Room Temperature | 2°C to 8°C |
| Phosphate-Buffered Saline, pH 7.4 | | √ | Room Temperature | 2°C to 8°C |
| Opti-MEM® I Reduced Serum Medium | | √ | Room Temperature | 2°C to 8°C (keep in the dark) |
| Geneticin® Selective Antibiotic (50 mg/ml) | | √ | Room Temperature | -20°C or 2°C to 8°C |
| Trypan Blue Stain | | √ | Room Temperature | Room Temperature |
| Fetal Bovine Serum | | √ | Dry ice | -5° to -20°C |
| 200 mM L-Glutamine (100X) | | √ | Dry ice | -5° to -20°C |
| Penicillin-Streptomycin | | √ | Dry ice | -5° to -20°C |
| Trypsin-EDTA | | √ | Dry ice | -5° to -20°C |

Continued on next page

Kit Contents and Storage, continued

293FT Cell Line

The 293FT Cell Line is used for the production of lentiviral stocks. The 293FT Cell Line is supplied as one vial containing 3×10^6 frozen cells in 1 ml of Freezing Medium. **Upon receipt, store in liquid nitrogen until use.**



Handle as potentially biohazardous material under at least Biosafety Level 2 containment. This product contains Dimethyl Sulfoxide (DMSO), a hazardous material. Review the Material Safety Data Sheet before handling.

BioModule™ Lentiviral 293 Unit

The following reagents are provided with the BioModule™ Lentiviral 293 Unit:

| Component | Composition | Quantity |
|---|---|----------------|
| Dulbecco's Modified Eagle Medium | D-MEM high glucose (1X), containing 4,500 mg/L D-glucose, and 4 mM L-glutamine, but no sodium pyruvate. | 2 x 1000 ml |
| 10 mM MEM Non-Essential Amino Acids Solution (100X) | 890 mg/L L-Alanine 1320 mg/L L-Asparagine 1330 mg/L L-Aspartic Acid 1470 mg/L L-Glutamic Acid 750 mg/L Glycine 1150 mg/L L-Proline 1050 mg/L L-Serine | 100 ml |
| MEM Sodium Pyruvate Solution (100X) | 100 mM Sodium Pyruvate Solution (11,004 mg/L) | 100 ml |
| Phosphate-Buffered Saline, pH 7.4 | 0.144 g/L KH_2PO_4 9.00 g/L NaCl 0.795 g/L Na_2HPO_4 pH 7.4 | 500 ml |
| Opti-MEM® I Reduced Serum Medium | See below | 500 ml |
| Geneticin® Selective Antibiotic (50 mg/ml) | 50 mg/ml active Geneticin® Selective Antibiotic in distilled water | 20 ml |
| Trypan Blue Stain | 0.4% Trypan Blue solution in 0.85% NaCl | 100 ml |
| Fetal Bovine Serum | Fetal Bovine Serum, Certified (US) | 2 x 100 ml |
| 200 mM L-Glutamine (100X) | 200 mM L-Glutamine (29.2 mg/ml) in 0.85% NaCl | 100 ml |
| Penicillin-Streptomycin | 5,000 units/ml penicillin (base) 5,000 µg/ml streptomycin (base) in 0.85% NaCl | 100 ml |
| Trypsin-EDTA | 0.5 g/L trypsin (1:250) 0.2 g/L EDTA•4Na in Hanks' Balanced Salt Solution without CaCl_2 , $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ Contains phenol red | 100 ml |

Continued on next page

Overview

Introduction

This manual is provided with the 293FT Cell Line and BioModule™ Lentiviral 293 Unit. The 293FT Cell Line is a very suitable host for lentiviral production, while the BioModule™ Lentiviral 293 Unit contains the reagents for optimal growth and lentiviral production of the 293FT Cell Line. Below the characteristics of the 293FT Cell Line and BioModule™ Lentiviral 293 Unit are explained.

293FT Cell Line

The 293FT Cell Line is derived from the 293F Cell Line (see below) and stably expresses the SV40 large T antigen from the pCMVSPORT6TAg.neo plasmid. Expression of the SV40 large T antigen is controlled by the human cytomegalovirus (CMV) promoter and is high-level and constitutive. For more information about pCMVSPORT6TAg.neo, see the **Appendix**, page 10.

Use of the Cell Line

Studies have demonstrated maximal virus production in human 293 cells expressing SV40 large T antigen (Naldini *et al.*, 1996), making the 293FT Cell Line a particularly suitable host for generating lentiviral constructs using the ViraPower™ Lentiviral Expression System available from Invitrogen (Catalog nos. K4950-00 and K4960-00).

Parental Cell Lines

The 293 Cell Line is a permanent line established from primary embryonal human kidney transformed with sheared human adenovirus type 5 DNA (Graham *et al.*, 1977; Harrison *et al.*, 1977). The E1A adenovirus gene is expressed in these cells and participates in transactivation of some viral promoters, allowing these cells to produce very high levels of protein.

The 293-F Cell Line available from Invitrogen (Catalog no. 11625) is a fast-growing variant of the 293 cell line, and was originally obtained from Robert Horlick at Pharmacoceia.

Antibiotic Resistance

293FT cells stably express the neomycin resistance gene from pCMVSPORT6TAg.neo and should be maintained in medium containing Geneticin® at the concentration listed below. Expression of the neomycin resistance gene in 293FT cells is controlled by the SV40 enhancer/promoter.

Continued on next page

Introduction

Overview

Introduction

The ViraPower™ Promoterless Lentiviral Gateway® Expression System combines Invitrogen's ViraPower™ Lentiviral and MultiSite Gateway® technologies to facilitate lentiviral-based expression of a gene of interest from any promoter of choice in dividing or non-dividing mammalian cells. The System includes:

- The pENTR™5'-TOPO® TA Cloning Kit for production of an entry clone containing your eukaryotic promoter of interest. The pENTR™5'-TOPO® entry vector is adapted with MultiSite Gateway® Technology to facilitate transfer of the promoter sequence into the lentiviral expression plasmid.
- A promoterless pLenti6/R4R2/V5-DEST destination vector into which the promoter and gene of interest are transferred. This expression plasmid contains elements that allow packaging of the construct into virions and the Blasticidin resistance marker for selection of stably transduced cell lines.
- Components of the ViraPower™ Lentiviral System (Catalog no. K5910-00 only) for production of a replication-incompetent lentivirus that transiently or stably expresses the gene of interest in both dividing and non-dividing mammalian cells.

For more information about the ViraPower™ Lentiviral Technology and the MultiSite Gateway® Technology, see pages 6-7.

Advantages of the ViraPower™ Promoterless Lentiviral Gateway® Expression System

Use of the ViraPower™ Promoterless Lentiviral Gateway® Expression System to facilitate lentiviral-based expression of the gene of interest provides the following advantages:

- Allows production of a lentiviral construct that facilitates expression of a gene of interest under the control of a promoter of choice.
- Generates replication-incompetent lentivirus that effectively transduces both dividing and non-dividing mammalian cells, thus broadening the potential applications beyond those of traditional retroviral systems (Naldini, 1998).
- Efficiently delivers the gene of interest to mammalian cells in culture or *in vivo* (Dull *et al.*, 1998).
- Provides stable, long-term expression of a target gene beyond that offered by adenoviral-based systems (Dull *et al.*, 1998; Naldini *et al.*, 1996).
- Produces a pseudotyped virus with a broad host range (Yee *et al.*, 1994).
- The expression vector in the System is adapted with MultiSite Gateway® Technology for easy, simultaneous, recombination-based cloning of multiple DNA fragments in a defined order and orientation.
- Includes multiple features designed to enhance the biosafety of the system.

continued on next page

Overview, continued

ViraPower™ Lentiviral Technology

The ViraPower™ Lentiviral Technology facilitates highly efficient, *in vitro* or *in vivo* delivery of a target gene or RNA to dividing and non-dividing mammalian cells using a replication-incompetent lentivirus. Based on the lentikat™ system developed by Cell Genesys (Dull *et al.*, 1998), the ViraPower™ Lentiviral Technology possesses features which enhance its biosafety while allowing high-level expression in a wider range of cell types than traditional retroviral systems. The main components of the ViraPower™ Lentiviral Expression System include:

- A pLenti-based expression vector into which the DNA sequence (or sequences) are cloned. This vector contains elements required to allow packaging of the expression construct into virions and an antibiotic resistance marker to allow selection of stably transduced cell lines. For more information, see page 5.
- The ViraPower™ Packaging Mix, an optimized mixture of the three packaging plasmids required for production of the lentivirus.
- A 293FT producer cell line to facilitate optimal production of virus.

For more information about the ViraPower™ lentiviral components in this kit, see page 4. For more information about the biosafety features of the System, see page 8.

Purpose of this Manual

This manual provides an overview of the ViraPower™ Promoterless Lentiviral Gateway® Expression System and provides instructions and guidelines to:

1. Generate entry clones containing the promoter and gene of interest, one in pENTR™5'-TOPO® and the second in any Gateway® entry vector (guidelines only provided).
2. Use the pLenti6/R4R2/V5-DEST vector and two entry clones containing the promoter and gene of interest in a MultiSite Gateway® LR recombination reaction to generate an expression clone.
3. Cotransfect the pLenti6/R4R2/V5-DEST expression construct and the ViraPower™ Packaging Mix into the 293FT Cell Line to produce a lentiviral stock.
4. Titer the lentiviral stock.
5. Transduce the mammalian cell line of choice with the Lenti6/R4R2/V5-DEST lentiviral construct.
6. Assay for “transient” expression of your recombinant protein or generate a stably transduced cell line, if desired.

For details and instructions to generate the entry clone containing the promoter of interest, refer to the pENTR™5'-TOPO® TA Cloning Kit manual. For instructions to generate the entry clone containing the gene of interest, refer to the manual for the entry vector you select. For instructions to culture and maintain the 293FT producer cell line, refer to the 293FT Cell Line manual. The pENTR™5'-TOPO® TA Cloning® Kit and 293FT Cell Line manuals are supplied with Catalog no. K5910-00. All manuals are available for downloading from www.invitrogen.com or by contacting Technical Support (see page 56).

continued on next page

Overview, continued




Important

The ViraPower™ Promoterless Lentiviral Expression System is designed to help you create a lentivirus to deliver and express a gene of interest from a promoter of choice in mammalian cells. Although the system has been designed to help you express your recombinant protein of interest in the simplest, most direct fashion, use of the system is geared towards those users who are familiar with the principles of retrovirus biology and retroviral vectors. In addition, we highly recommend that users possess a working knowledge of:

- Viral and tissue culture techniques
- Gateway® Technology and site-specific recombination

For more information about these topics, refer to the following published reviews:

- Retrovirus biology and the retroviral replication cycle: see Buchschacher and Wong-Staal (2000) and Luciw (1996).
 - Retroviral and lentiviral vectors: see Naldini (1999), Naldini (1998), and Yee (1999)
 - Gateway® Technology and site-specific recombination: see Hartley *et al.* (2000) and Landy (1989)
-



Note

The One Shot® Stbl3™ Chemically Competent *E. coli*, LR Clonase™ II Plus Enzyme Mix, and Lipofectamine™ 2000 Reagent included in the ViraPower™ Promoterless Lentiviral Gateway® Expression System are available separately from Invitrogen and are each supplied with individual documentation detailing general use of the product. **For instructions to use these products specifically with the ViraPower™ Promoterless Lentiviral Gateway® Expression System, follow the recommended protocols in this manual.**

The ViraPower™ Promoterless Lentiviral Gateway® Expression System

Components of the ViraPower™ Promoterless Lentiviral Gateway® Expression System

The ViraPower™ Promoterless Lentiviral Gateway® Expression System facilitates highly efficient, lentiviral-based, *in vitro* or *in vivo* expression of a gene of interest under the control of a promoter of choice in dividing and non-dividing mammalian cells. The kit includes the following major components:

- The pENTR™5'-TOPO® TA Cloning Kit containing the pENTR™5'-TOPO® vector for production of an entry clone containing the promoter of interest. The vector is TOPO®-adapted and MultiSite Gateway®-adapted to allow TOPO® Cloning of a *Taq* polymerase-amplified PCR product encoding the promoter of interest and easy transfer of the promoter sequence into the pLenti6/R4R2/V5-DEST vector, respectively. For more information about the MultiSite Gateway® Technology, see page 6. For detailed information about the pENTR™5'-TOPO® vector and instructions to generate an entry clone, refer to the pENTR™5'-TOPO® TA Cloning® Kit manual.

Important: To generate the pLenti6/R4R2/V5-DEST expression construct, you will also need to generate an entry clone containing your gene of interest. In this instance, you may use any standard Gateway® entry vector except pENTR™5'-TOPO®. For more information, see page 6.

- The pLenti6/R4R2/V5-DEST expression vector into which the promoter and gene of interest will be simultaneously cloned using MultiSite Gateway® Technology. The vector also contains the elements required for packaging of the expression construct into virions (e.g. 5' and 3' LTRs, ψ packaging signal) and the Blasticidin resistance marker to allow generation of stable cell lines. For more information about the pLenti6/R4R2/V5-DEST vector, see page 5.
- The ViraPower™ Packaging Mix that contains an optimized mix of the three packaging plasmids, pLP1, pLP2, and pLP/VSVG. These plasmids supply the helper functions as well as structural and replication proteins *in trans* required to produce the lentivirus. For more information about the packaging plasmids, see the **Appendix**, pages 50-55.
- An optimized 293FT producer cell line that stably expresses the SV40 large T antigen under the control of the human CMV promoter and facilitates optimal production of virus. For more information about the 293FT Cell Line, refer to the 293FT Cell Line manual.

After you have generated the pLenti6/R4R2/V5-DEST expression construct containing your promoter and gene of interest, you will cotransfect the plasmid and the ViraPower™ Packaging Mix into 293FT cells to produce a replication-incompetent lentiviral stock. This lentiviral stock may then be transduced into the mammalian cell line of interest to express your recombinant protein.

How Lentivirus Works

Once the lentivirus enters the target cell, the viral RNA is reverse-transcribed, actively imported into the nucleus (Lewis & Emerman, 1994; Naldini, 1999), and stably integrated into the host genome (Buchschacher & Wong-Staal, 2000; Luciw, 1996). After the lentiviral construct has integrated into the genome, you may assay for transient expression of your recombinant protein or use antibiotic selection to generate a stable cell line for long-term expression studies.

continued on next page

The ViraPower™ Promoterless Lentiviral Gateway® Expression System, continued

VSV Envelope Glycoprotein

Most retroviral vectors are limited in their usefulness as gene delivery vehicles by their restricted tropism and generally low titers. In the ViraPower™ Promoterless Lentiviral Gateway® Expression System, this limitation has been overcome by use of the G glycoprotein gene from Vesicular Stomatitis Virus (VSV-G) as a pseudotyping envelope, thus allowing production of a high titer lentivirus with a significantly broadened host cell range (Burns *et al.*, 1993; Emi *et al.*, 1991; Yee *et al.*, 1994).

In vivo Gene Delivery

The ViraPower™ Promoterless Lentiviral Expression System is suitable for *in vivo* gene delivery applications. Many groups have successfully used lentiviral vectors to express a target gene in tissues including brain, retina, pancreas, muscle, liver, and skin (Gallichan *et al.*, 1998; Kafri *et al.*, 1997; Miyoshi *et al.*, 1997; Naldini, 1998; Pfeifer *et al.*, 2001; Pfeifer *et al.*, 2001; Takahashi *et al.*, 1999). For more information about target genes that have been successfully expressed *in vivo* using lentiviral-based vectors, refer to the references above as well as the following additional references (Baek *et al.*, 2001; Dull *et al.*, 1998; Lois *et al.*, 2002; Park & Kay, 2001; Peng *et al.*, 2001).

Features of the pLenti6/R4R2/V5-DEST Vector

The pLenti6/R4R2/V5-DEST vector contains the following elements:

- Rous Sarcoma Virus (RSV) enhancer/promoter for Tat-independent production of viral mRNA in the producer cell line (Dull *et al.*, 1998)
 - Modified HIV-1 5' and 3' Long Terminal Repeats (LTR) for viral packaging and reverse transcription of the viral mRNA (Dull *et al.*, 1998; Luciw, 1996)
Note: The U3 region of the 3' LTR is deleted (Δ U3) and facilitates self-inactivation of the 5' LTR after transduction to enhance the biosafety of the vector (Dull *et al.*, 1998)
 - HIV-1 psi (Ψ) packaging sequence for viral packaging (Luciw, 1996)
 - HIV Rev response element (RRE) for Rev-dependent nuclear export of unspliced viral mRNA (Kjems *et al.*, 1991; Malim *et al.*, 1989)
 - Two recombination sites, *attR4* and *attR2* for recombinational cloning of the promoter and gene of interest from two separate entry clones
 - The *ccdB* gene located between the *attR* sites for negative selection
 - Chloramphenicol resistance gene (Cm^R) located between the two *attR* sites for counterselection
 - C-terminal V5 epitope for detection of the recombinant protein of interest (Southern *et al.*, 1991)
 - Blastocidin resistance gene for selection in *E. coli* and mammalian cells (Izumi *et al.*, 1991; Kimura *et al.*, 1994; Takeuchi *et al.*, 1958; Yamaguchi *et al.*, 1965)
 - Ampicillin resistance gene for selection in *E. coli*
 - pUC origin for high-copy replication of the plasmid in *E. coli*
-

Biosafety Features of the System

Introduction

The lentiviral and packaging vectors supplied in the ViraPower™ Promoterless Lentiviral Gateway® Expression System are third-generation vectors based on lentiviral vectors developed by Dull *et al.*, 1998. This third-generation HIV-1-based lentiviral system includes a significant number of safety features designed to enhance its biosafety and to minimize its relation to the wild-type, human HIV-1 virus. These safety features are described below.

Biosafety Features of the ViraPower™ Promoterless Lentiviral System

The ViraPower™ Promoterless Lentiviral Gateway® Expression System includes the following key safety features:

- The pLenti6/R4R2/V5-DEST vector contains a deletion in the 3' LTR (Δ U3) that does not affect generation of the viral genome in the producer cell line, but results in "self-inactivation" of the lentivirus after transduction of the target cell (Yee *et al.*, 1987; Yu *et al.*, 1986; Zufferey *et al.*, 1998). Once integrated into the transduced target cell, the lentiviral genome is no longer capable of producing packageable viral genome.
- The number of genes from HIV-1 that are used in the system has been reduced to three (*i.e.* *gag*, *pol*, and *rev*).
- The VSV-G gene from Vesicular Stomatitis Virus is used in place of the HIV-1 envelope (Burns *et al.*, 1993; Emi *et al.*, 1991; Yee *et al.*, 1994).
- Genes encoding the structural and other components required for packaging the viral genome are separated onto four plasmids (*i.e.* three packaging plasmids and pLenti6/R4R2/V5-DEST). All four plasmids have been engineered not to contain any regions of homology with each other to prevent undesirable recombination events that could lead to the generation of a replication-competent virus (Dull *et al.*, 1998).
- Although the three packaging plasmids allow expression *in trans* of proteins required to produce viral progeny (*e.g.* *gal*, *pol*, *rev*, *env*) in the 293FT producer cell line, none of them contain LTRs or the Ψ packaging sequence. This means that none of the HIV-1 structural genes are actually present in the packaged viral genome, and thus, are never expressed in the transduced target cell. No new replication-competent virus can be produced.
- The lentiviral particles produced in this system are replication-incompetent and only carry the gene of interest. No other viral species are produced.
- Expression of the *gag* and *pol* genes from pLP1 has been rendered Rev-dependent by virtue of the HIV-1 RRE in the *gag/pol* mRNA transcript. Addition of the RRE prevents *gag* and *pol* expression in the absence of Rev (Dull *et al.*, 1998).
- A constitutive promoter (RSV promoter) has been placed upstream of the 5' LTR in the pLenti6/R4R2/V5-DEST vector to offset the requirement for Tat in the efficient production of viral RNA (Dull *et al.*, 1998).

continued on next page

Biosafety Features of the System, continued

Biosafety Level 2



Despite the inclusion of the safety features discussed on the previous page, the lentivirus produced with this System can still pose some biohazardous risk since it can transduce primary human cells. For this reason, **we highly recommend that you treat lentiviral stocks generated using this System as Biosafety Level 2 (BL-2) organisms and strictly follow all published BL-2 guidelines with proper waste decontamination.** Furthermore, exercise extra caution when creating lentivirus carrying potential harmful or toxic genes (e.g. activated oncogenes).

For more information about the BL-2 guidelines and lentivirus handling, refer to the document, "Biosafety in Microbiological and Biomedical Laboratories", 4th Edition, published by the Centers for Disease Control (CDC). This document may be downloaded at the following address:

<http://www.cdc.gov/od/ohs/biosfty/bmbl4/bmbl4toc.htm>



Important

Handle all lentiviruses in compliance with established institutional guidelines. Since safety requirements for use and handling of lentiviruses may vary at individual institutions, we recommend consulting the health and safety guidelines and/or safety officer(s) at your institution prior to use of the ViraPower™ Promoterless Lentiviral Gateway® Expression System.

Cell Biology

| | | | | |
|------------------|--|--|--------------------|-------------------|
| ATCC® Number: | CRL-11268™ | <input type="button" value="Order this Item"/> | Price: | \$264.00 |
| Designations: | 293T/17 [HEK 293T/17] | | Depositors: | Rockefeller Univ. |
| Biosafety Level: | 2 [Cells contain Adeno and SV-40 viral DNA sequences] | | Shipped: | frozen |
| Medium & Serum: | See Propagation | | Growth Properties: | adherent |
| Organism: | <i>Homo sapiens</i> (human) | | Morphology: | epithelial |
| Source: | Organ: kidney | | | |
| Permits/Forms: | In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location. | | | |

This material is cited in a U.S. and/or other Patent or Patent Application, and may not be used to infringe on the patent claims. ATCC is required to inform the Patent Depositor of the party to which the material was furnished.

Related Cell Culture Products

| | |
|---------------------|--|
| Restrictions: | The line is available with the following restriction: 1. The cell line was deposited at the ATCC by Rockefeller University and is provided for research purposes only. Neither the cell line nor the products derived from it may be sold or used for commercial purposes. Nor can the cells be distributed to third parties for purposes of sale, or producing for sale, cells or their products. The cells are provided as a service to the research community. They are provided without warranty of merchantability or fitness for a particular purpose or any other warranty, expressed or implied. 2. Any proposed commercial use of the cells, or their products, must first be negotiated with Cell Genesys, 500 Forbes Boulevard, South San Francisco, CA 94080 Attn: Robert H. Tidwell; Senior Vice President, Corporate Development. |
| Antigen Expression: | SV40 T antigen [45408] |
| Age: | fetus |
| Comments: | The 293T/17 cell line is a derivative of the 293T (293tsA1609neo) cell line. 293T is a highly transfectable derivative of the 293 cell line into which the temperature sensitive gene for SV40 T-antigen was inserted. 293T cells were cloned and the clones tested with the pBND and pZAP vectors to obtain a line capable of producing high titers of infectious retrovirus, 293T/17. These cells constitutively express the simian virus 40 (SV40) large T antigen, and clone 17 was selected specifically for its high transfectability. 293T/17 cells were cotransfected with the pCRIPenv- and the pCRIPgag-2 vectors to obtain the ANJOU 65 (see ATCC CRL-11269) cell line. ANJOU 65 cells were cotransfected with the pCRIPgag-2 and pGPT2E vectors to obtain the BOSC 23 (see ATCC CRL-11270) ecotropic envelope-expression packaging cell line. ANJOU 65 cells were also cotransfected with the pCRIPAMgag vector along with a plasmid expressing the gpt resistance gene to obtain the Bing (see ATCC CRL-11554) amphotropic envelope-expression packaging cell line. |

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.

Temperature: 37.0°C

Protocol:

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin- 0.53 mM EDTA solution to remove all traces of serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).

Subculturing: Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.

4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessels.
6. Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:4 to 1:8 is recommended

Medium Renewal: Every 2 to 3 days

Preservation: **Freeze medium:** Complete growth medium supplemented with 5% (v/v) DMSO
Storage temperature: liquid nitrogen vapor phase

derivative:ATCC CRL-11269

Related Products: recommended serum:ATCC 30-2020

Recommended medium (without the additional supplements or serum described under ATCC Medium):ATCC 30-2002

45408: Sena-Esteves M, et al. Single-step conversion of cells to retrovirus vector producers with herpes simplex virus-Epstein-Barr virus hybrid amplicons. *J. Virol.* 73: 10426-10439, 1999. PubMed: 10559361

References: 57446: Pensiero M, et al. Retroviral vectors produced by producer cell lines resistant to lysis by human serum. US Patent 5,952,225 dated Sep 14 1999

57447: Pensiero M, et al. Retroviral vectors produced by producer cell lines resistant to lysis by human serum. US Patent 6,329,199 dated Dec 11 2001

57448: Pear WS, et al. Production of High-Titer Helper-Free Retroviruses by Transient Transfection. *Proc. Natl. Acad. Sci. USA* 90: 8392-8396, 1993.

PubMed: 7690960

[Return to Top](#)



Your Discoveries
Begin with Us.™

Search Catalog:

[Home](#) | [Ordering Info](#) | [Quick Order](#) | [Cart](#) | [Te](#)

Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's [Material Transfer Agreement](#) or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Japan, Hong Kong, Korea, New Zealand, Singapore and Taiwan, R.O.C. must contact a local distributor for pricing information and to place an order for ATCC cultures and products.

| Cell Lines | |
|--|---|
| ATCC® Number: CCL-107™ | Order this item Price: \$185.00 |
| Designations: C6 | Depositors: G Sato |
| Biosafety Level: 1 | Shipped: frozen |
| Medium & Serum: See Propagation | Growth Properties: adherent |
| Organism: <i>Rattus norvegicus</i> (rat) | Morphology: fibroblast |
| Source: Organ: brain Cell type: glial cell Disease: glioma | |
| Cellular Products: S-100 protein; produce glyceryl phosphate dehydrogenase in response to glucocorticoids; somatotrophin | |
| Permits/Forms: In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location. | |
| Related Cell Culture Products | |
| Receptors: | glucocorticoid |
| Virus Susceptibility: | vesicular stomatitis (Indiana); vaccinia; herpes simplex |
| Virus Resistance: | poliovirus 3 |
| Reverse Transcript: | negative |
| Cytogenetic Analysis: | Stemline number is diploid. Karyotype is stable within the stemline number and is that of a normal male. Three cells with breaks; one with a secondary constriction, one with a dicentric, one with a rearrangement and four with terminal or centromere associations. |
| Comments: | The glial cell strain, C6, was cloned from a rat glial tumor induced by N-nitrosomethylurea by Benda et al. after a series of alternate culture and animal passages [PubMed: 4873531]. S-100 production increases ten fold as cells grow from low density to confluency. |
| Propagation: | ATCC complete growth medium: Ham's F12K medium with 2 mM L-glutamine adjusted to contain 1.5 g/L sodium bicarbonate, 82.5%; horse serum, 15%; fetal |

| | |
|--------------------------|---|
| | bovine serum, 2.5% Temperature: 37.0C Atmosphere: air, 95%; carbon dioxide (CO ₂), 5% |
| Subculturing: | <p>Protocol:</p> <ol style="list-style-type: none"> 1. Remove and discard culture medium. 2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin - 0.53 mM EDTA solution to remove all traces of serum which contains trypsin inhibitor. 3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37C to facilitate dispersal. 4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting. 5. Add appropriate aliquots of the cell suspension to new culture vessels. 6. Incubate cultures at 37C. <p>Subcultivation ratio: A subcultivation ratio of 1:2 to 1:3 is recommended</p> <p>Medium renewal: 2 to 3 times per week</p> |
| Preservation: | Freeze medium: culture medium, 95%; DMSO, 5% Storage temperature: liquid nitrogen vapor phase |
| Related Products: | Recommended medium (without the additional supplements or serum described under ATCC Medium): ATCC 30-2004 recommended serum: ATCC 30-2020 recommended serum: ATCC 30-2040 0.25% (w/v) Trypsin - 0.53 mM EDTA in Hank' BSS (w/o Ca ⁺⁺ , Mg ⁺⁺): ATCC 30-2101 Cell culture tested DMSO: ATCC 4-X |
| References: | 1022: Benda P , et al. Differentiated rat glial cell strain in tissue culture. Science 161: 370-371, 1968. PubMed: 4873531 25965: Lightbody JJ , et al. Establishment of differentiated clonal strains of glial brain cells in culture. Fed. Proc. 27: 720, 1968. 32720: Chen Y , et al. Demonstration of binding of dengue virus envelope protein to target cells. J. Virol. 70: 8765-8772, 1996. PubMed: 8971005 |

Notices and Disclaimers

ATCC products are intended for laboratory research purposes only. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States and Canada. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your word search in Cell Biology selections by typing in your search criteria below or returning to the [Cell Biology](#) menu. To search another product line, choose one from the dropdown box at the top. For complex searches using boolean operators, the following characters must be used: & (for AND), | (for OR), ^ (for AND NOT). An asterisk (*) is used as the wildcard. For more information please review the [Search Help](#).

C6 glioma

Word Search

Clear Search

Home Page Archive

[Home](#) [Ordering Info](#) [Quick Order](#) [Support](#) [About ATCC](#) [Contact Us](#)
[Privacy Policy](#) [Terms of Use](#) [ATCC MTA](#)

© 2005 American Type Culture Collection (ATCC).
All rights reserved.



Your Discoveries
Begin with US.™

Search: Choose Op

[Home](#) || [Ordering Info](#) || [Quick Order](#) || [Cart](#) || [Tech Support](#)

Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's Material Transfer Agreement or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Japan, Hong Kong, Korea, New Zealand, Singapore and Taiwan, R.O.C. must contact a local distributor for pricing information and to place an order for ATCC cultures and products.

| Cell Lines | |
|--|--|
| ATCC® Number: CCL-229™ | Order this item Price: \$185.00 |
| Designations: LoVo | Depositors: M Romsdahl |
| Biosafety Level: 1 | Shipped: frozen |
| Medium & Serum: See Propagation | Growth Properties: adherent |
| Organism: <i>Homo sapiens</i> (human) | Morphology: epithelial |
| Source: Organ: colon Disease: colorectal adenocarcinoma Tumor stage: Dukes' type C, grade IV Derived from metastatic site: left supraclavicular region | |
| Cellular Products: carcinoembryonic antigen (CEA) 908 ng/10 exp6 cells/10 days | |
| Permits/Forms: In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location. | |
| Related Cell Culture Products | |
| Tumorigenic: | Yes, in nude mice (Tumors developed within 21 days at 100% frequency (5/5) in nude mice inoculated subcutaneously with 10(7) cells) |
| Reverse Transcript: | negative |
| Oncogene: | myc +; myb +; ras +; fos +; p53 +; sis -; abl -; ros -; src - |
| Antigen Expression: | HLA A11, B15, B17, Cw1, Cw3; blood type B |
| Cytogenetic Analysis: | The stemline chromosome number is hyperdiploid with the 2S component occurring at about 2.7% and 3 marker chromosomes were common to all S metaphases. Karyotypes were generally homogeneous and stable. |
| Isoenzymes: | ES-D, 1; G6PD, B; PGD, A; PGM1, 2; PGM3, 1-2 |
| Age: | 56 years |
| Gender: | male |
| Comments: | LoVo was initiated in 1971 from a fragment of a metastatic tumor nodule in the left |

| | |
|--------------------------|--|
| | <p>supraclavicular region of a 56-year-old Caucasian male patient with a histologically proven diagnosis of adenocarcinoma of the colon. [1049] The cells are negative for expression of CSAp (CSAp-) and colon antigen 3. The line is positive for expression of c-myc, K-ras, H-ras, N-ras, Myb, sis and fos oncogenes. [22861] Myb, and fos oncogenes. [22861] N-myc and sis oncogene expression were not detected. [22861] Tumor specific nuclear matrix proteins CC-3 and CC-4 are expressed. [23341]</p> |
| Propagation: | <p>ATCC complete growth medium: Ham's F12K medium with 2 mM L-glutamine adjusted to contain 1.5 g/L sodium bicarbonate, 90%; fetal bovine serum, 10% Temperature: 37.0C</p> |
| Subculturing: | <p>Remove medium, and rinse with 0.25% trypsin, 0.03% EDTA solution. Remove the solution and add an additional 1 to 2 ml of trypsin-EDTA solution. Allow the flask to sit at room temperature (or at 37C) until the cells detach. Add fresh culture medium, aspirate and dispense into new culture flasks. Subcultivation ratio: A subcultivation ratio of 1:3 to 1:10 is recommended Medium renewal: 2 to 3 times per week</p> |
| Preservation: | <p>culture medium 95%; DMSO, 5%</p> |
| Related Products: | <p>Recommended medium (without the additional supplements or serum described under ATCC Medium): ATCC 30-2004 recommended serum: ATCC 30-2020</p> |
| References: | <p><u>1047</u>: Drewinko B , et al. Further biologic characteristics of a human carcinoembryonic antigen-producing colon carcinoma cell line. J. Natl. Cancer Inst. 61: 75-83, 1978. PubMed: <u>276641</u> <u>1048</u>: Drewinko B , Yand LY . Restriction of CEA synthesis to the stationary phase of growth of cultured human colon carcinoma cells. Exp. Cell Res. 101: 414-416, 1976. PubMed: 964319 <u>1049</u>: Drewinko B , et al. Establishment of a human carcinoembryonic antigen-producing colon adenocarcinoma cell line. Cancer Res. 36: 467-475, 1976. PubMed: 1260746 <u>22861</u>: Trainer DL , et al. Biological characterization and oncogene expression in human colorectal carcinoma cell lines. Int. J. Cancer 41: 287-296, 1988. PubMed: <u>3338874</u> <u>23341</u>: Keesee SK , et al. Nuclear matrix proteins in human colon cancer. Proc. Natl. Acad. Sci. USA 91: 1913-1916, 1994. PubMed: 8127905 <u>26057</u>: Drewinko B , et al. Response of exponentially growing, stationary-phase, and synchronized cultured human colon carcinoma cells to treatment with nitrosourea derivatives. Cancer Res. 39: 2630-2636, 1979. PubMed: 445465 <u>32913</u>: Miranda L , et al. Isolation of the human PC6 gene encoding the putative host protease for HIV-1 gp160 processing in CD4+ T lymphocytes. Proc. Natl. Acad. Sci. USA 93: 7695-7700, 1996. PubMed: 8755538</p> |

Notices and Disclaimers

ATCC products are intended for laboratory research purposes only. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States and Canada. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your word search in Cell Biology selections by typing in your search criteria below or returning to the Cell Biology menu. To search another product line, choose one from the dropdown box at the top. For complex searches using boolean operators, the following characters must be used: & (for

AND), | (for OR), ^ (for AND NOT). An asterisk (*) is used as the wildcard. For more information please review the [Search Help](#).

Home Page Archive

[Home](#) [Ordering Info](#) [Quick Order](#) [Support](#) [About ATCC](#) [Contact Us](#)
[Privacy Policy](#) [Terms of Use](#) [ATCC MTA](#)

© 2005 American Type Culture Collection (ATCC).
All rights reserved.



Your Discoveries
Begin with Us.

Search: --- Choose Op

[Home](#) || [Ordering Info](#) || [Quick Order](#) || [Cart](#) || [Tech Support](#)

Search Results

The search engine found 1 entries for **c2c12**.

Entries 1 to 1 are currently being displayed. [Continue Search](#).

[View Selections](#) | [View All](#) | [Clear Selections](#)

Cell Biology

| ATCC® Number | Description | Designation | View |
|--------------------------|-----------------------------|-------------|------|
| CRL-1772 | <i>Mus musculus</i> (mouse) | C2C12 | |

[View Selections](#) | [View All](#) | [Clear Selections](#)

You may continue your word search in Cell Biology selections by typing in your search criteria below or returning to the [Cell Biology](#) menu. To search another product line, choose one from the dropdown box at the top. For complex searches using boolean operators, the following characters must be used: & (for AND), | (for OR), ^ (for AND NOT). An asterisk (*) is used as the wildcard. For more information please review the [Search Help](#).

[Word Search](#)

[Clear Search](#)

Home Page Archive

[Home](#) [Ordering Info](#) [Quick Order](#) [Support](#) [About ATCC](#) [Contact Us](#)
[Privacy Policy](#) [Terms of Use](#) [ATCC MTA](#)

© 2005 American Type Culture Collection (ATCC).
All rights reserved.



Your Discoveries
Begin with Us.™

Search: --- Choose Op

[Home](#) || [Ordering Info](#) || [Quick Order](#) || [Cart](#) || [Tech Support](#)

Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's [Material Transfer Agreement](#) or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Japan, Hong Kong, Korea, New Zealand, Singapore and Taiwan, R.O.C. must contact a [local distributor](#) for pricing information and to place an order for ATCC cultures and products.

| Cell Lines | |
|--|--|
| ATCC® Number: CRL-1772™ | Order this item Price: \$185.00 |
| Designations: C2C12 | Depositors: B Paterson |
| Biosafety Level: 1 | Shipped: frozen |
| Medium & Serum: See Propagation | Growth Properties: adherent |
| Organism: <i>Mus musculus</i> (mouse) | Morphology: fibroblast  |
| Source: Tissue: muscle Cell type: myoblast; myoblast | |
| Permits/Forms: | In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location. |
| Related Cell Culture Products | |
| Strain: | C3H |
| Comments: | This is a subclone (produced by H. Blau, et al) of the mouse myoblast cell line established by D. Yaffe and O. Saxel. [22903] The C2C12 cell line differentiates rapidly, forming contractile myotubes and producing characteristic muscle proteins. [22953] Treatment with bone morphogenic protein 2 (BMP-2) cause a shift in the differentiation pathway from myoblastic to osteoblastic. [23427] Tested and found negative for ectromelia virus (mousepox). |
| Propagation: | ATCC complete growth medium: Dulbecco's modified Eagle's medium with 4 mM L-glutamine adjusted to contain 1.5 g/L sodium bicarbonate and 4.5 g/L glucose, 90%; fetal bovine serum, 10% Temperature: 37.0C |
| Subculturing: | Protocol: IMPORTANT - DO NOT ALLOW CULTURES TO BECOME CONFLUENT. Cultures must not be allowed to become confluent as this will deplete the myoblastic population in the culture. Myotube formation is enhanced when the medium is supplemented with 10% horse serum instead of fetal bovine serum. |

| | |
|--------------------------|---|
| | <ol style="list-style-type: none"> 1. Remove and discard culture medium. 2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin- 0.53 mM EDTA solution to remove all traces of serum which contains trypsin inhibitor. 3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes). Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal. 4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting. 5. Add appropriate aliquots of the cell suspension to new culture vessels. Inoculate at a cell concentration between 1.5×10^5 and 1.0×10^6 viable cells/75 cm². 6. Incubate cultures at 37°C. <p>Medium renewal: Every two to three days</p> |
| Preservation: | Freeze medium: Complete growth medium supplemented with 5% (v/v) DMSO Storage temperature: liquid nitrogen vapor phase |
| Related Products: | Recommended medium (without the additional supplements or serum described under ATCC Medium): ATCC 30-2002 recommended serum: ATCC 30-2020 |
| References: | <p>22903: Yaffe D , Saxel O . Serial passaging and differentiation of myogenic cells isolated from dystrophic mouse muscle. Nature 270: 725-727, 1977. PubMed: 563524</p> <p>22953: Blau HM , et al. Plasticity of the differentiated state. Science 230: 758-766, 1985. PubMed: 2414846</p> <p>23427: Katagiri T , et al. Bone morphogenetic protein-2 converts the differentiation pathway of C2C12 myoblasts into the osteoblast lineage [published erratum appears in J Cell Biol 1995 Feb;128(4):following 713]. J. Cell Biol. 127: 1755-1766, 1994. PubMed: 7798324</p> <p>28236: Chow YH , et al. Improvement of hepatitis B virus DNA vaccines by plasmids coexpressing hepatitis B surface antigen and interleukin-2. J. Virol. 71: 169-178, 1997. PubMed: 8985336</p> <p>32828: Kessler PD , et al. Gene delivery to skeletal muscle results in sustained expression and systemic delivery of a therapeutic protein. Proc. Natl. Acad. Sci. USA 93: 14082-14087, 1996. PubMed: 8943064</p> <p>33069: Hsu DK , et al. Identification of a murine TEF-1-related gene expressed after mitogenic stimulation of quiescent fibroblasts and during myogenic differentiation. J. Biol. Chem. 271: 13786-13795, 1996. PubMed: 8662936</p> |

Notices and Disclaimers

ATCC products are intended for laboratory research purposes only. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States and Canada. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your word search in Cell Biology selections by typing in your search criteria below or returning to the [Cell Biology](#) menu. To search another product line, choose one from the dropdown box at the top. For complex searches using boolean operators, the following characters must be used: & (for AND), | (for OR), ^ (for AND NOT). An asterisk (*) is used as the wildcard. For more information please review the [Search Help](#).

c2c12

Word Search

Clear Search

Home Page Archive

[Home](#) [Ordering Info](#) [Quick Order](#) [Support](#) [About ATCC](#) [Contact Us](#)
[Privacy Policy](#) [Terms of Use](#) [ATCC MTA](#)

© 2005 American Type Culture Collection (ATCC).
All rights reserved.

Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's Material Transfer Agreement or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Canada, China, Hong Kong, India, Japan, Korea, Macau, Mexico, New Zealand, Singapore, and Taiwan R.O.C. must contact a local distributor for pricing information and to place an order for ATCC cultures and products.

[Print this](#)

Cell Biology

| | | | | |
|----------------------------|---------------------------------|---------------------------------|---------------------------|---------------------|
| ATCC® Number: | CRL-5803™ | Order this Item | Price: | \$264.00 |
| Designations: | NCI-H1299 | | Depositors: | AF Gazdar, JD Minna |
| Biosafety Level: | 1 | | Shipped: | frozen |
| Medium & Serum: | See Propagation | | Growth Properties: | adherent |
| Organism: | <i>Homo sapiens</i> (human) | | Morphology: | epithelial |

Source: **Organ:** lung
Disease: carcinoma; non-small cell lung cancer
Derived from metastatic site: lymph node

Cellular Products: neuromedin B

Permits/Forms: In addition to the [MTA](#) mentioned above, other [ATCC and/or regulatory permits](#) may be required for the transport of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location.

Related Cell Culture Products

Restrictions: The line is available with the following restrictions: 1. This cell line was deposited at the ATCC by Dr. A. Gazdar and Dr. J. Minna and is provided for research purposes only. Neither the cell line nor products derived from it may be sold or used for commercial purposes. Nor can the cells be distributed to third parties for purposes of sale, or producing for sale, cells or their products. The cells are provided as service to the research community. They are provided without warranty of merchantability or fitness for a particular purpose or any other warranty expressed or implied. 2. Any proposed commercial use of the these cells, or their products must first be negotiated with the University of Texas Southwestern Medical Center at Dallas, 5323 Harry Hines Blvd., Dallas, Texas 75235. Telephone (214) 699-8056, FAX (214) 688-7233.

Applications: transfection host (technology from amaxa [Roche FuGENE® Transfection Reagents](#))

Age: 43 years adult

Gender: male

Ethnicity: Caucasian

Comments: The cells have a homozygous partial deletion of the p53 protein, and lack expression of p53 protein. They are reported to be able to synthesize the peptide neuromedin B (NMB) at 0.1 pmol/mg protein, but not the growth releasing peptide (GRP).

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated RPMI-1640 Medium Catalog No. 30-2001. To make the complete growth medium, add the following components to the medium: fetal bovine serum to a final concentration of 10%.

Temperature: 37.0°C

Subculturing: **Protocol:**

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin - 0.53 mM EDTA solution to remove all trace serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope. Cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37C to facilitate dispersal.
4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessels.
6. Incubate cultures at 37C.

Subcultivation Ratio: A subcultivation ratio of 1:3 to 1:6 is recommended
Medium Renewal: Every 2 to 3 days

- Preservation:** **Freeze medium:** Complete growth medium supplemented with 5% (v/v) DMSO
Storage temperature: liquid nitrogen vapor phase
- Related Products:** recommended serum: ATCC 30-2020
purified DNA: ATCC CRL-5803D
Recommended medium (without the additional supplements or serum described under ATCC Medium): ATCC 2001
- References:** 23517: Giaccone G, et al. Neuromedin B is present in lung cancer cell lines. Cancer Res. 52: 2732s-2735s, 1992. PubMed: 1563005
23570: . NCI-Navy Medical Oncology Branch Cell Line Supplement. J. Cell. Biochem. suppl. 24: 1996.
33177: Lin DL, Chang C. p53 is a mediator for radiation-repressed human TR2 orphan receptor expression in MCF-7 cells, a new pathway from tumor suppressor to member of the steroid receptor superfamily. J. Cell. Physiol. 167: 14649-14652, 1996. PubMed: 8663350

[Return to top](#)

Notices and Disclaimers

ATCC products are intended for laboratory research purposes only, unless noted otherwise. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States. Cultures that are ordered as test tubes or flasks will carry an additional labor fee. Fees for permits, shipping, and handling may apply.

[Back to my Search](#)

Customize your ATCC Web experience: [Login](#) >

[Home](#) | [Site Map](#) | [FAQ](#) | [Privacy Policy](#) | [Careers](#) | [Contact Us](#)

© 2008 ATCC. All Rights Reserved

Product Description

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's Material Transfer Agreement or, in certain cases, an MTA specified by the depositing institution.

Customers in Europe, Australia, Canada, China, Hong Kong, India, Japan, Korea, Macau, Mexico, New Zealand, Singapore, and Taiwan, R.O.C. must contact a local distributor for pricing information and to place an order for ATCC cultures and products.

[Print this](#)

Cell Biology

| | | | | |
|---------------------------------|---|---------------------------------|---------------------------|---|
| ATCC[®] Number: | CRL-1435™ | Order this Item | Price: | \$256.00 |
| Designations: | PC-3 | | Depositors: | ME Kaighn |
| Biosafety Level: | 1 | | Shipped: | frozen |
| Medium & Serum: | See Propagation | | Growth Properties: | adherent (The cells form clusters in soft agar and can be adapted to suspension growth) |
| Organism: | <i>Homo sapiens</i> (human) | | Morphology: | epithelial |
| | | | |  <small>PHOTO</small> |
| Source: | Organ: prostate Tumor Stage: grade IV Disease: adenocarcinoma Derived from metastatic site: bone | | | |
| Permits/Forms: | In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transport of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for information regarding the specific requirements for shipment to your location. | | | |
| | | | | Related Cell Culture Products |
| Applications: | transfection host (technology from amaxa Roche FuGENE [®] Transfection Reagents) | | | |
| Tumorigenic: | YES | | | |
| Antigen Expression: | HLA A1, A9 | | | |
| DNA Profile (STR): | Amelogenin: X CSF1PO: 11 D13S317: 11 D16S539: 11 D5S818: 13 D7S820: 8,11 THO1: 6,7 TPOX: 8,9 vWA: 17 | | | |
| Cytogenetic Analysis: | The line is near-triploid with a modal number of 62 chromosomes. There are nearly 20 marker chromosomes commonly found in each cell; and normal N2, N3, N4, N5, N12, and N15 are not found. No normal chromosomes could be detected by Q-band analysis. | | | |
| Age: | 62 years adult | | | |
| Gender: | male | | | |

Ethnicity: Caucasian

Comments: The PC-3 was initiated from a bone metastasis of a grade IV prostatic adenocarcinoma from a 62-year-old Caucasian. [22363]
The cells exhibit low acid phosphatase and testosterone-5-alpha reductase activities.

Propagation: **ATCC complete growth medium:** The base medium for this cell line is ATCC-formulated F-12K Me Catalog No. 30-2004. To make the complete growth medium, add the following components to the medium: fetal bovine serum to a final concentration of 10%.
Atmosphere: air, 95%; carbon dioxide (CO₂), 5%
Temperature: 37.0°C

Subculturing: **Protocol:**

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 0.25% (w/v) Trypsin- 0.53 mM EDTA solution to remove all trace serum that contains trypsin inhibitor.
3. Add 2.0 to 3.0 ml of Trypsin-EDTA solution to flask and observe cells under an inverted microscope cell layer is dispersed (usually within 5 to 15 minutes).
Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 ml of complete growth medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessels.
6. Incubate cultures at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:3 to 1:6 is recommended

Medium Renewal: 2 to 3 times per week

Preservation: **Freeze medium:** Complete growth medium supplemented with 5% (v/v) DMSO

Storage temperature: liquid nitrogen vapor phase

Related Products: Recommended medium (without the additional supplements or serum described under ATCC Medium):ATCC 2004
recommended serum:ATCC [30-2020](#)

References: 22363: Kaighn ME, et al. Establishment and characterization of a human prostatic carcinoma cell line (F-12K). Invest. Urol. 17: 16-23, 1979. PubMed: 447482
22470: Chen TR. Chromosome identity of human prostate cancer cell lines, PC-3 and PPC-1. Cytogenet Genet. 62: 183-184, 1993. PubMed: 8428522
26302: Ohnuki Y, et al. Chromosomal analysis of human prostatic adenocarcinoma cell lines. Cancer Res 524-534, 1980. PubMed: 7471073
32341: Sheng S, et al. Maspin acts at the cell membrane to inhibit invasion and motility of mammary prostatic cancer cells. Proc. Natl. Acad. Sci. USA 93: 11669-11674, 1996. PubMed: 8876194
32344: Umekita Y, et al. Human prostate tumor growth in athymic mice: inhibition by androgens stimulation by finasteride. Proc. Natl. Acad. Sci. USA 93: 11802-11807, 1996. PubMed: 8876218
32460: Carter RE, et al. Prostate-specific membrane antigen is a hydrolase with substrate and pharmacological characteristics of a neuropeptidase. Proc. Natl. Acad. Sci. USA 93: 749-753, 1996. PubMed: 8570628
32486: Nupponen NN, et al. Genetic alterations in prostate cancer cell lines detected by comparative genomic hybridization. Cancer Genet. Cytogenet. 101: 53-57, 1998. PubMed: 9460501
32488: Geiger T, et al. Antitumor activity of a PKC-alpha antisense oligonucleotide in combination with standard chemotherapeutic agents against various human tumors transplanted into nude mice. Anticancer Drug Devel. 35-45, 1998. PubMed: 9474241
32916: Su ZZ, et al. Surface-epitope masking and expression cloning identifies the human prostate carcinoma tumor antigen gene PCTA-1 a member of the galectin gene family. Proc. Natl. Acad. Sci. USA 93: 7252-7256, 1996. PubMed: 8692978

Return to

Notices and Disclaimers

ATCC products are intended for laboratory research purposes only, unless noted otherwise. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States. Cultures that are ordered as test tubes or flasks will carry an additional labor fee. Fees for permits, shipping, and handling may apply.

[Back to my Search](#)

Customize your ATCC Web experience: [Login](#) 

[Home](#) | [Site Map](#) | [FAQ](#) | [Privacy Policy](#) | [Careers](#) | [Contact Us](#)

© 2008 ATCC. All Rights Res