

**Practice Term Test 2**

1. Problems from PS 3.
2. Exercise 7.9. (Note that the integral in definition of function  $F$  here is the Lebesgue integral!)
3. Exercise 7.11.
4. Exercise 7.13.
5. Exercise 7.15.
6. Exercise 8.5.
7. Exercise 8.7.
8. Exercise 11.8.
9. Let  $n \geq 2$  and let  $S$  be a standard  $n$ -simplex in  $\mathbb{R}^n$  with base of length  $a$ , for some  $a > 0$ . That is,

$$S := \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_i \geq 0, \sum_{i=1}^n x_i \leq a\}.$$

Use Fubini Theorem (and induction) to find the Lebesgue integral  $\int_{\mathbb{R}^n} \chi_S$ .

For Problems 10 and 11, let  $(X, \mathcal{M}, \mu)$  be a  $\sigma$ -finite measure space, and let  $f : X \rightarrow \mathbb{R}$  be an  $\mathcal{M}$ -measurable function. Define the *distribution function* of  $f$  by

$$\mu_f(t) := \mu(\{x \in X : |f(x)| \geq t\}), \quad t > 0.$$

10. Show that  $\mu_f : (0, \infty) \rightarrow [0, \mu(X)]$  is non-increasing and Borel measurable.
11. Prove that, for any  $p \in [1, \infty)$ ,

$$\int_X |f(x)|^p d\mu(x) = \int_0^\infty \mu_f(t) p t^{p-1} dt.$$

Hint:  $|f(x)|^p = \int_0^{|f(x)|} p t^{p-1} dt$ .